

Magnetic skyrmion crystal at a topological insulator surface

Stefan Divic^{1,*}, Henry Ling², T. Pereg-Barnea^{3,†}, and Arun Paramekanti^{4,‡}

¹*Department of Physics, University of California, Berkeley, California 94720, USA*

²*208-5800 Cooney Road, Richmond, British Columbia V6X3A8, Canada*

³*Department of Physics and the Centre for the Physics of Materials, McGill University, Montréal, Québec H3A 2T8, Canada*

⁴*Department of Physics, University of Toronto, Toronto, Ontario M5S 1A7, Canada*



(Received 10 April 2021; revised 5 November 2021; accepted 5 January 2022; published 31 January 2022; corrected 7 February 2022)

We consider a magnetic skyrmion crystal formed at the surface of a topological insulator. Incorporating the exchange interaction between the helical Dirac surface states and the periodic Néel or Bloch skyrmion texture, we obtain the resulting electronic band structure and discuss the constraints that symmetries impose on the energies and Berry curvature. We find substantive qualitative differences between the Néel and Bloch cases, **with the latter generically permitting a multiband low energy tight-binding representation whose parameters are tightly constrained by symmetries.** We explicitly compute the associated Wannier orbitals, which resemble the ringlike chiral bound states of helical Dirac fermions coupled to a single skyrmion in a ferromagnetic background. We construct a two-band tight-binding model with real nearest-neighbor hoppings which captures the salient topological features of the low-energy bands. Our results are relevant to magnetic topological insulators (TIs), as well as to TI-magnetic thin film heterostructures, in which skyrmion crystals may be stabilized.

DOI: [10.1103/PhysRevB.105.035156](https://doi.org/10.1103/PhysRevB.105.035156)

I. INTRODUCTION

Massless Dirac fermions emerge in condensed matter as low energy excitations of systems whose Fermi levels sit close to a band crossing. Notable examples of this phenomenon in dimensions $d > 1$ include graphene, Weyl/Dirac semimetals, and the surface states of strong topological insulators (TIs) [1–7]. In certain cases, such band touchings may be protected by lattice or time-reversal symmetries, so that breaking these symmetries induces a Dirac mass gap, leading to physically observable consequences [8–12]. For instance, inducing a mass gap in graphene by breaking inversion symmetry leads to a valley Hall effect [13], while breaking time-reversal symmetry for a TI surface Dirac cone via a perpendicular Zeeman field leads to a gapped half-integer quantum Hall state [14] with $\sigma_{xy} = e^2/2h$. Such symmetry breaking may be induced by proximity coupling with a symmetry-broken substrate or by spontaneous ordering of magnetic moments. Domain walls of such broken symmetries, where the Dirac mass changes sign, act as channels which support chiral edge modes [15–17].

Going beyond the impact of uniform symmetry breaking orders and isolated domain walls, it is interesting to consider the effect of periodically modulated potentials on massless Dirac fermions. Such modulations have been extensively studied in the context of superlattices in graphene [18–23] and bilayer graphene [24,25], where they have been shown to produce emergent Dirac fermion excitations. The superlattice reconstruction of low-energy bands has also recently come to

the fore in studies of twisted bilayer graphene [26–40] and multilayer transition metal dichalcogenides [41–48], where the moiré pattern leads to an enlarged unit cell, as well as in recent work examining moiré potential modulations on TI surface states [49,50].

In this paper, inspired by these previous developments, we study magnetic skyrmion lattices on a TI surface and explore the resulting electronic states. Our work is also motivated by the desire to understand the interplay of the momentum space topology of TIs, as reflected in their helical Dirac surface states, with the topological real space texture of magnetic skyrmions. For instance, materials such as topological Kondo insulators (TKIs) can have Dirac surface states together with soft magnetic modes in the bulk due to strong correlation effects [51]. Such materials might thus be prone to spontaneous magnetic ordering and time-reversal breaking at the surface [52–55]. The inversion breaking at the TKI surface can also enhance the role of chiral Dzyaloshinskii-Moriya magnetic exchange interactions [56,57], which could favor the formation of skyrmions at the surface. Magnetic topological materials such as MnBi_2Te_4 [58] are another proposed candidate for realizing skyrmions [59]. Further possibilities of realizing magnetic skyrmions at TI surfaces include the ordering of impurity magnetic moments of dopants induced by RKKY interactions [60], proximity coupling to a magnetic substrate hosting these textures [61–65], or spontaneous magnetic ordering due to hexagonal warping of the surface Dirac cone [66].

Previous theoretical work demonstrated the electrical charging of nonuniform magnetic textures, such as vortices and domain walls of Néel and Bloch type, by coupling to Dirac TI surface states [67]. Focusing on an isolated Bloch skyrmion texture, it was subsequently shown that chiral bound states, confined to the skyrmion perimeter and analogous to

*stefan_divic@berkeley.edu

†tamipb@physics.mcgill.ca

‡arun.paramekanti@utoronto.ca

the Jackiw-Rebbi zero mode, offer a complementary mechanism for inducing electric charge [15,68]. Later studies investigated the scattering of Dirac electrons off of single skyrmions [69,70] and showed that the in-gap bound states modify the skyrmion-skyrmion interaction potential [71].

In ordinary magnetic metals, the presence of skyrmions is often inferred from an additional Hall contribution generated by the real-space Berry curvature induced by skyrmions. Termed the topological Hall effect (THE) [72–77], this transport phenomenon is distinct from the anomalous Hall effect that derives from momentum-space Berry curvature [63]. Recent experiments on magnetically doped TI superlattices have interpreted anomalies in the Kerr effect [78] as arising from skyrmions. However, we emphasize that the experimental identification and disentanglement of the skyrmion contribution from magnetic inhomogeneities can be difficult in practice, as showcased by recent work on SrRuO₃ films [79–81]. More importantly, as we discuss below, massless Dirac fermions moving in a skyrmion background do not sense the topological charge of the skyrmions as a magnetic flux and should not exhibit the THE.

This paper is organized as follows. In Sec. II, we begin with a band theoretic analysis of Dirac fermions coupled to the periodic Zeeman texture of the skyrmion lattice. This ‘nearly-free Dirac electron’ approach allows us to determine the energy bands and their topological invariants, as well as symmetries. In Sec. III B, we pass to a tight-binding description of the low-energy Bloch skyrmion bands, drawing inspiration from the bound states of the single-skyrmion problem when constructing Wannier orbitals. We conclude with a summary of important results, possible limitations, and a discussion of promising future directions.

II. CONTINUUM BAND THEORY

A. Single skyrmion and skyrmion crystal ansatz

The unit vector magnetization of an isolated 2D skyrmion centered at the origin may be written in the following form:

$$\mathbf{n}(r, \phi) = \begin{pmatrix} \sqrt{1 - n_z(r)^2} \cos(\phi + \phi_0) \\ \sqrt{1 - n_z(r)^2} \sin(\phi + \phi_0) \\ n_z(r) \end{pmatrix}. \quad (1)$$

We assume that $n_z(r)$ is a function which increases monotonically from the value $n_z(0) = -1$ at the skyrmion center to $n_z(r \geq R_0) = +1$ beyond a cutoff radius R_0 . The fixed angle ϕ_0 determines the skyrmion handedness. We highlight two special cases: “hedgehog”-type Néel skyrmions characterized by $\phi_0 = 0$ and “vortex”-type Bloch skyrmions which have $\phi_0 = \pm\pi/2$. The skyrmion topological charge

$$Q_{\text{top}} = -\frac{1}{4\pi} \int d^2\mathbf{r} \, \mathbf{n} \cdot \partial_x \mathbf{n} \times \partial_y \mathbf{n} = 1 \quad (2)$$

is independent of ϕ_0 and is invariant under local deformations of the texture. We do not discuss the energetic stability of the various skyrmion types but instead present results for both Bloch and Néel skyrmions.

The radial function $n_z(r)$ may in general have a sharp, i.e., domain-wall-like, or more gradual transition as a function of

r . We parametrize this freedom by

$$n_z(r) = \begin{cases} -1 & r \in [0, \alpha R_0] \\ 2 \sin^2\left(\frac{\pi(r - \alpha R_0)}{2(R_0 - \alpha R_0)}\right) - 1 & r \in (\alpha R_0, R_0) \\ +1 & r \geq R_0 \end{cases}. \quad (3)$$

In the limit $\alpha \rightarrow 1$, this ansatz leads to a sharp transition, with $n_z(r < R_0) = -1$ and $n_z(r > R_0) = +1$; in this case, the skyrmion approaches the form of a minority domain droplet with no in-plane magnetization component, and the distinction between Néel versus Bloch skyrmion loses its significance. On the other hand, the transition is smooth for all $\alpha < 1$, with the most gradual transition $n_z(r) = 2 \sin^2(\pi r / 2R_0) - 1$ obtained when $\alpha = 0$. Tuning $\alpha \in [0, 1)$ allows us to interpolate between these two limits, and we find that many of our results concerning the Chern numbers of the skyrmion bands depend crucially on this parameter. For later use, we also define the skyrmion core size by

$$R = R_0(1 + \alpha)/2, \quad (4)$$

which is where $n_z(r)$ undergoes a change in sign. This radius R will later be found to determine the radius of ringlike Wannier functions obtained from the skyrmion bands and, more broadly, is a convenient tuning parameter for studying the skyrmion bands and their wave-function topology. We remark that a variety of alternative forms for $n_z(r)$ in the interval $(\alpha R_0, R_0)$ have been investigated, but these lead to only minor quantitative differences as compared to Eq. (3). To simplify the discussion, we therefore focus on this particular form.

We construct the skyrmion crystal ansatz as a triangular lattice of skyrmions centered at Bravais vectors

$$\mathbf{R} = m_1 \mathbf{a}_1 + m_2 \mathbf{a}_2, \quad m_{1,2} \in \mathbb{Z}, \quad (5)$$

where $\mathbf{a}_1 = a(1, 0)$ and $\mathbf{a}_2 = a(\frac{1}{2}, \frac{\sqrt{3}}{2})$. We assume for simplicity that $a > 2R_0$, so that individual skyrmions in the crystal do not directly overlap, therefore allowing the magnetization $n_z(\mathbf{r}) = +1$ of adjacent skyrmions to join smoothly at their Wigner-Seitz cell boundaries. Note that the spacing between skyrmions is controlled solely by the parameter a , not by the quantities R , R_0 or α . We will denote the skyrmion reciprocal lattice by \mathfrak{G} and its primitive vectors by

$$\mathbf{Q}_1 = Q(\sqrt{3}/2, -1/2), \quad \mathbf{Q}_2 = (0, Q) \quad (6)$$

with $Q = 4\pi/\sqrt{3}a$. The skyrmion lattice texture can be written as a Fourier series in these reciprocal lattice vectors, $\mathbf{n}(\mathbf{r}) = \sum_{\mathbf{G} \in \mathfrak{G}} \mathbf{n}_{\mathbf{G}} e^{i\mathbf{r} \cdot \mathbf{G}}$.

B. Coupling helical Dirac fermions to the magnetic skyrmion crystal

The continuum Hamiltonian for the TI Dirac surface states coupled to the skyrmion spin texture via a Hund’s term is given by $H = H_0 + H_1$, where

$$\begin{aligned} H_0 &= v_F \int_{\mathbb{R}^2} d^2\mathbf{r} \, c^\dagger(\mathbf{r}) \left[\left(-i\hbar \frac{\partial}{\partial \mathbf{r}} \times \boldsymbol{\sigma} \right) \cdot \hat{\mathbf{z}} \right] c(\mathbf{r}) \\ H_1 &= J_{\text{eff}} \int_{\mathbb{R}^2} d^2\mathbf{r} \, c^\dagger(\mathbf{r}) \mathbf{n}(\mathbf{r}) \cdot \boldsymbol{\sigma} c(\mathbf{r}). \end{aligned} \quad (7)$$

Here, the fermion operators are implicitly spinors. The quantities v_F and J_{eff} denote the Dirac velocity and spin-fermion coupling strength, respectively.

Here we argue that contributions from external magnetic fields may be neglected in various cases. First, we note that a nonvanishing Zeeman shift may be absorbed into the skyrmion texture without modifying any Hamiltonian symmetries. Thus, we only need to discuss the impact of an orbital magnetic field. In cases where the Zeeman field is necessary to stabilize skyrmions, the corresponding orbital magnetic flux per skyrmion lattice unit cell is expected to be small compared to the flux quantum in the case of small skyrmions. For example, in MnBi_2Te_4 , the required magnetic field is of the order of $B \sim 0.02 \text{ meV}/g\mu_B \sim 0.2 \text{ T}$. This gives a magnetic length of $\ell_B \sim 60 \text{ nm}$, so that the magnetic flux is small, $Ba^2 \sim \Phi_0 a^2/2\pi\ell_B^2 \ll \Phi_0$ with $a \sim 4 \text{ nm}$ [59]. Finally, we note that it may be possible to interface TI surface states with materials hosting skyrmions at zero field, such as those recently reported to be stabilized by frustration [82] or soft x-ray illumination [83].

Henceforth, we will measure energies in units of $\hbar v_F/a$. Let us denote the dimensionless spin-fermion exchange coupling by $J = (a/\hbar v_F)J_{\text{eff}}$. To estimate this in the magnetic topological insulator MnBi_2Te_4 , we set $\hbar v_F \sim 1 \text{ eV \AA}$ as measured from angle resolved photoemission spectroscopy (ARPES) [58]. The spin-fermion coupling J_{eff} may be crudely estimated via $T_N \sim J_{\text{eff}}^2/W$ within an RKKY picture, where the bandwidth is $W \sim 1 \text{ eV}$ based on ARPES and the experimental Néel temperature is $T_N \sim 25 \text{ K}$ [58]. This leads to $J_{\text{eff}} \sim \sqrt{T_N W} \sim 50 \text{ meV}$. To estimate the skyrmion lattice constant, we note that skyrmions in MnSi have $a \sim 20 \text{ nm}$, but MnBi_2Te_4 has been argued to support skyrmions with a smaller lattice constant $\sim 4 \text{ nm}$ [59]. Taking $a \sim 4\text{--}20 \text{ nm}$ translates to a range of a dimensionless spin-fermion couplings $J \sim 1\text{--}10$.

Moving to momentum space and folding into the first Brillouin zone (BZ) of the skyrmion lattice,

$$c_{s\mathbf{G}}(\mathbf{k}) \equiv c_s(\mathbf{k} + \mathbf{G}) = \int_{\mathbb{R}^2} d^2\mathbf{r} e^{-i(\mathbf{k}+\mathbf{G})\cdot\mathbf{r}} c_s(\mathbf{r}) \quad (8)$$

the Hamiltonian is block diagonal in the crystal momentum \mathbf{k} due to discrete translational symmetry:

$$H = \int_{\text{BZ}} \frac{d^2\mathbf{k}}{(2\pi)^2} \sum_{\mathbf{G}, \mathbf{G}' \in \mathfrak{G}} c_{\mathbf{G}}^\dagger(\mathbf{k}) (\mathcal{H}_{\mathbf{k}})_{\mathbf{G}, \mathbf{G}'} c_{\mathbf{G}'}(\mathbf{k}). \quad (9)$$

In this basis, the Hamiltonian has matrix elements

$$(\mathcal{H}_{\mathbf{k}})_{\mathbf{G}, \mathbf{G}'} = \delta_{\mathbf{G}, \mathbf{G}'} ((\mathbf{k} + \mathbf{G}) \times \boldsymbol{\sigma})_z - J \mathbf{n}_{\mathbf{G}-\mathbf{G}'} \cdot \boldsymbol{\sigma} \quad (10)$$

where the Pauli matrices correspond to spin. This defines for us the electronic band structure problem for the skyrmion crystal. Since the set of skyrmion reciprocal lattice vectors \mathfrak{G} is infinite, each matrix $\mathcal{H}_{\mathbf{k}}$ possesses infinitely many components. Computation of the band structure therefore requires truncating \mathfrak{G} to some finite number of reciprocal vectors nearest zero. This truncation is justified by the absence of any singularity in the skyrmion spin texture, corresponding to rapid decay of its Fourier components. For the range of parameters $0 \leq J \lesssim 10$ we find that truncation to 300 momenta, and therefore 600 bands due to spin, is sufficient to

attain convergence in the energies, Berry curvature, and tight-binding parameters considered later.

The Bloch vectors $u_{s\mathbf{G},n}(\mathbf{k})$ are defined as the eigenvectors of $\mathcal{H}_{\mathbf{k}}$ and allow us to define the Bloch operators

$$\psi_{kn}^\dagger = \sum_{s\mathbf{G}} c_{s\mathbf{G}}^\dagger(\mathbf{k}) u_{s\mathbf{G},n}(\mathbf{k}). \quad (11)$$

By construction, the second-quantized Hamiltonian is diagonal in the band basis,

$$H = \int_{\text{BZ}} \frac{d^2\mathbf{k}}{(2\pi)^2} \sum_n \epsilon_n(\mathbf{k}) \psi_{kn}^\dagger \psi_{kn}. \quad (12)$$

Before we separately present results for the band structure for Bloch and Néel skyrmions, we note that the skyrmion crystal reduces the continuous rotational symmetry of the isolated skyrmion problem to a sixfold symmetry. We fix the rotation axis to be parallel to \hat{z} , passing through a skyrmion center. We represent the sixfold operator on the continuum states by

$$C_6 c(\mathbf{r}) C_6^\dagger = e^{i\frac{\pi}{3} \frac{\sigma_z}{2}} c(C_6 \mathbf{r}), \quad (13)$$

where C_6 acts as a sixfold counterclockwise rotation on vectors. By appeal to the identity $C_6 \mathbf{n}(C_6^{-1} \mathbf{r}) = \mathbf{n}(\mathbf{r})$, one finds that C_6 is a symmetry (see Supplemental Material [84]), independent of the value of ϕ_0 which sets the skyrmion type.

III. BLOCH SKYRMION LATTICE

A. Continuum model

When the magnetization texture $\mathbf{n}(\mathbf{r})$ describes a lattice of Bloch skyrmions, given by setting $\phi_0 = \pi/2$ or $-\pi/2$, we are granted several additional symmetries which constrain the energy bands and Berry curvature. Setting ϕ_0 accordingly in Eq. (1), one observes that the magnetization of an isolated Bloch skyrmion satisfies the property $\mathbf{r} \cdot \mathbf{n}(\mathbf{r}) = 0$. Consequently, the in-plane divergence must vanish identically, $\nabla^{2D} \cdot \mathbf{n}(\mathbf{r}) = 0$ due to the divergence theorem [68]. In this case, it has been shown [69] that the in-plane component of $\mathbf{n}(\mathbf{r})$ may be entirely removed from the Hamiltonian by the gauge transformation $\mathcal{U}(\mathbf{r}) = \exp(iJ \int_0^\rho d\rho' \sqrt{1 - n_z^2(\rho')})$. All of these statements remain true when we promote the texture to a triangular lattice of skyrmions rather than an isolated skyrmion. One must simply interpret $\mathbf{r} = (\rho, \phi)$ in Eq. (1) as a quantity measured relative to the nearest skyrmion center.

The fact that the symmetry $\mathcal{U}(\mathbf{r})$ is periodic in the skyrmion lattice ensures that the band structure is invariant under its action. This is because it performs a unitary convolution on each subspace of definite crystal momentum. Therefore, we may drastically simplify the arguments that follow by setting the in-plane magnetization to zero outright. We denote the resulting texture by a separate symbol to distinguish it from the original periodic texture:

$$\mathfrak{n}(\mathbf{r}) \equiv (0, 0, n_z(\mathbf{r})). \quad (14)$$

Note that $\mathfrak{n}(\mathbf{r})$ is neither normalized nor does it possess a winding number. Moreover, the texture crucially possesses the symmetries

$$\mathfrak{n}(\mathbf{r}) = \mathfrak{n}(-\mathbf{r}) = \mathfrak{n}(M_y \mathbf{r}) \quad (15)$$

with M_y defined as the matrix which flips the second component of a vector. We now proceed with a description the symmetries manifesting in the continuum description of the Bloch skyrmion lattice problem.

1. Particle-hole symmetry

Consider the unitary transformation which flips spins and exchanges electrons with holes at fixed momentum,

$$Uc(\mathbf{k})U^\dagger = c^\dagger(\mathbf{k})\sigma_y, \quad UiU^\dagger = i. \quad (16)$$

Leveraging the identity $\mathbf{n}(\mathbf{r}) = \mathbf{n}(-\mathbf{r})$, we demonstrate (see Supplemental Material [84]) that U is a symmetry of the Hamiltonian, satisfying $[U, H] = 0$. Such a symmetry is commonly referred to as a ‘particle-hole’ symmetry [85]. Analyzing this symmetry in reciprocal space reveals that the band structure is symmetric about zero energy at each value of the crystal momentum. Indeed, in the notation of Eq. (10) we find that the particle-hole symmetry is expressed by the matrix relation

$$\sigma_y(\mathcal{H}_k)^*\sigma_y = -\mathcal{H}_k. \quad (17)$$

Since \mathcal{H}_k is Hermitian then its spectrum is invariant under both complex conjugation and change of basis. The above equation thus implies that its spectrum is symmetric, i.e., that its eigenvalues come in pairs $\epsilon_{-n}(\mathbf{k}) = -\epsilon_n(\mathbf{k})$. We have adopted a convention where the valence (conduction) bands are labeled by negative (positive) integers in order of their energy.

2. Chiral symmetry

Continuing with our analysis of the Bloch skyrmion lattice, we now show that the Hamiltonian Eq. (7) satisfies a ‘chiral’ symmetry [85] which constrains the band structure and Berry curvature. Consider the anti-unitary operator which flips spin and exchanges electrons with holes at mirror-related momenta,

$$Ac(\mathbf{k})A^\dagger = c_{M_y\mathbf{k}}^\dagger(M_y\mathbf{k})\sigma_x, \quad AiA^\dagger = -i. \quad (18)$$

At the level of the Hamiltonian kernel, the symmetry $[A, H] = 0$ derives from the identity

$$(\mathcal{H}_k)_{G,G'} = -(\sigma_x \mathcal{H}_{M_y\mathbf{k}} \sigma_x)_{M_yG, M_yG'}. \quad (19)$$

For each eigenvector $u_n(\mathbf{k})$ of \mathcal{H}_k , we therefore have a related eigenstate $u_{-n}(M_y\mathbf{k})$ of $\mathcal{H}_{M_y\mathbf{k}}$ with opposite energy. As shown explicitly in the Supplemental Material [84], this provides a relation between the Berry curvature of the opposing $\pm n$ bands at these mirror-related momenta,

$$F^{(n)}(\mathbf{k}) = -F^{(-n)}(M_y\mathbf{k}). \quad (20)$$

Upon integrating over the Brillouin zone, we immediately see that these bands must carry opposite Chern number, $\mathcal{C}_{-n} = -\mathcal{C}_n$. As we later discuss, this property is absent from the Néel band structure.

3. Topological bands and DOS

In Fig. 2 we illustrate the band structure for the continuum Hamiltonian Eq. (7) in the Bloch skyrmion case. In accordance with the particle-hole symmetry presented in Eq. (17),




FIG. 1. (a) Bravais vectors $\mathbf{a}_1, \mathbf{a}_2$ and overhead view of the triangular lattice of Bloch skyrmions at $\alpha = 0.5$ and cutoff radius $R_0/a = 0.49$. The color corresponds to the out-of-plane magnetization n_z whereas the vector field indicates the in-plane magnetization. (b) The single-skyrmion magnetization profile n_z plotted for various values of α which interpolates smoothly between the sinusoidal and domain wall limits. For each α we mark in orange the corresponding radius R at which the out-of-plane magnetization changes sign.

the energy spectrum is symmetric everywhere in the skyrmion Brillouin zone. We exhibit the bands for fixed parameter values $J = 3$ and $R_0/a = 0.49$, with the latter chosen so as to encourage hybridization between the single-skyrmion bound states. We recall that R_0 sets the cutoff radius at which the skyrmion magnetization is purely polarized in the \hat{z} direction whereas the core size R , as illustrated in Fig. 1, sets the radius at which the Dirac mass n_z changes sign.

At $R/a = 0.28$ the four bands nearest half filling each carry zero Chern number and are continuously connected to an atomic insulator phase in the limit of large J . When the skyrmion core size R/a is increased up to 0.37 we observe a gap closure between the $n = \pm 1$ bands at the Γ point upon which the system enters a new topological sector with Chern numbers $\mathcal{C} = (0, -1 : 1, 0)$ for the four bands nearest half filling. As R is increased toward its maximal value $R_0/a = 0.49$, we reach the value $R/a = 0.44$ at which a final topological transition takes us into the $\mathcal{C} = (1, -1 : 1, -1)$ sector, this time due to a gap closure at the Γ point between the $|n| = 2, 3$ bands. Unlike in a related moiré study [50], these remote Dirac cones are not protected by time reversal, nor can their velocity be easily tuned to zero within the present model.

It is important to distinguish this ‘‘intrinsic’’ anomalous Hall effect, permitted by broken time-reversal symmetry, from the well-studied THE which arises when nonrelativistic electrons are coupled to a skyrmion lattice [72,73]. The THE is absent for Dirac electrons since the effective magnetic flux density $\mathbf{B}_{\text{eff}} \sim \nabla^{2D} \cdot \mathbf{n}$ seen by the Dirac electron in a skyrmion texture does not depend on the topological charge Eq. (2) of the magnetic texture [86]. Indeed, this effective flux density, when integrated over the skyrmion unit cell, vanishes for any generic skyrmion texture due to the divergence theorem (for any type of skyrmion). For Bloch skyrmions, an even stronger condition holds, that \mathbf{B}_{eff} itself vanishes everywhere in space.

The symmetry $\mathcal{C}_{-n} = -\mathcal{C}_n$ of these Chern numbers is consistent with the chiral symmetry constraint, Eq. (20). In each




FIG. 2. Bloch skyrmion bands and density of states (DOS) at $J = 3$ and cutoff radius $R_0/a = 0.49$. (a) At small R/R_0 , the four lowest-energy bands carry zero Chern number and we observe windows of vanishing DOS. (b) As the core size R is increased there is a gap closure (circled) between the $n = \pm 1$ bands at the Γ point, resulting in Chern numbers $\mathcal{C} = (0, -1:1, 0)$. The dispersion and DOS at this transition are linear. (c) As R is increased towards its maximum value R_0 , the bands undergo a final transition into the $\mathcal{C} = (1, -1:1, -1)$ sector upon direct gap closure between the $|n| = 2, 3$ bands at the Γ point. Increasing J leads to narrower Chern bands and more pronounced peaks in the DOS.

case, the Chern numbers were computed from the band eigenstates following the methodology and sign conventions of Ref. [87]. In Fig. 3 we display a phase diagram indicating the Chern numbers carried by the four lowest-energy bands over the range of parameters $J \in [1, 8]$ and $R \in [R_0/2, R_0]$ at fixed $R_0/a = 0.49$. We omit the range $J \in [0, 1]$ where the bands too closely resemble those of the free Dirac Hamiltonian. We observe that all horizontal cuts along the phase diagram realize the same topological phases. To aid in visualizing the appearance of the corresponding bands, we mark in this figure those values of R whose bands are displayed in Fig. 2.

As the cutoff radius R_0 is decreased, the $\mathcal{C} = (1, -1:1, -1)$ region recedes completely, followed by the $\mathcal{C} = (0, -1:1, 0)$ phase. Once the cutoff radius has reached $R_0 = 0.35a$, all four bands carry zero Chern number over the entire




FIG. 3. Chern number phase diagram for the four lowest-energy bands in the Bloch skyrmion case. The chosen value $R_0/a = 0.49$ corresponds to a near-maximally dense packing of skyrmions. The Chern numbers in each tuple \mathcal{C} are listed in order of increasing energy with a colon marking half filling. In agreement with the Berry curvature constraint Eq. (20), in each sector we observe that the Chern numbers of bands of opposite energy are opposite, $\mathcal{C}_{-n} = -\mathcal{C}_n$. For ease of comparison, we mark points corresponding to the band structures presented in Fig. 2.

range of parameters (R, J) . This phase can be understood as being continuously connected to the limit of well-separated Wannier orbitals which resemble the electronic bound states of isolated skyrmions. On the other hand, fixing $R_0/a \sim 0.49$ as in Fig. 3 and increasing the skyrmion core size R toward R_0 , the domain wall limit, we find that increasing J leads to the $|n| = 1, 2$ bands bunching together and flattening, reminiscent of the single-particle physics of magic angle bilayer graphene [26]. In the trivial Chern sector, increasing J more simply increases the number of in-gap bands, which flatten and are completely isolated.

B. Tight-binding model for the Bloch skyrmion lattice problem

We can approach the problem of Dirac electrons coupled to a Bloch skyrmion texture from a complementary angle, namely a tight-binding approach. In the previous sections we argued that the continuum problem Eq. (7) features both a particle-hole symmetry, responsible for a symmetric energy spectrum, and a chiral symmetry, which dictates that the particle-hole-related bands must carry opposite Chern number. Well-known theoretical results about the interplay between Wannier functions and band topology guarantee the existence [88] of a Wannier representation for any even number of nondegenerate bands nearest half filling. In the limit of large interskyrmion distance $a \gg R_0$, we expect the Wannier orbitals to approach single-skyrmion electronic bound states, whose features we review in the following section. Application of the Wannier projection method [89,90] for the two gapped bands nearest half filling, which we discuss in Sec. III C 1, reveals well-localized Wannier states whose

qualitative features match those of the single-skyrmion bound states.

1. Single-skyrmion bound states

The problem of Dirac fermions coupled to a single skyrmion was addressed by Hurst *et al.* in Ref. [68]. In that work, the authors considered a two-dimensional Dirac model with a position-dependent Dirac mass representing the skyrmion. The corresponding wave functions were found to be strongly peaked at the skyrmion perimeter where the Dirac mass changes sign. Further studies [71] found similar states for a more realistic description of the skyrmions. Most importantly, the skyrmion bound states are a discrete set of states with energies inside the bulk electronic gap and a well-defined out-of-plane angular momentum $j = \pm 1/2, \pm 3/2, \dots$ arising from the rotational symmetry of the single skyrmion texture. The bound state wave functions take the form

$$\Psi_j(\mathbf{r}) = \begin{pmatrix} e^{i(j-\frac{1}{2})\phi} \chi_{j,\uparrow}(r) \\ e^{i(j+\frac{1}{2})\phi} \chi_{j,\downarrow}(r) \end{pmatrix}. \quad (21)$$

The asymptotic behavior of the radial wave functions $\chi_{j,s}(r)$ has been extensively studied [68,69,71]. While their exact form is not important for our purposes, they are known to decay exponentially at long distances.

2. From Bloch states to Wannier states

Combining Bloch skyrmions together in a hexagonal lattice, we expect the skyrmion bound states to hybridize to form orthogonal Wannier states. Supposing we have identified some set of isolated bands

$$n \in \{\pm 1, \pm 2, \dots, \pm n_0\} \equiv \mathcal{B} \quad (22)$$

nearest half filling, the Wannier states are merely the Fourier transform of a smoothening of the Bloch states

$$d_{\mathbf{R}j}^\dagger = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} d_{\mathbf{k}j}^\dagger e^{-i\mathbf{k} \cdot \mathbf{R}} \quad (23)$$

for a choice of unitaries $\mathcal{U}_{nj}(\mathbf{k})$ such that the rotated states

$$d_{\mathbf{k}j}^\dagger = \sum_{n \in \mathcal{B}} \psi_{kn}^\dagger \mathcal{U}_{nj}(\mathbf{k}) \quad (24)$$

are smooth in the variable \mathbf{k} . Less abstractly, these orbitals correspond to smooth functions $d_{\mathbf{s}kj}(\mathbf{r}) = \langle \Omega | c_{\mathbf{s}}(\mathbf{r}) d_{\mathbf{k}j}^\dagger | \Omega \rangle$. In Sec. III C 1 we implement a technique, known as the projection method [89,90], for constructing the unitaries $\mathcal{U}_{nj}(\mathbf{k})$ directly from an initial guess for the Wannier orbitals. In particular, the constructed Wannier orbitals will carry an eigenvalue j under C_6 , justifying the choice of j as a label.

In a tight-binding representation of a manifold of bands \mathcal{B} , the Hamiltonian data is encoded in a set of amplitudes

$$t_{\delta jj'} = \langle \delta + \mathbf{R} j | H | \mathbf{R} j' \rangle \quad (25)$$

where δ, \mathbf{R} are skyrmion Bravais vectors. The independence of the hoppings on \mathbf{R} is due to discrete translational symmetry. Since the Hamiltonian does not mix states with different angular momentum then the on-site overlap matrix $t_{0jj'} \equiv \varepsilon_j \delta_{jj'}$ is

diagonal. The orthogonal projection of the continuum Hamiltonian Eq. (12) into the bands \mathcal{B} then reads

$$H_{\mathcal{B}} = \sum_{\mathbf{k} \in \text{BZ}} \sum_{jj'} d_{\mathbf{k}j}^\dagger H_{\mathbf{k}jj'} d_{\mathbf{k}j'} \quad (26)$$

with momentum space kernel given by

$$H_{\mathbf{k}jj'} \equiv \delta_{jj'} \varepsilon_j + \sum_{\delta} e^{-i\mathbf{k} \cdot \delta} t_{\delta jj'}. \quad (27)$$

In the following, we discuss how the rotational, particle-hole, and chiral symmetries act on the Wannier states as well as the constraints they impose on the hopping parameters. We focus here on nearest-neighbor hoppings, leaving a detailed analysis of next and next-next-nearest neighbor hoppings to the Supplemental Material [84]. In the nearest-neighbor case, the displacement vector δ runs over the six nearest neighbors

$$\delta = \pm \mathbf{a}_1, \pm \mathbf{a}_2, \pm \mathbf{a}_3 \quad (28)$$

where $\mathbf{a}_{1,2}$ are defined in Eq. (5) and $\mathbf{a}_3 = \mathbf{a}_2 - \mathbf{a}_1$. At this stage we do not limit our analysis to any particular number of bands $|\mathcal{B}|$.

3. Representing the sixfold symmetry

While the lattice of skyrmions breaks the continuous rotation symmetry of the single skyrmion texture, the C_6 of Eq. (13) still remains. We choose to implement it on the Wannier orbitals by

$$C_6 d_{\mathbf{R}j} C_6^\dagger = e^{i\frac{2\pi}{3}j} d_{C_6 \mathbf{R}j} \quad (29)$$

which conveniently leads to Wannier functions which transform under C_6 like the single-skyrmion bound states, Eq. (21). Moreover, the hoppings are constrained (see Supplemental Material [84]) to satisfy

$$t_{C_6 \delta jj'} = e^{-i\frac{2\pi}{3}(j-j')} t_{\delta jj'}. \quad (30)$$

This property allows us to express all the nearest neighbor hopping parameters in terms of $t_{jj'} \equiv t_{\mathbf{a}_1 jj'}$ alone. Correspondingly, Eq. (27) takes the compact form

$$H_{\mathbf{k}jj'} = \delta_{jj'} \varepsilon_j + s_{\mathbf{k}jj'} t_{jj'} \quad (31)$$

where $s_{\mathbf{k}jj'}$, a function of the nearest-neighbor lattice geometry whose precise form is inessential to the present discussion, may be found in the Supplemental Material along with the generalization to further neighbors [84].

For later reference, we remark that Hermiticity and C_2 , whose action is obtained by three applications of the C_6 constraint Eq. (30), together guarantee

$$t_{j'j} = e^{-i\pi(j-j')} (t_{jj'})^*. \quad (32)$$

Finally, while our tight-binding construction is nominally performed for the Bloch case for topological reasons, we remark that these C_6 results apply equally well to the Néel case provided that one is modeling a gapped subset of bands with net zero Chern number.

4. Representing the particle-hole symmetry

The particle-hole and chiral symmetries hold only in the Bloch case. We represent the former on the Wannier states by

$$U d_{kj} U^\dagger = (d_{k-j})^\dagger e^{i\pi j}. \quad (33)$$

By equating the band-projected Hamiltonian H_B to the particle-hole conjugated expression

$$U H_B U^\dagger = - \sum_{k,j,j'} d_{kj}^\dagger [H_{k-j'-j} e^{i\pi(j'-j)}] d_{kj'} + \sum_{kj} H_{kjj},$$

we immediately deduce that

$$H_{k-j'-j} e^{i\pi(j'-j)} = -H_{kjj}. \quad (34)$$

In the Supplemental Material [84] we show that this condition is satisfied if and only if

$$\begin{aligned} \varepsilon_{-j} &= -\varepsilon_j \\ t_{-j'-j} &= -e^{i\pi(j'-j)} t_{jj'}. \end{aligned} \quad (35)$$

5. Representing the chiral symmetry

The final constraints on the hopping parameters derive from the chiral symmetry Eq. (18), which we represent on the Wannier orbitals by

$$A d_{kj} A^\dagger = (d_{M,k-j})^\dagger. \quad (36)$$

As in our particle-hole analysis, comparison of H_B to

$$A H_B A^\dagger = - \sum_{k,j,j'} d_{kj}^\dagger [H_{M,k-j-j'}] d_{kj'} + \sum_{kj} H_{kjj} \quad (37)$$

demands immediately that

$$H_{M,k-j-j'} = -H_{kjj}. \quad (38)$$

In the Supplemental Material [84] we show that this condition is satisfied if and only if

$$\begin{aligned} \varepsilon_{-j} &= -\varepsilon_j \\ t_{-j-j'} &= -t_{jj'}. \end{aligned} \quad (39)$$

Together with Eqs. (32) and (35), these expressions constitute all independent constraints on the parameters ε_j and $t_{jj'}$ due to the symmetries of the Bloch skyrmion lattice system.

C. Two-band Wannier construction

We now perform an explicit construction of Wannier functions for the gapped low-energy bands of the Bloch skyrmion problem. We treat the two bands of lowest energy, $n \in \mathcal{B} = \{\pm 1\}$. In our analysis of the continuum theory, these bands were found to maintain a finite energy gap to the remaining higher-energy states over the entire phase diagram Fig. 3, with a gap closure occurring only mutually between them at zero energy.

The resulting tight-binding model features two Wannier states, labeled by their out-of-plane angular momentum $j = \pm 1/2$, and localized at each skyrmion site. Their wave functions are found to be tightly concentrated at the radius R and exponentially decaying at long distances, which is reminiscent of the single-skyrmion bound states discussed in Sec. III B 1, even for a reasonably closely packed skyrmion lattice.

Finally, we numerically observe that the nearest-neighbor hopping amplitudes are generically dominant. This inspires the independent study of a nearest-neighbor two band toy model whose Chern sectors can be characterized analytically as a function of the hopping amplitudes $t_{-1/2, +1/2}$ and $t_{-1/2, -1/2}$ of Eq. (31). We find that the hopping amplitudes realized by the continuum model account for only half of the topological phases present in the two band toy model, thereby motivating further study into variants of Eq. (7) as a means of realizing novel Chern insulator phases.

1. Wannier orbitals of Bloch skyrmion bands

To construct the $j = \pm 1/2$ Wannier states from the pair $n = \pm 1$ of lowest-lying energy bands, we employ the projection method [89,90]. One begins with a set of trial orbitals $g_{sj}(\mathbf{r})$ which serve as a best guess for the Wannier orbitals centered in the home unit cell $\mathbf{R} = \mathbf{0}$. These trial states are then projected into the manifold of Bloch states at each wave vector \mathbf{k} ,

$$|\phi_{kj}\rangle = \sum_{n=\pm 1} |\psi_{kn}\rangle \langle \psi_{kn} | g_j \rangle, \quad (40)$$

where the latter inner product is taken over all of space. To ensure that these states are orthonormal, one performs a Löwdin transformation

$$|d_{kj}\rangle = \sum_{j'} |\phi_{kj'}\rangle (\mathcal{S}_k^{-1/2})_{j'j} \quad (41)$$

where the overlap matrix is given by

$$\mathcal{S}_k = \mathcal{A}_k^\dagger \mathcal{A}_k, \quad (\mathcal{A}_k)_{nj} \equiv \langle \psi_{kn} | g_j \rangle. \quad (42)$$

The Bloch-like states $|d_{kj}\rangle$ will be smooth in \mathbf{k} , and the corresponding Wannier orbitals Eq. (23) will be exponentially localized, provided that the overlap matrix is finite everywhere in the BZ. As an immediate consequence of the above definitions, we obtain the unitary transformation taking us between the Bloch-like states d_{kj} and the original Bloch states. In the notation of Eq. (24), the unitary is given by

$$\mathcal{U}_{nj}(\mathbf{k}) = \sum_{j'} (\mathcal{A}_k)_{nj'} (\mathcal{S}_k^{-1/2})_{j'j}. \quad (43)$$

The Wannier orbitals can be shown to inherit the transformation properties of the trial orbitals under the symmetries of the Hamiltonian. In the Supplemental Material [84] we demonstrate that the representations Eqs. (29), (33), (36) may be enforced by the choice of trial orbitals

$$g_{sj}(r, \phi) = e^{i(j-\frac{s}{2})\phi} e^{-(r-\mu)^2/2\xi^2} \quad (44)$$

where the integer-valued term $j - s/2 = j \pm 1/2$ has the interpretation of orbital angular momentum. In Fig. 4 we plot the resulting Wannier function densities corresponding to trial parameters with annular peak at $\mu = R$ and spread equal to the width of skyrmion wall, $\xi = R_0 - R$. This choice of the parameters (μ, ξ) yields a favorable ratio of the maximum and minimum values of $\det S_k$ over the BZ, namely < 2 for all parameters (α, J) considered, therefore indicating a smooth fit. Remarkably, the same Wannier functions result from instead inputting Gaussian trial functions $\mu = 0$ with various ξ , indicating that the annular features of the Wannier functions are




FIG. 4. (a) Wannier function densities $\sum_s |d_{s0j}(\mathbf{r})|^2$ derived from the lowest two Bloch skyrmion bands at $R_0/a = 0.49$. The density is equal for opposite j . The black hexagon indicates the boundary of a Wigner-Seitz cell of the skyrmion lattice and the orange ring indicates the skyrmion core size, $R = R_0(1 + \alpha)/2$. The Wannier function density accumulates at R and is more tightly localized for larger Hund's coupling J . For small skyrmion size, the Wannier functions closely resemble the single skyrmion bound states. For larger skyrmion radius R (i.e., larger α), the lower sixfold symmetry of the Wannier function density is more apparent. (b) Hopping parameters derived from the Wannier functions at $(J, R_0/a) = (3, 0.49)$. The nearest-neighbor hoppings $\tau, \tilde{\tau}$ are most prominent. The shaded region marks the transition from the $\mathcal{C} = (0:0)$ sector to $\mathcal{C} = (-1:1)$, whereas the vertical line at $\tau/\varepsilon = -1/6$ marks where this boundary shifts upon truncating the hoppings at nearest neighbor.

a product of the Hamiltonian and not the trial functions. Further indication of our successful application of the projection method is provided by the Wannier functions $d_{s0j}(\mathbf{r})$ decaying exponentially with distance from the skyrmion center, which we verified numerically over the range of several unit cells.

The Wannier functions Fig. 4(a) are found to localize around the radius R at which the Dirac mass $n_z(r)$ changes sign. Moreover, the states localize more tightly around R as the Hund's coupling J is increased. These features are consistent with the behavior of the previously studied single-skyrmion bound states [68,71]. Where the lattice problem differs, however, is in the breaking of continuous rotational symmetry in the Wannier states at large $R \rightarrow R_0 \sim a/2$ where the hybridization between the single-skyrmion bound states is largest due to proximity.

2. Nearest-neighbor toy model

When the skyrmion cutoff radius is not too large compared to the Hund's length scale, $R_0 J_{\text{eff}}/\hbar v_F \lesssim 3$, only the two lowest-lying bands are uniformly gapped from continuum of states at higher energies. Correspondingly, it is shown in Ref. [68] that only two electronic bound states accompany an isolated skyrmion for sufficiently small skyrmion radius. These observations motivate further investigation into the two-band tight-binding problem in particular. We note that Chern insulators in similar two-band lattice models with local orbitals having nonzero angular momentum have been discussed previously in the context of spin-orbit coupled ferromagnets [91,92].

In Fig. 4(b) we plot the two-band hopping parameters as a function of the skyrmion radius for fixed $J = 3$ and

$R_0/a = 0.49$. As discussed above and further detailed in the Supplemental Material [84], constraints from symmetry ensure that the parameters $\varepsilon, t_{jj'}$, and $t_{jj'}^n$ characterizing the hopping Hamiltonian up to next-nearest neighbor are all real. Crucially, we find that the nearest-neighbor terms $t_{jj'}$ are more prominent than longer-range terms for a wide range of parameters (J, R_0) , thereby motivating a full analytical investigation of the two-band *nearest-neighbor* toy model.

In this case, the symmetry constraints Eqs. (32), (35), (39) imply that the only free parameters are

$$\begin{aligned} \varepsilon &\equiv \varepsilon_{-1/2} \\ \tau &\equiv t_{-1/2, -1/2} \\ \tilde{\tau} &\equiv t_{+1/2, -1/2} \end{aligned} \quad (45)$$

and that they must all be real. The Hamiltonian Eq. (31) then takes the simple form $H_{\mathbf{k}} = h_{\mathbf{k}} \cdot \boldsymbol{\sigma}$, where

$$h_{\mathbf{k}} \equiv (\tilde{\tau} \text{Re} Q_{\mathbf{k}}, -\tilde{\tau} \text{Im} Q_{\mathbf{k}}, -(\varepsilon + \tau P_{\mathbf{k}})), \quad (46)$$

and $P_{\mathbf{k}} \equiv s_{kjj}$, $Q_{\mathbf{k}} \equiv s_{kjj-1}$. This gives rise to energies

$$E_{\mathbf{k}\pm} = \pm |h_{\mathbf{k}}| = \pm \sqrt{|\tilde{\tau} Q_{\mathbf{k}}|^2 + (\varepsilon + \tau P_{\mathbf{k}})^2}. \quad (47)$$

Within this toy model, the Chern numbers of the bands $n = \pm$ are given by the number of times the unit vector $h_{\mathbf{k}}$ covers the unit sphere as the momentum is scanned through the BZ. Consequently, flipping the sign of $\tilde{\tau}$ has no bearing on the Chern number since it merely changes the helicity of the momentum space texture Eq. (46) and not its winding. Due to continuity, the phase boundaries must occur at gap closures between the two bands, i.e., where $|h_{\mathbf{k}}|$ vanishes. Invoking Eq. (47), these




FIG. 5. Topological phase diagram for the Bloch two-band nearest-neighbor toy model. We have assumed $\varepsilon > 0$, which is consistent with Fig. 4(b), but the $\varepsilon < 0$ phase diagram may be obtained by negating the Chern numbers. Both the $\mathcal{C} = (-1 : 1)$ and $(0 : 0)$ Chern sectors are realized in the continuum model, as exhibited in Fig. 3.

gap closures are realized at the parameter values

$$\tau/\varepsilon = -1/6, 1/3, 1/2, \quad (48)$$

with the gap closures located at the Γ point, three M points, and two K points, respectively, as shown in the Supplemental Material [84]. Special attention must be paid to the axis $\tilde{\tau}/\varepsilon = 0$ along which the gap is closed for $\tau/\varepsilon \geq 1/3$ and $\tau/\varepsilon \leq -1/6$. Having identified the locations of all gap closures of the two-band toy model, we may compute the Chern numbers numerically. The results are summarized in Fig. 5.

Comparing the Chern number diagram to that which we derived from the continuum model, Fig. 3, we observe that only the $\mathcal{C} = (-1 : 1)$ and $(0 : 0)$ Chern sectors are accessible by tuning the parameters R_0, R, J of the continuum Hamiltonian. Moreover, we remark that the behavior at the boundary between these two Chern sectors is consistent between the continuum and toy models. As exhibited in Fig. 4(b), which displays the hopping parameters obtained from the continuum model at $(J, R_0/a) = (3, 0.49)$, truncating the tight-binding model at nearest-neighbor hopping causes only a small shift in the boundary between the two Chern sectors. Similar agreement is seen at other parameter values (J, R_0) . Finally, the gap closure at this transition is Dirac-like and occurs at the Γ point in both the full tight-binding model and truncated toy model. We caution, however, that longer-range hoppings must be preserved in order to accurately fit to energies away from the Γ point. Finally, given the experimental capacity to simulate magnetic fluxes and the Hofstadter Hamiltonian using atoms in optical lattices [93], we note that it would be interesting to attempt to realize our family of tight-binding models in the ultracold atom setting.

IV. NÉEL SKYRMION LATTICE

In contrast to the Bloch skyrmion case, the Néel in-plane divergence $\nabla^{2D} \cdot \mathbf{n}(\mathbf{r})$ does not vanish identically but only in total when integrated over a single unit cell [67]. Consequently, there exists no gauge transformation which removes the in-plane magnetization component, thereby sacrificing both the particle-hole and chiral symmetries. The bands in the Néel case are therefore not symmetric across zero energy and do not have related Berry curvature.

Nonetheless, computation of the band eigenstates again reveals isolated low-energy bands with nontrivial Chern number. As in the Bloch case, tuning the skyrmion proximity, radial profile, and the effective Hund's coupling leads to a collection of phases distinguished by these Chern numbers. In Fig. 6 we plot the bands for fixed parameters $(J, R_0/a) = (3, 0.3)$ and for three values of the skyrmion size R . Two of these values $R/a = 0.2, 0.29$ sit on the boundaries between topological sectors. In Fig. 6(c) we observe the emergence of particle-hole symmetry. This is due to the Bloch and Néel skyrmions losing their distinction in the limit $R \rightarrow R_0$, with both textures approaching a domain wall droplet with no in-plane magnetization.

In Fig. 7 we display the full topological phase diagram for fixed $R_0/a = 0.3$. Let us continue to label the bands by integers n where $n < 0$ ($n > 0$) denote the bands below (above) half-filling. We find that the bands $n \in \{-1, 1, 2, 3\}$ are consistently gapped from the remaining bands and therefore have well-defined Chern numbers. We denote them by $(\mathcal{C}_{-1} : \mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3)$ with the colon marking half filling. The transition between the $\mathcal{C} = (0 : 0, 1, 1)$ and $(0 : 0, 0, 2)$ sectors upon increasing R and J occurs due to a gap closure at the Γ point, Fig. 6(a). Two further topological sectors are then accessible, including an island of $\mathcal{C} = (0 : 0, 0, -1)$ near the minimal value $R = R_0/2$, as well as a robust Chern-trivial region which persist for large J . We remark that the chosen value of $R_0/a = 0.3$ showcases that Chern bands can be obtained in the Néel case for smaller cutoff radii R_0 and smaller skyrmion core sizes R than in the Bloch case.

V. SUMMARY AND FUTURE DIRECTIONS

We have determined the band structure of TI helical Dirac surface states coupled to skyrmion crystal textures, revealing a strong dependence on the single-skyrmion radial profile n_z which we studied by interpolating continuously between a sinusoidal and domain wall limit. This stands in contrast to previous single-skyrmion [68,69] and skyrmion lattice studies [77,94] in which the textures were approximated by domain wall droplets. The question of tuning the radial profile, or of determining its most stable realization, is likely material dependent and remains open to further investigation.

Significant qualitative differences between Néel and Bloch-type skyrmions were also elucidated. In the latter case, additional particle-hole and chiral symmetries led to constraints on the eigenstates and Berry curvature of the associated bands, producing a rich topological phase diagram upon varying the skyrmion separation, core size, and the spin-fermion coupling. In both cases we demonstrated the appearance of topologically nontrivial bands whose Chern




FIG. 6. Néel skyrmion bands and density of states (DOS) at Hund's coupling $J = 3$ and cutoff radius $R_0/a = 0.30$. Colored red are the bands whose Chern numbers are displayed in Fig. 7, namely the first band below half filling and the three bands above it. (a) Whereas the bands carry Chern number $\mathcal{C} = (0 : 0, 1, 1)$ at the minimum value $R = R_0/2 = 0.15a$ of the core size, increasing R leads to a gap closure (circled) between the $n = 2, 3$ bands. (b) These four bands enter the $\mathcal{C} = (0 : 0, 0, 2)$ Chern sector. The nontrivial $\mathcal{C}_{+3} = 2$ Chern band hovers near the continuum before eventually closing a gap and descending toward zero energy. (c) While the Néel bands generically do not exhibit the particle-hole symmetry that they enjoyed in the Bloch case, we observe its reemergence in the limit $R \rightarrow R_0$ where both skyrmion types approach a domain wall droplet.

numbers depend sensitively on the skyrmion radial profile. This intrinsic contribution to the Hall conductance contrasts with the topological Hall effect observed when nonrelativistic electrons couple to a skyrmion texture [72].

Going beyond the work described in this paper, we expect the coupling between skyrmion texture and fermions to be anisotropic [17], with the coupling to n_z being different from that to n_x, n_y . For Bloch skyrmions, this anisotropy has no impact since the in-plane component of the skyrmion texture can still be gauged away. It may, however, be interesting to investigate the impact of this anisotropy on the bands of Néel skyrmion crystals.

Using band theory techniques, we have constructed localized, symmetric Wannier orbitals for Dirac surface states coupled to a skyrmion crystal. To the best of our knowledge, such Wannier functions have not been extracted in previous

work on this subject. For well-separated skyrmions, we have shown that the Wannier functions are ‘ring’-like states which resemble the previously studied single-skyrmion bound states. Within a two-band study, the truncation of the associated tight-binding model to nearest-neighbor hopping is found to capture the most relevant topological features.

In future studies it could prove fruitful to explore the effects of electron-electron interactions in such lattice models, which could support fractional Chern insulator phases [95] or other correlated states due to the presence of narrow Chern bands at large skyrmion core size and effective Hund's coupling strength. Furthermore, it would be interesting to explore how external magnetic fields tune the density and core size of skyrmions in the lattice, thereby allowing for systematic exploration of the topological phase diagrams discussed above. In this work, we assumed that a weak external magnetic field is sufficient to stabilize skyrmions [96,97]. However, in materials where stronger external fields are needed to stabilize skyrmions, the Landau levels of the helical Dirac surface states could begin to play an important role in the skyrmion-skyrmion interaction as well as the nature of electronic states, leading to distinct topological features deserving of a separate investigation.

Note added. Recently, we came across a related paper [94] which examines the impact of spiral textures on TI Dirac surface states and the energetic stabilization of Néel skyrmions in intrinsically magnetic TIs. Our band theory calculations for Néel and Bloch skyrmion crystals, construction of Wannier orbitals, and tight-binding model results provide a complementary perspective on this topic.

ACKNOWLEDGMENTS

T.P.-B. acknowledges useful discussions with Yafis Barlas, Kunal Tiwari, and William Coish, and funding from NSERC and FRQNT. S.D. and A.P. acknowledge funding from NSERC and useful discussions with Sopheak Sorn. S.D. acknowledges helpful discussions with Shubhayu Chatterjee, Stephen Gant, and Zachary Weinstein. S.D. is supported by the NSERC PGSD fellowship. H.L. thanks Richard Man-Wai Ling for his unfailing and enduring support.




FIG. 7. $R_0/a = 0.3$. Chern number phase diagram in the Néel skyrmion case for a single band below half filling and three bands above half filling. The Chern numbers in each tuple \mathcal{C} are listed in order of increasing energy, with the colon marking half filling. The marked points correspond to the band-structure plots presented in Fig. 6.

- [1] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [2] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [3] N. P. Armitage, E. J. Mele, and A. Vishwanath, *Rev. Mod. Phys.* **90**, 015001 (2018).
- [4] L. Fu, C. L. Kane, and E. J. Mele, *Phys. Rev. Lett.* **98**, 106803 (2007).
- [5] J. E. Moore and L. Balents, *Phys. Rev. B* **75**, 121306(R) (2007).
- [6] R. Roy, *Phys. Rev. B* **79**, 195321 (2009).
- [7] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, *Phys. Rev. B* **78**, 195424 (2008).
- [8] G. Dresselhaus and M. S. Dresselhaus, *Phys. Rev.* **140**, A401 (1965).
- [9] F. D. M. Haldane, *Phys. Rev. Lett.* **61**, 2015 (1988).
- [10] X.-F. Wang and T. Chakraborty, *Phys. Rev. B* **75**, 033408 (2007).
- [11] T. Thonhauser and D. Vanderbilt, *Phys. Rev. B* **74**, 235111 (2006).
- [12] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 226801 (2005).
- [13] D. Xiao, W. Yao, and Q. Niu, *Phys. Rev. Lett.* **99**, 236809 (2007).
- [14] C.-X. Liu, S.-C. Zhang, and X.-L. Qi, *Annu. Rev. Condens. Matter Phys.* **7**, 301 (2016).
- [15] R. Jackiw and C. Rebbi, *Phys. Rev. D* **13**, 3398 (1976).
- [16] G. W. Semenoff, V. Semenoff, and F. Zhou, *Phys. Rev. Lett.* **101**, 087204 (2008).
- [17] R. Wakatsuki, M. Ezawa, and N. Nagaosa, *Sci. Rep.* **5**, 13638 (2015).
- [18] A. Isacsson, L. M. Jonsson, J. M. Kinaret, and M. Jonson, *Phys. Rev. B* **77**, 035423 (2008).
- [19] C.-H. Park, L. Yang, Y.-W. Son, M. L. Cohen, and S. G. Louie, *Nat. Phys.* **4**, 213 (2008).
- [20] M. Barbier, P. Vasilopoulos, and F. M. Peeters, *Phys. Rev. B* **81**, 075438 (2010).
- [21] F. Guinea and T. Low, *Philos. Trans. R. Soc., A* **368**, 5391 (2010).
- [22] P. Bursat, A. L. Yeyati, L. Brey, and H. A. Fertig, *Phys. Rev. B* **83**, 195434 (2011).
- [23] S. Dubey, V. Singh, A. K. Bhat, P. Parikh, S. Grover, R. Sensarma, V. Tripathi, K. Sengupta, and M. M. Deshmukh, *Nano Lett.* **13**, 3990 (2013).
- [24] M. Killi, S. Wu, and A. Paramekanti, *Phys. Rev. Lett.* **107**, 086801 (2011).
- [25] S. Wu, M. Killi, and A. Paramekanti, *Phys. Rev. B* **85**, 195404 (2012).
- [26] R. Bistritzer and A. H. MacDonald, *Proc. Natl. Acad. Sci.* **108**, 12233 (2011).
- [27] J. M. B. Lopes dos Santos, N. M. R. Peres, and A. H. Castro Neto, *Phys. Rev. B* **86**, 155449 (2012).
- [28] L. Zou, H. C. Po, A. Vishwanath, and T. Senthil, *Phys. Rev. B* **98**, 085435 (2018).
- [29] W. Yan, M. Liu, R.-F. Dou, L. Meng, L. Feng, Z.-D. Chu, Y. Zhang, Z. Liu, J.-C. Nie, and L. He, *Phys. Rev. Lett.* **109**, 126801 (2012).
- [30] E. J. Mele, *Phys. Rev. B* **81**, 161405(R) (2010).
- [31] J. M. B. Lopes dos Santos, N. M. R. Peres, and A. H. Castro Neto, *Phys. Rev. Lett.* **99**, 256802 (2007).
- [32] G. Trambly de Laissardière, D. Mayou, and L. Magaud, *Nano Lett.* **10**, 804 (2010).
- [33] S. Shallicross, S. Sharma, E. Kandelaki, and O. A. Pankratov, *Phys. Rev. B* **81**, 165105 (2010).
- [34] E. Suárez Morell, J. D. Correa, P. Vargas, M. Pacheco, and Z. Barticevic, *Phys. Rev. B* **82**, 121407(R) (2010).
- [35] A. Luican, G. Li, A. Reina, J. Kong, R. R. Nair, K. S. Novoselov, A. K. Geim, and E. Y. Andrei, *Phys. Rev. Lett.* **106**, 126802 (2011).
- [36] T. Ohta, J. T. Robinson, P. J. Feibelman, A. Bostwick, E. Rotenberg, and T. E. Beechem, *Phys. Rev. Lett.* **109**, 186807 (2012).
- [37] G. Li, A. Luican, J. M. B. Lopes dos Santos, A. H. Castro Neto, A. Reina, J. Kong, and E. Y. Andrei, *Nat. Phys.* **6**, 109 (2010).
- [38] Y. Cao, J. Y. Luo, V. Fatemi, S. Fang, J. D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras, and P. Jarillo-Herrero, *Phys. Rev. Lett.* **117**, 116804 (2016).
- [39] P. J. Ledwith, G. Tarnopolsky, E. Khalaf, and A. Vishwanath, *Phys. Rev. Research* **2**, 023237 (2020).
- [40] J. Jung, A. Raoux, Z. Qiao, and A. H. MacDonald, *Phys. Rev. B* **89**, 205414 (2014).
- [41] D. Xiao, G.-B. Liu, W. Feng, X. Xu, and W. Yao, *Phys. Rev. Lett.* **108**, 196802 (2012).
- [42] F. Wu, T. Lovorn, E. Tutuc, and A. H. MacDonald, *Phys. Rev. Lett.* **121**, 026402 (2018).
- [43] F. Wu, T. Lovorn, E. Tutuc, I. Martin, and A. H. MacDonald, *Phys. Rev. Lett.* **122**, 086402 (2019).
- [44] Y. Tang, L. Li, T. Li, Y. Xu, S. Liu, K. Barmak, K. Watanabe, T. Taniguchi, A. H. MacDonald, J. Shan *et al.*, *Nature (London)* **579**, 353 (2020).
- [45] E. C. Regan, D. Wang, C. Jin, M. I. Bakti Utama, B. Gao, X. Wei, S. Zhao, W. Zhao, Z. Zhang, K. Yumigeta *et al.*, *Nature (London)* **579**, 359 (2020).
- [46] Z. Zhang, Y. Wang, K. Watanabe, T. Taniguchi, K. Ueno, E. Tutuc, and B. J. LeRoy, *Nat. Phys.* **16**, 1093 (2020).
- [47] L. Wang, E.-M. Shih, A. Ghiotto, L. Xian, D. A. Rhodes, C. Tan, M. Claassen, D. M. Kennes, Y. Bai, B. Kim *et al.*, *Nat. Mater.* **19**, 861 (2020).
- [48] Y. Xu, S. Liu, D. A. Rhodes, K. Watanabe, T. Taniguchi, J. Hone, V. Elser, K. F. Mak, and J. Shan, *Nature (London)* **587**, 214 (2020).
- [49] T. Wang, N. F. Q. Yuan, and L. Fu, *Phys. Rev. X* **11**, 021024 (2021).
- [50] J. Cano, S. Fang, J. H. Pixley, and J. H. Wilson, *Phys. Rev. B* **103**, 155157 (2021).
- [51] M. Dzero, J. Xia, V. Galitski, and P. Coleman, *Annu. Rev. Condens. Matter Phys.* **7**, 249 (2016).
- [52] D. K. Efimkin and V. Galitski, *Phys. Rev. B* **90**, 081113(R) (2014).
- [53] S. Wolgast, Y. S. Eo, T. Öztürk, G. Li, Z. Xiang, C. Tinsman, T. Asaba, B. Lawson, F. Yu, J. W. Allen, K. Sun, L. Li, C. Kurdak, D. J. Kim, and Z. Fisk, *Phys. Rev. B* **92**, 115110 (2015).
- [54] Y. Nakajima, P. Syers, X. Wang, R. Wang, and J. Paglione, *Nat. Phys.* **12**, 213 (2016).
- [55] K. L. Tiwari, W. A. Coish, and T. Pereg-Barnea, *Phys. Rev. B* **96**, 235120 (2017).
- [56] H. Yang, A. Thiaville, S. Rohart, A. Fert, and M. Chshiev, *Phys. Rev. Lett.* **115**, 267210 (2015).
- [57] C. Moreau-Luchaire, C. Moutafis, N. Reyren, J. Sampaio, C. A. F. Vaz, N. Van Horne, K. Bouzehouane, K. Garcia, C. Deranlot, P. Warnicke *et al.*, *Nat. Nanotechnol.* **11**, 444 (2016).

- [58] Y.-J. Hao, P. Liu, Y. Feng, X.-M. Ma, E. F. Schwier, M. Arita, S. Kumar, C. Hu, R. Lu, M. Zeng, Y. Wang, Z. Hao, H. Y. Sun, K. Zhang, J. Mei, N. Ni, L. Wu, K. Shimada, C. Chen, Q. Liu, and C. Liu, *Phys. Rev. X* **9**, 041038 (2019).
- [59] B. Li, J.-Q. Yan, D. M. Pajerowski, E. Gordon, A.-M. Nedić, Y. Sizyuk, L. Ke, P. P. Orth, D. Vaknin, and R. J. McQueeney, *Phys. Rev. Lett.* **124**, 167204 (2020).
- [60] Q. Liu, C.-X. Liu, C. Xu, X.-L. Qi, and S.-C. Zhang, *Phys. Rev. Lett.* **102**, 156603 (2009).
- [61] K. Yasuda, R. Wakatsuki, T. Morimoto, R. Yoshimi, A. Tsukazaki, K. S. Takahashi, M. Ezawa, M. Kawasaki, N. Nagaosa, and Y. Tokura, *Nat. Phys.* **12**, 555 (2016).
- [62] J. Jiang, D. Xiao, F. Wang, J.-H. Shin, D. Andreoli, J. Zhang, R. Xiao, Y.-F. Zhao, M. Kayyalha, L. Zhang *et al.*, *Nat. Mater.* **19**, 732 (2020).
- [63] P. Li, J. Ding, S. S.-L. Zhang, J. Kally, T. Pillsbury, O. G. Heinonen, G. Rimal, C. Bi, A. DeMann, S. B. Field *et al.*, *Nano Lett.* **21**, 84 (2020).
- [64] X. Z. Yu, Y. Onose, N. Kanazawa, J. H. Park, J. H. Han, Y. Matsui, N. Nagaosa, and Y. Tokura, *Nature (London)* **465**, 901 (2010).
- [65] F. S. Nogueira, I. Eremin, F. Katmis, J. S. Moodera, J. van den Brink, and V. P. Kravchuk, *Phys. Rev. B* **98**, 060401(R) (2018).
- [66] D. Mandler, P. Kotetes, and G. Schön, *Phys. Rev. B* **91**, 155405 (2015).
- [67] K. Nomura and N. Nagaosa, *Phys. Rev. B* **82**, 161401(R) (2010).
- [68] H. M. Hurst, D. K. Efimkin, J. Zang, and V. Galitski, *Phys. Rev. B* **91**, 060401(R) (2015).
- [69] Y. Araki and K. Nomura, *Phys. Rev. B* **96**, 165303 (2017).
- [70] C.-Z. Wang, H.-Y. Xu, and Y.-C. Lai, *Phys. Rev. Research* **2**, 013247 (2020).
- [71] K. L. Tiwari, J. Lavoie, T. Pereg-Barnea, and W. A. Coish, *Phys. Rev. B* **100**, 125414 (2019).
- [72] A. Neubauer, C. Pfleiderer, B. Binz, A. Rosch, R. Ritz, P. G. Niklowitz, and P. Böni, *Phys. Rev. Lett.* **102**, 186602 (2009).
- [73] M. Lee, W. Kang, Y. Onose, Y. Tokura, and N. P. Ong, *Phys. Rev. Lett.* **102**, 186601 (2009).
- [74] K. Hamamoto, M. Ezawa, and N. Nagaosa, *Phys. Rev. B* **92**, 115417 (2015).
- [75] B. Göbel, A. Mook, J. Henk, and I. Mertig, *Phys. Rev. B* **95**, 094413 (2017).
- [76] S. Sorn, S. Divic, and A. Paramekanti, *Phys. Rev. B* **100**, 174411 (2019).
- [77] J. L. Lado and J. Fernández-Rossier, *Phys. Rev. B* **92**, 115433 (2015).
- [78] J. Liu, A. Singh, B. Kuerbanjiang, C. H. W. Barnes, and T. Hesjedal, *Nanotechnology* **31**, 434001 (2020).
- [79] G. Kimbell, P. M. Sass, B. Woltjes, E. K. Ko, T. W. Noh, W. Wu, and J. W. A. Robinson, *Phys. Rev. Materials* **4**, 054414 (2020).
- [80] G. Kim, K. Son, Y. E. Suyolcu, L. Miao, N. J. Schreiber, H. P. Nair, D. Putzky, M. Minola, G. Christiani, P. A. van Aken, K. M. Shen, D. G. Schlom, G. Logvenov, and B. Keimer, *Phys. Rev. Materials* **4**, 104410 (2020).
- [81] F. M. Bartram, S. Sorn, Z. Li, K. Hwangbo, S. Shen, F. Frontini, L. He, P. Yu, A. Paramekanti, and L. Yang, *Phys. Rev. B* **102**, 140408(R) (2020).
- [82] S. Meyer, M. Perini, S. von Malottki, A. Kubetzka, R. Wiesendanger, K. von Bergmann, and S. Heinze, *Nat. Commun.* **10**, 3823 (2019).
- [83] Y. Guang, I. Bykova, Y. Liu, G. Yu, E. Goering, M. Weigand, J. Gräfe, S. K. Kim, J. Zhang, H. Zhang, *et al.*, *Nat. Commun.* **11**, 949 (2020).
- [84] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.105.035156> for derivations of the action of various symmetries and further details concerning the Wannier gauge transformation and tight-binding model.
- [85] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, *Rev. Mod. Phys.* **88**, 035005 (2016).
- [86] J. H. Han, *Skyrmions in condensed matter* (Springer, Cham, 2017).
- [87] T. Fukui, Y. Hatsugai, and H. Suzuki, *J. Phys. Soc. Jpn.* **74**, 1674 (2005).
- [88] C. Brouder, G. Panati, M. Calandra, C. Mourougane, and N. Marzari, *Phys. Rev. Lett.* **98**, 046402 (2007).
- [89] N. Marzari and D. Vanderbilt, *Phys. Rev. B* **56**, 12847 (1997).
- [90] N. Marzari, A. A. Mostofi, J. R. Yates, I. Souza, and D. Vanderbilt, *Rev. Mod. Phys.* **84**, 1419 (2012).
- [91] A. M. Cook, C. Hickey, and A. Paramekanti, *Phys. Rev. B* **90**, 085145 (2014).
- [92] S. Baidya, U. V. Waghmare, A. Paramekanti, and T. Saha-Dasgupta, *Phys. Rev. B* **94**, 155405 (2016).
- [93] M. Aidelsburger, M. Atala, M. Lohse, J. T. Barreiro, B. Paredes, and I. Bloch, *Phys. Rev. Lett.* **111**, 185301 (2013).
- [94] N. Paul and L. Fu, *Phys. Rev. Research* **3**, 033173 (2021).
- [95] N. Regnault and B. A. Bernevig, *Phys. Rev. X* **1**, 021014 (2011).
- [96] A. Soumyanarayanan, M. Raju, A. L. Gonzalez Oyarce, A. K. C. Tan, M.-Y. Im, A. P. Petrović, P. Ho, K. H. Khoo, M. Tran, C. K. Gan *et al.*, *Nat. Mater.* **16**, 898 (2017).
- [97] A. Chacon, L. Heinen, M. Halder, A. Bauer, W. Simeth, S. Mühlbauer, H. Berger, M. Garst, A. Rosch, and C. Pfleiderer, *Nat. Phys.* **14**, 936 (2018).

Correction: The first sentence of the caption to Figure 6 contained a typographical error and has been fixed.