

Substrate chemistry distributions — cluster extremes
Top 15 most CHEMICALLY UNIFORM clusters
Glucocorticoid receptor (n=2, median=0.74)
Glucosylceramidase (n=4, median=0.72)
Microsomal triglyceride transfer protein large subunit (n=3, median=0.70)
Hedgehog protein (n=2, median=0.66)
Hormone-sensitive lipase (n=3, median=0.66)
Apolipoprotein A-I (n=2, median=0.66)
Hedgehog protein (n=2, median=0.66)
Bile acid receptor (n=2, median=0.66)
Acyl-coenzyme A diphosphatase NUDT19 (n=2, median=0.66)
Glucosylceramidase (n=7, median=0.64)
Bile salt export pump (n=10, median=0.61)
Multidrug resistance-associated protein 4 (n=9, median=0.60)
G-protein coupled estrogen receptor 1 (n=10, median=0.59)
Acyl-coenzyme A diphosphatase NUDT19 (n=17, median=0.57)
Hormone-sensitive lipase (n=8, median=0.53)
random-pair median = 0.20
0.0
0.2
0.4
0.6
0.8
1.0
Pairwise Tanimoto (ECFP4)
Top 15 most CHEMICALLY DIVERSE clusters
(3R)-3-hydroxyacyl-CoA dehydrogenase (n=17, median=0.17)
G-protein coupled estrogen receptor 1 (n=27, median=0.18)
Bile acid receptor (n=29, median=0.19)
Sterol carrier protein 2 (n=5, median=0.19)
Glucosylceramidase (n=20, median=0.19)
Glucosylceramidase (n=20, median=0.19)
Hormone-sensitive lipase (n=39, median=0.20)
Hormone-sensitive lipase (n=22, median=0.20)
Non-lysosomal glucosylceramidase (n=30, median=0.20)
11-beta-hydroxysteroid dehydrogenase 1 (n=87, median=0.21)
Oxysterol-binding protein-related protein 8 (n=65, median=0.21)
3-hydroxyacyl-CoA dehydrogenase type-2 (n=108, median=0.21)
Cytochrome P450 3A (n=42, median=0.21)
Carboxylic ester hydrolase (n=11, median=0.22)
Cholesterol 24-hydroxylase (n=44, median=0.22)
0.0
0.2
0.4
0.6
0.8
1.0
Pairwise Tanimoto (ECFP4)