

ASSUMING SCHINZEL'S HYPOTHESIS H: A SIX-UNIVERSAL-QUANTIFIER DEFINITION OF $\mathbb{Z}$ INSIDE $\mathbb{Q}$

H10Q PROJECT

ABSTRACT. Assuming classical Schinzel's Hypothesis H, we prove that $\mathbb{Z}$ has a definition in $\mathbb{Q}$ with six universal quantifiers and that $\text{efd}_{\mathbb{Q}}(\mathbb{Q} \setminus \mathbb{Z}) \leq 5$. This is a conditional result, not an unconditional solution of Hilbert's Tenth Problem over $\mathbb{Q}$. For every cell (w, z) , the class construction fixes

$$\tau^\dagger = \frac{1 + 2a^2}{1 + 4a^2}, \quad \delta_{\tau^\dagger} = -\frac{4a^4}{1 + 4a^2},$$

uses $f = w$, and proves normalization, coprimality, absence of a fixed divisor, frozen local data, and positivity uniformly.

The last non-Schinzel obligation is removed by a fixed-fiber theorem. For every fixed $Z \in \mathbb{Q}^\times$, equal endpoints and Newton polygons at A , $a = 0$, and $a = \infty$ eliminate every possible factor of

$$H = \frac{A}{4}P = a^8 Z^4 N_g^2 + 4A^3 D^2 b^4 - 2A^4 (a - 1)^2 D^2 b^5.$$

The degree-four case would force $Z^4 = 1024(1 - Z)^2$, whose two quadratic branches have nonsquare discriminants 1152 and 896; the degenerate value $Z = 1$ has an exact degree-eight certificate modulo 11. Hence $P(a, Z)$ is irreducible in $\mathbb{Q}(a)[b]$ for every nonzero rational Z .

Quantitative Hilbert irreducibility selects a good a inside the existing character-admissible progression. The L20 class and the L19 fixed-pair implication then reduce member existence exactly to classical Schinzel H; the resulting simultaneous prime values give an emergent-free member and activate L6. An independent reciprocal-trace and complete 2-isogeny-descent proof at $a = 1$ corroborates the vertical theorem; that $a = 1$ is a specialization witness, not the selected class parameter. Thus Schinzel H is the sole conjectural input and no per-cell irreducibility hypothesis remains.

The L23 half-sieve gives, unconditionally, $\gg X/(\log X)^{3/2}$ prime-sequence candidates clean at every bad prime below $X^{49/100}$. These are small-prime-clean members of the progression, not full emergent-free members: degree-eight values may retain up to 16 large factors, and reciprocity leaves $0, 2, \dots, 16$ bad ones. L24 proves that both proposed *unscaled* self-coupled diagonal orientations are Φ -empty over $\mathbb{Q}_2$. L25 shows that fixed scaling removes this universal dyadic wall but leaves parity and target compatibility in those formulas. L26 gives an even-pullback reciprocal tie with a quartic trace character. L28 proves uniformly that the trace quartic and its bad-sign Capell cover are irreducible over $\mathbb{Q}_2$. L29 completely classifies the distinct reciprocal lift and gives a target-preserving dyadic compressor. L27 gives one triangular shear with complete dyadic and aligned odd-target local points. L30 then replaces its degree-eight cover by the constant-two even quartic, proves the same complete displayed local data, and establishes broad square-branch/shear rigidity. L31 proves one exact globally good fixed-field reciprocal member at $w = 13$. It also finds an exact rational constant-two point immediately before the mandatory cube pullback, reduces that cube condition to two genus-2 curves, and identifies AP1 as the exact parity replacement for Schinzel H. No uniform member theorem, controlled cube-compatible quartic point, AP1 theorem, or five-count improvement is claimed, and the conditional theorem is unchanged.

CONTENTS

1. Introduction	3
1.1. The record chain and its audit	3
1.2. Contributions and scope	4
2. Model-field architecture: selection, witness tying, and the open norm problem	5
2.1. The finite-field selector is exact	5

Date: Working draft, 2026-08-23 — not submitted.

2.2.	The tied model and its six-variable bookkeeping	5
2.3.	Local structure and the exact global obstruction	6
2.4.	Structural corollaries and the remaining lemma	7
3.	Obstruction walls and reciprocity steering	7
3.1.	The alignment wall	7
3.2.	Reciprocity steering	8
3.3.	The generalized coprimality wall	9
3.4.	Composite squarefree wall	9
4.	Class Dichotomy, Characterization, and Escapes	10
4.1.	The L10 dichotomy	10
4.2.	Free branches, an immovable wall	10
4.3.	Non-coprime escapes and complete grid coverage	11
4.4.	Why no square-class shortcut remains	12
5.	Uniform existence of an aligned class	12
6.	The verified quantitative layer	14
6.1.	L14: instance verification	14
6.2.	L15: the exact remainder and counting laws	14
6.3.	L16: why the small layer has sieve shape	15
6.4.	L17: matched validation	15
6.5.	L17 cofactor layer: reciprocity and its limit	16
7.	The conditional record and the Schinzel audit	16
7.1.	The statement, with its hypothesis exposed	16
7.2.	What is proved, and what Schinzel must supply	17
7.3.	PROVED finite audit of the Schinzel pair	17
7.4.	Residual certificates and the literature boundary	18
8.	Schinzel's Hypothesis H: the fixed-class implication	19
8.1.	The operational contract	19
8.2.	The theorem	19
8.3.	Uniform admissibility status	19
9.	Fixed-branch irreducibility for every vertical fiber	20
9.1.	The off-branch warning	21
9.2.	Endpoint and Newton elimination	21
9.3.	Hilbert selection inside the admissible progression	23
9.4.	Independent reciprocal-trace corroboration	23
9.5.	Route discipline	24
10.	Unconditional density, algebraic walls, and the off-grid frontier	24
10.1.	An unconditional local bound	24
10.2.	The L23 half-sieve and its exact parity wall	25
10.3.	The L23 fibration and norm walls	26
10.4.	The L24 diagonal route is refuted	26
10.5.	The L25 scaled couplings are not dyadically killed	27
10.6.	Two new non-diagonal constructions	28
10.7.	The off-grid horizon	32
11.	Conclusion and outlook	33
	Reproducibility	34
	References	35

1. INTRODUCTION

Hilbert’s Tenth Problem over $\mathbb{Q}$ asks for an algorithm which, given a finite system of polynomials with rational coefficients, decides whether the system has a common rational zero. The problem remains open in both directions. One standard route to undecidability would be an existential definition of $\mathbb{Z}$ in $\mathbb{Q}$, since such a definition would transport the undecidability over $\mathbb{Z}$ to rational points. The converse is not known: failure of that route would not imply decidability over $\mathbb{Q}$.

There is already a sharp first obstruction. Cornelissen–Zahidi proved that $\mathbb{Z}$ cannot be the projection of the rational points of a curve over $\mathbb{Q}$; in particular, no polynomial with one existential witness defines $\mathbb{Z}$ in $\mathbb{Q}$ [4]. Thus two witnesses, equivalently a surface as witness variety, form the first open existential case. This is also exactly where the available topological and thin-set lower-bound methods stop. A different elliptic-curve route has its own cliff. If only finitely many primes are withheld from an S -integer ring, the indices n for which $x(nP)$ is integral are a finite union of congruence classes, relative to the chosen integral Weierstrass model. When nonempty this set has positive density and is archimedeanly equidistributed on a real component, rather than discrete. Consequently the prime-index mechanism used in elliptic Diophantine models degenerates at the cofinite- S boundary; this statement does not rule out unrelated predicates.

Koenigsmann changed the quantitative landscape by proving that $\mathbb{Z}$ is purely universally definable in $\mathbb{Q}$ [10]. Complementation turns this into the study of the existential rank $\text{rk}_{\mathbb{Q}}^{\exists}(\mathbb{Q} \setminus \mathbb{Z})$. Daans–Dittmann–Fehm then supplied the geometric interpretation: for a non-quantifier-free Diophantine set D over $\mathbb{Q}$,

$$\text{rk}_{\mathbb{Q}}^{\exists}(D) = \text{efd}_{\mathbb{Q}}(D) + 1,$$

and $\text{efd}_{\mathbb{Q}}(D) \leq d$ precisely when D is the image of a morphism all of whose fibres have dimension at most d [6]. The quantifier race is therefore a fibre-dimension problem. In particular, a six-universal-quantifier definition would amount to the bound $\text{efd}_{\mathbb{Q}}(\mathbb{Q} \setminus \mathbb{Z}) \leq 5$, not merely to more efficient logical notation.

1.1. The record chain and its audit. The refereed benchmark is Daans’s ten-quantifier universal definition [5]. Our source-level audit reconstructs its budget before applying any fusion: two quaternion parameters, three witnesses for the auxiliary Φ condition, one splitting witness, and two three-variable trace-norm certificates give twelve variables. Two uses of the intersection theorem of Daans–Dittmann–Fehm save one coordinate each, giving ten; pairwise reassociation cannot improve that proof shape. The same audit isolates the real places to seek an improvement: stronger simultaneous fusion, a cheaper trace-norm primitive, or a joint treatment of the nested valuation-ring conditions.

Sun subsequently announced an unrefereed seven-quantifier construction [16]. It retains Daans’s bridge but replaces the six-witness intersection block by a finite disjunction of three-witness quadrics, and then uses one further fusion. Our A2 audit checked the exact quaternion identities, the block’s soundness, the residue-field combinatorics, and explicit bridge certificates. It also traced the local–global completeness argument through parameterized Hensel lifting, weak and norm approximation, and Hasse–Minkowski. No substantive missing implication was found, but the posted preprint has stale cross-references at the decisive step, and its local–global chain was hand-traced rather than machine-verified. We therefore call seven an *unrefereed announcement*; ten remains the refereed record.

PROVED. The audit also corrects one ingredient of the unrefereed announcement [16]. Its finite-field selection lemma assumes a residue field larger than 25. Exhaustion first showed that the small fields do not fail; more conceptually, we proved the exact identity

$$N(\tau) = \frac{\#E_{\tau^{-2}}(\mathbb{F}_q)}{4}, \quad E_{\tau^{-2}} : y^2 = x(x-1)(x-\tau^{-2}).$$

Hasse's bound makes this positive over every odd finite field admitting an admissible τ . Hence the threshold can be deleted: $\mathbb{F}_3$ is exceptional only because it has no $\tau \notin \{0, \pm 1\}$, while the preprint's other, data-dependent exceptional places remain untouched.

1.2. Contributions and scope. The contributions of this paper are deliberately separated by status.

- (1) **CONDITIONAL ON CLASSICAL SCHINZEL H — a six-quantifier definition.** We tie a witness in Sun's quadratic block to Daans's congruence parameter by $a = 1 + 2s$. Soundness, denominator clearing, the exact target-place selector, the global norm obstruction, canonical-branch irreducibility, and every Schinzel-pair admissibility condition are proved. Assuming classical Schinzel's Hypothesis H, $\mathbb{Q} \setminus \mathbb{Z}$ has existential rank at most six, $\mathbb{Z}$ has a six-universal-quantifier definition, and $\text{efd}_{\mathbb{Q}}(\mathbb{Q} \setminus \mathbb{Z}) \leq 5$.
- (2) **PROVED — wall theorems.** We prove the two-adic source–target alignment wall, show that branch choice cannot remove the reciprocity obstruction, and prove the generalized wall: every admissible b whose odd support is coprime to the cell data fails. The only available door is controlled non-coprimality.
- (3) **PROVED — complete class characterization.** For the prime- b family, on every branch, an aligned class exists exactly when the numerator of z has a prime divisor congruent to 1 modulo 4. The necessity holds for every admissible rational a , not only when $1 + 4a^2$ is prime; the sufficiency is constructive.
- (4) **PROVED — uniform class existence.** For every cell, choose the fixed numerator prime $f = w$ and an odd a with $\left(\frac{1+4a^2}{w}\right) = -1$. A nonempty explicit CRT system and Dirichlet's theorem then give infinitely many square-branch aligned classes. Class existence is therefore no longer part of the global hypothesis.
- (5) **PROVED — exhaustive finite verification.** On the declared grid, every one of the 353 cells has a replayable soluble witness, and every one of the 293 aligned classes has a certified emergent-free member. These are bounded, per-row theorems under the proven-primality engine, not an extrapolation to all primes.
- (6) **EVIDENCE — a mechanistic sieve with matched validation.** Exact root lifting turns each recurring small-prime obstruction into an odd-valuation p -adic mass. On the identical 8,760-member sample, the matched marginal ratio is 0.9993, the hit-conditioned union residual is -2.9 , and a marginal-preserving permutation gives $z = -0.09$. These data validate the small-layer mechanism in the measured window; they do not supply a tail bound or a uniform density theorem.
- (7) **PROVED — fixed-fiber canonical-branch irreducibility.** Blanket irreducibility is false on the explicit $a = 1$, square- δ locus, but the fixed branch

$$\tau^\dagger = \frac{1 + 2a^2}{1 + 4a^2}, \quad \delta_{\tau^\dagger} = -\frac{4a^4}{1 + 4a^2}$$

avoids it. For every fixed $Z \in \mathbb{Q}^\times$, the endpoint–Newton elimination proves $P(a, Z)$ irreducible in $\mathbb{Q}(a)[b]$. Quantitative Hilbert irreducibility then places a good a in the existing character-admissible progression. An independent reciprocal proof uses $a = 1$ only as a specialization witness, not as the selected class parameter.

- (8) **PROVED — off-grid horizon extension.** The same certificate machinery now closes and replays $[179, [-1, 1]]$, the last searched open cell. Every off-grid cell tried to date has a certified member, but this remains a finite instance statement, not a uniform member theorem.

What is not claimed. The chain is exact: fixed- Z vertical irreducibility gives Hilbert selection in the admissible a progression; the L20 class certificate has $f = w$ and all pair conditions; the L19 fixed-pair implication turns classical Schinzel H into an emergent-free member; and L6 yields the

definition. Thus Schinzel H is the sole conjectural input. We claim *no new unconditional quantifier record*: Daans's ten remains the refereed record, Sun's seven remains an unrefereed announcement, and Hilbert's Tenth Problem over $\mathbb{Q}$ remains open.

2. MODEL-FIELD ARCHITECTURE: SELECTION, WITNESS TYING, AND THE OPEN NORM PROBLEM

2.1. The finite-field selector is exact.

Theorem 2.1 (PROVED: L4, the small-field exhaustion). *Let q be an odd prime power with $q \leq 25$, let χ be the quadratic character of $\mathbb{F}_q$ with $\chi(0) = 0$, and let $\tau \in \mathbb{F}_q \setminus \{0, \pm 1\}$. There is an $a \in \mathbb{F}_q$ such that*

$$\chi(1 + 4a^2) = -1, \quad \chi(-(1 + 4a^2)(1 - \tau^2(1 + 4a^2))) = 1.$$

For $q = 3$ the assertion is vacuous, because there is no admissible τ . For $q \in \{5, 7, 9, 11, 13\}$ the minimum number of choices is 2, and for $q \in \{17, 19, 23, 25\}$ it is 4.

Thus the constant 25 in the unrefereed character-sum proof of [16] is not a genuine small-field obstruction: among odd residue fields, only the field of order 3 remains, and there the issue is admissibility rather than failure of the selector.

Theorem 2.2 (PROVED: L5, the Legendre identity). *For every odd prime power q , every $\tau \in \mathbb{F}_q \setminus \{0, \pm 1\}$, and $\lambda = \tau^{-2}$, let $E_\lambda : y^2 = x(x-1)(x-\lambda)$. Then*

$$N(\tau) := \#\{a \in \mathbb{F}_q : \chi(1 + 4a^2) = -1, \chi(-(1 + 4a^2)(1 - \tau^2(1 + 4a^2))) = 1\} = \frac{\#E_\lambda(\mathbb{F}_q)}{4}.$$

Consequently

$$\#E_\lambda(\mathbb{F}_q) \geq (\sqrt{q} - 1)^2 > 0, \quad N(\tau) \geq (\sqrt{q} - 1)^2/4 > 0,$$

so the hypothesis $|k| > 25$ can be deleted from the lemma.

The proof is an exact point count, not an asymptotic substitution. With $u_a = 1 + 4a^2$ and $w_a = -u_a(1 - \tau^2 u_a)$, expand the corrected indicator; the hyperbola identity $\sum_b \chi(b^2 + c) = -1$ for $c \neq 0$ evaluates the quadratic pieces. Fibre-counting the map $a \mapsto u_a$ then gives

$$4N = q + 1 + \sum_u \chi(u(u-1)(u-\lambda)) = \#E_\lambda(\mathbb{F}_q).$$

This also explains divisibility by 4: the Legendre curve has its full rational 2-torsion.

2.2. The tied model and its six-variable bookkeeping. Set $K = \mathbb{Q}$, $S = \{2\}$, $\pi = 2$, and $u = 1$. Pull the bridge back along $z \mapsto z^3$ and put $A = 1 + 4a^2$, $B = 2b$, $\delta_\tau = 1 - A\tau^2$, and $c = h(a, b, z^3) = a^2 z^6 g(a, b)/(1 - z^3 - a^2 z^6)$. The original block is

$$\Psi_\tau(a, b, c) : \exists y, r, s [\delta_\tau(c^2 - Ay^2) - 16B(r^2 - As^2) = 16].$$

The architectural step is to identify its witness s with the congruence parameter by imposing $a = 1 + 2s$. We use the finite branch menu

$$\tau \in \left\{ 0, \frac{2a}{A}, \tau^\dagger = \frac{1 + 2a^2}{A} \right\}.$$

After clearing the nonzero denominators, the resulting two-witness block is

$$\begin{aligned} \Theta_*(a, b, c, s) : \quad & \exists y, r \quad [c^2 - Ay^2 - 16Br^2 = 16 - 16ABs^2] \\ & \vee [c^2 - Ay^2 - 16ABr^2 = 16A - 16A^2Bs^2] \\ & \vee [a^4(c^2 - Ay^2) + 4ABr^2 = 4A^2Bs^2 - 4A]. \end{aligned}$$

The three equations correspond respectively to $\tau = 0$, $\tau = 2a/A$, and $\tau^\dagger$; on the last branch $\delta_{\tau^\dagger} = -4a^4/A$ and $\alpha = (2a^2)^2$ is a square. A finite disjunction is represented by the product of

these polynomials in the same witnesses (y, r) , so adding the square branch costs no variable. The candidate model is therefore

$$F(z) := \exists b, s \left[(1 + 2s, b) \in \Phi_1^{\{2\}} \wedge \Theta_*(1 + 2s, b, h(1 + 2s, b, z^3), s) \right].$$

Here $\Phi_1^{\{2\}}$ enforces $v_2(s) \geq 0$ and $b \in \mathcal{O}_2^\times$. Membership in it has existential rank at most 3; the tied block has rank 2 in (y, r) . The intersection theorem of [6] replaces their conjunction by $3 + 2 - 1 = 4$ witnesses, and projection of the two base coordinates (b, s) gives

$$2 + (3 + 2 - 1) = 6.$$

If an input set has rank zero, the elementary conjunction bound is no larger. The two branches share (y, r) and are combined by multiplying their equations; denominator clearing introduces no inverse witness because $A \equiv 5 \pmod{8}$ at 2 and $1 - z^3 - a^2 z^6$ cannot vanish over $\mathbb{Q}$.

For use by the later local model, write $Z = z^3$, $D_z = 1 - z^3 - a^2 z^6$, and $N_g(b) = 16a^4 b^2 - A(b-1)^4$. The denominator-cleared value polynomial is

$$P(b) = 16D_z^2 A^2 b^4 - \delta_\tau a^4 Z^4 N_g(b)^2 - 32A^3 s^2 D_z^2 b^5.$$

Finally define the tied target value

$$M_\tau = 16 - \delta_\tau c^2 - 16ABs^2.$$

PROVED: soundness. CONDITIONAL ON SCHINZEL H: completeness. Soundness, $F(z) \Rightarrow z \in \bigcup_{w \notin S} \mathfrak{m}_w$, is proved. The converse is open unconditionally; Sections 7–9 prove it assuming classical Schinzel H. If that lemma holds, the union outside S is $\exists_6$; adjoining the finitely many omitted maximal ideals preserves that count, so $\mathbb{Q} \setminus \mathbb{Z}$ is $\exists_6$, $\mathbb{Z}$ is $\forall_6$, and $\text{efd}_{\mathbb{Q}}(\mathbb{Q} \setminus \mathbb{Z}) \leq 5$.

2.3. Local structure and the exact global obstruction. PROVED (W0).

The first criterion, W0, is uniform even in the residue field $\mathbb{F}_3$. For every finite field k of odd order,

$$\sum_{a \in k} \chi(1 + 4a^2) = -1.$$

Hence some a makes A a nonsquare, and the canonical selector is

$$\tau_*(a) = \begin{cases} 0, & \chi(-1) = -1, \\ 2a/A, & \chi(-1) = 1. \end{cases}$$

It satisfies $\delta_{\tau_*} \neq 0$ and $\chi(-\delta_{\tau_*} A) = 1$: the two values of $-\delta_{\tau_*} A$ are $-A$ and -1 . Since $s \mapsto 1 + 2s$ is a bijection, tying loses no residue class and removes L5's $\mathbb{F}_3$ vacancy.

PROVED (W1).

For W1, let w be odd with $v_w(A) = v_w(\delta_\tau) = 0$, $v_w(b) = 1$, $v_w(z) \geq 1$, and $v_w(s) \geq 0$. The cube pullback gives either $c = 0$ or

$$v_w(c) = 2v_w(a) + 6v_w(z) - 2 \geq 6v_w(z) - 2 \geq 4.$$

Then the tied conic

$$-\delta_\tau A y^2 - 16B r^2 = 16 - \delta_\tau c^2 - 16ABs^2$$

is soluble over $\mathbb{Q}_w$ if and only if $\chi_w(-\bar{\delta}_\tau \bar{A}) = 1$. The right side is a unit congruent to 16; its two subtracted terms have valuations at least 8 and 1. Thus the target-place test is independent of the tied value of s .

PROVED (W2).

W2 identifies the remaining obstruction. Put $E_\tau = \mathbb{Q}(\sqrt{-\delta_\tau AB})$. The conic is equivalent to

$$N_{E_\tau/\mathbb{Q}}(\delta_\tau A y + 4r\sqrt{-\delta_\tau AB}) = -\delta_\tau A M_\tau, \quad M_\tau = 16 - \delta_\tau c^2 - 16ABs^2.$$

Its global solubility is exactly

$$M_\tau = 0 \quad \text{or} \quad -\delta_\tau A M_\tau \in N_{E_\tau/\mathbb{Q}}(E_\tau^\times).$$

The zero alternative is genuine. In the two branches the data are

τ	E_τ	M_τ
0	$\mathbb{Q}(\sqrt{-AB})$	$16 - c^2 - 16ABs^2$
$2a/A$	$\mathbb{Q}(\sqrt{-B})$	$16 - c^2/A - 16ABs^2$.

2.4. Structural corollaries and the remaining lemma. PROVED (L7a). At an odd place with $v(\delta_\tau A) = v(B) = 0$ and $M_\tau \neq 0$, the tied conic is soluble when $v(M_\tau)$ is even; when that valuation is odd, it is soluble exactly when $\chi_v(-\delta_\tau AB) = 1$. If also $v(A) = 0$, the corresponding untied ternary form is universal over $\mathbb{Q}_v$. Accordingly the full candidate obstruction set is

$$\{2, \infty\} \cup \{v : v(M_\tau) \text{ odd}\} \cup \{v : v(A) \neq 0 \text{ or } v(B) \neq 0\}.$$

Coefficient-bad places are genuine and cannot be replaced merely by the support of AB , because valuations can cancel in that product.

CONDITIONAL (L7b, conditional on L6 soundness). No fixed admissible (s, b) and branch admits rational functions $Y(z), R(z) \in \mathbb{Q}(z)$ satisfying

$$-\delta_\tau A Y(z)^2 - 16B R(z)^2 = M_\tau(z)$$

identically. This rules out a uniform rational section, not pointwise coverage of all relevant rational fibres by some fixed pair.

PROVED (L7c). For $M_\tau \neq 0$, local solubility is equivalent to

$$t_v := (-16\delta_\tau AB, -\delta_\tau AM_\tau)_v = 1, \quad \prod_v t_v = 1.$$

Thus a failing tied conic is obstructed at an even number of places, at least two, and repairs must change places in pairs.

CONDITIONAL ON SCHINZEL H / OPEN UNCONDITIONALLY (assembly).

For every odd target w and every z with $v_w(z) \geq 1$, one must choose (s, b) with $(1 + 2s, b) \in \Phi_1^{\{2\}}$, $s \equiv (\bar{a}_w - 1)/2 \pmod{\mathfrak{m}_w}$, $v_w(b) = 1$, and a retained branch such that $M_\tau = 0$ or $-\delta_\tau AM_\tau \in N_{E_\tau/\mathbb{Q}}(E_\tau^\times)$. In this local formulation the free b -direction must steer $E_\tau = \mathbb{Q}(\sqrt{-\delta_\tau AB})$ against the moving odd-valuation and coefficient-bad places. Later sections implement the required choice on the fixed canonical branch: L22 supplies irreducibility, L20 supplies the admissible class, and classical Schinzel H supplies the member. Thus the weak/norm-approximation problem remains open only unconditionally, not as an extra premise of the conditional theorem.

EVIDENCE. Bounded witness-switching searches support this steering mechanism, but are not a proof of assembly and are not used to upgrade the conditional six-quantifier conclusion.

3. OBSTRUCTION WALLS AND RECIPROCITY STEERING

3.1. The alignment wall. Put

$$a = 1 + 2s, \quad A = 1 + 4a^2, \quad B = 2b, \quad P = 1 - ABs^2.$$

For the two original tied branches, the source square classes are $D_0 = P$ and $D_1 = AP$, whereas the target norm fields are $E_0 = \mathbb{Q}(\sqrt{-2Ab})$ and $E_1 = \mathbb{Q}(\sqrt{-2b})$. The exact value identities show that the branch value u_τ is a norm from $\mathbb{Q}(\sqrt{D_\tau})$, while solubility asks that Au_τ be a norm from E_τ . In both branches the mismatch is measured by

$$D_\tau \text{ disc}(E_\tau) \equiv -2AbP =: \kappa \pmod{\mathbb{Q}^{\times 2}}.$$

Proposition 3.1 (L8: absorption and the parity wall). **PROVED.** *At an odd place v , assume the branch value M_τ is nonzero and each of $v(\delta_\tau A)$, $v(B)$, and $v(M_\tau)$ is even. Then the tied conic has a $\mathbb{Q}_v$ -point. Consequently an odd obstruction can occur only where A , B , or M_τ has odd valuation. On the other hand, for every admissible pair, $v_2(P) = 0$ and $v_2(\kappa) = 1$.*

Proof. After scaling the two conic variables and dividing by the even power of v contributed by M_τ , one obtains a nondegenerate binary quadratic form with unit coefficients representing a unit. Over the residue field it represents every nonzero class: in the anisotropic case it is the norm form of the quadratic extension, whose norm is surjective. Hensel lifting gives the local point. This is the absorption mechanism.

Admissibility gives $v_2(ABs^2) \geq 1$, hence P is a 2-adic unit. Thus $-2AbP$ has odd 2-adic valuation. The target discriminants $-2b$ and $-2Ab$ likewise have valuation one, so the target is always a field and can never equal the source quadratic field. $\square$

The two exact-matching devices are therefore empty on the admissible locus. Indeed, matching source and target would require $\text{disc}(E_\tau) \equiv D_\tau$, equivalently $\kappa \equiv 1$ modulo squares; splitting the target would require $\text{disc}(E_\tau)$ itself to be a square. The first condition contradicts $v_2(\kappa) = 1$, and the second contradicts the odd valuation of the target discriminant. This rules out precisely these two identities, not every possible fixed square-class relation: for a general class j , one would still have to prove that an odd-multiplicity value prime inert in $\mathbb{Q}(\sqrt{j})$ must occur, and no such theorem is asserted here.

EVIDENCE. The frozen rescues do not exhibit a further support-controlled identity: all ten have odd-multiplicity wild primes, and all measured symbols at those primes are $+1$. In the recorded session sweep, thirteen of 161 admissible pairs succeeded, while every observed obstruction set had even cardinality. These observations are alignment statistics only; they do not prove a uniform mechanism or an assembly theorem.

3.2. Reciprocity steering.

Lemma 3.2 (L9 steering lemma). **PROVED.** For $x, d \in \mathbb{Q}^\times$, let

$$T = \{2, \infty\} \cup \{q \text{ odd} : v_q(x) \neq 0 \text{ or } v_q(d) \neq 0\}.$$

Then $\prod_{v \in T} (x, d)_v = 1$. Hence, if all symbols in T except possibly one are $+1$, the remaining symbol is also $+1$.

Proof. Hilbert reciprocity gives the product over all places. Outside T , both entries are odd-adic units and their Hilbert symbol is $+1$, so only the displayed finite set remains. $\square$

For the tied conic take $x = \alpha M_\tau$, $d = \alpha B$, and $\alpha = -\delta_\tau A$. Once every controlled symbol is $+1$, a single wild odd-multiplicity prime q_0 with $v_{q_0}(d) = 0$ is not an obstruction: its symbol is forced by the lemma. Thus the useful arithmetic target is not complete smoothness, but a controlled part together with one proven prime-power cofactor.

Proposition 3.3 (L9a: the prime- b symbol). **PROVED.** Fix an admissible a , a cell (w, z) , and a branch τ . Let $b = \varepsilon q_1$, where q_1 is odd prime, and impose separately

$$v_{q_1}(\delta_\tau) = v_{q_1}(A) = v_{q_1}(a) = v_{q_1}(z) = v_{q_1}(Dz) = 0.$$

Then

$$(x, d)_{q_1} = \left(\frac{A}{q_1} \right).$$

In particular, the moving place is aligned by choosing a residue class with $\left(\frac{A}{q_1} \right) = +1$.

Proof. Clearing denominators and reducing at $b \equiv 0 \pmod{q_1}$ gives $N_g(0) = -A$. The hypotheses make this a unit, the denominator contributes valuation four, and $v_{q_1}(x) = -4$. Its unit part is A modulo squares, while $v_{q_1}(d) = 1$; the Hilbert-symbol parity formula gives the stated Legendre symbol. Quadratic reciprocity makes it constant on suitable congruence classes. $\square$

The corrected progression freezes only genuine controlled places. With $S_0 = \{2\} \cup \text{supp}(\alpha)$, the modulus contains the required 2-adic and A -character moduli and the Taylor exponents at places of S_0 ; it does not freeze the moving prime. For a coprime prime q_1 in the selected class,

$$\text{supp}(d) = S_0 \cup \{q_1\}, \quad \left(\frac{A}{q_1}\right) = +1.$$

Thus every place in the support of d is controlled, and only the odd-multiplicity primes of the remaining cofactor of x are wild.

The need for the corrected modulus is witnessed by the recorded cell $(w, u) = (73, 5)$. With $s = 0$, $b_0 = -29$, $\tau = 2/5$, the superseded recipe gives $N = 689120$. The prime $q = 244637629 = 29 + 355N$ has the same prescribed class, sign, and value of $(A \mid q)$, but at 59 the symbol changes: $v_{59}(x(b_0)) = 0$ with symbol $+1$, whereas $v_{59}(x(-q)) = 1$ with symbol -1 .

CONDITIONAL. Turning such a progression into an infinite assembly family still requires the named Schinzel–H/Bunyakovsky input that simultaneously makes q_1 prime and leaves a single prime cofactor of x . Reciprocity removes the last symbol once that input is supplied; it does not prove the prime-value assertion.

3.3. The generalized coprimality wall.

Theorem 3.4 (L12a). ***PROVED.** Let a be admissible and suppose $A = 1 + 4a^2$ is prime. Let τ be any branch, assume $v_A(z) \leq 0$, and suppose every odd-valuation place of an admissible rational b is coprime to $\{2, z, D_z, a, \delta_\tau, A\}$. If b_{sf} denotes the squarefree part of its odd part, then*

$$\prod_{p \in \{A\} \cup \text{oddsupp}(b)} (x, d)_p = -1.$$

Thus at least one of these places obstructs every branch.

Proof sketch. At each odd place p of b , whether p occurs in the numerator or denominator, the dominant-term calculation makes $v_p(x)$ even and its unit part

$$u_x \equiv A \pmod{\mathbb{Q}_p^{\times 2}}.$$

Therefore $(x, d)_p = (A \mid p) = (p \mid A)$, and multiplication over these places gives $(b_{\text{sf}} \mid A)$. At A , the branch-uniform parity calculation gives $(x, d)_A = -(b_{\text{sf}} \mid A)$: the minus sign is $(2 \mid A) = -1$, since $A \equiv 5 \pmod{8}$, while the branch-dependent unit of α is a residue. The two Jacobi factors cancel and leave -1 . $\square$

The only door in this argument is non-coprimality. If b shares a prime f with the cell data, the reduction to the constant term at f and hence $u_x \equiv A$ can fail; the telescoping proof then no longer applies. This is a door, not by itself a solubility claim.

3.4. Composite squarefree wall.

Theorem 3.5 (L13e). ***PROVED.** Let a be admissible, put $A = 1 + 4a^2$, and let K be the squarefree kernel of its rational square class; this includes composite, non-squarefree, and rational A . Assume $v_p(z) \leq 0$ for every odd $p \mid K$ and that every odd-valuation place of an admissible rational b is coprime to $\{2, z, D_z, a, \delta_\tau, A\}$. Then, for every odd $p \mid K$ and every odd ℓ occurring to odd order in b ,*

$$(x, d)_p = \left(\frac{2b}{p}\right), \quad (x, d)_\ell = \left(\frac{A}{\ell}\right),$$

and

$$\prod_{p \mid K} (x, d)_p \prod_{\ell \in \text{oddsupp}(b)} (x, d)_\ell = \left(\frac{2}{K}\right)^{1+v_2(b)}.$$

Proof sketch. At an odd $p \mid K$, the middle term $-\delta_\tau a^4 Z^4 N_g^2$ strictly dominates the other terms of the value polynomial, and $v_p(N_g) = 0$. Hence $v_p(x)$ is odd and its unit class contains the complementary factor K/p . Every prime divisor of K is $1 \pmod{4}$, because it divides a value $n^2 + 4m^2$ with the data coprime. In the Hilbert-symbol formula, the complementary factors from the two entries cancel. This yields $(2b \mid p)$ independently of the branch.

Multiplying over $p \mid K$ and over the moving primes of b produces the cross terms $(M \mid K)(K \mid M)$. Quadratic reciprocity cancels them factor by factor because every divisor of K is $1 \pmod{4}$. Moreover $K \equiv A \equiv 5 \pmod{8}$, so $(2 \mid K) = -1$. Admissibility is exactly $v_2(b) = 0$, for which the product is -1 ; at the first non-admissible boundary $v_2(b) = 1$, the product is $+1$ and this wall dissolves. Even-exponent primes of A do not enter K . $\square$

It follows that the necessity direction of L11h is a theorem for every admissible a : escaping the coprime wall requires a prime $p \mid K$ with $v_p(z) > 0$, and every such p is congruent to 1 modulo 4 and divides the numerator of z . Hence reachability implies that the numerator of z has a prime divisor congruent to 1 modulo 4. This conclusion does not supply the separate prime-value input needed for assembly.

4. CLASS DICHOTOMY, CHARACTERIZATION, AND ESCAPES

Fix an admissible parameter a , put

$$A = 1 + 4a^2, \quad \delta_\tau = 1 - A\tau^2, \quad \alpha = -\delta_\tau A,$$

and first consider the prime-member family $b = \varepsilon q$, with $\varepsilon \in \{\pm 1\}$ and q prime. An *aligned class* means a nonempty prime progression in which the Hilbert symbols at every controlled finite place, at the moving place q , and at the real place are all $+1$. This is deliberately a frozen-side notion: odd-valuation primes emerging from the value of $P(b)$ are a separate obligation.

4.1. The L10 dichotomy.

Proposition 4.1 (Canonical alignment and the opposite wall; PROVED in the stated scope). *On the canonical branch $\tau = 2a/A$, the prime- b family is aligned with the canonical square class -1 in the exact sense that*

$$\delta_\tau = \frac{1}{A}, \quad \alpha = -1.$$

On the branch $\tau = 0$, if a is odd, A is prime, $A \nmid 2zD_z$, and the prime $q \neq A$ satisfies the prescribed unit conditions, then no aligned class exists.

The first assertion is an identity, not permission to discard the places above A : although α is an A -unit, $\delta_\tau = 1/A$ forces those places back into the controlled set. This distinction is essential; the obsolete grid count obtained by omitting them is not used here. For the second assertion, $\tau = 0$ gives $\delta_\tau = 1$ and $\alpha = -A$. The local calculation at A yields

$$(x, d)_A \left(\frac{A}{q} \right) = \left(\frac{-2\varepsilon}{A} \right) = -1.$$

Indeed admissibility gives $A \equiv 5 \pmod{8}$, so $\left(\frac{-1}{A} \right) = +1$ and $\left(\frac{2}{A} \right) = -1$, independently of the sign ε . Two required symbols are therefore permanently anti-correlated. This is a theorem for prime integral A under the displayed hypotheses; it is not a claim extracted from a finite failure search.

4.2. Free branches, an immovable wall. Branch completion itself costs nothing. If $\tau = p/q$ is rational in the existing parameters and its denominator is proved nonzero on the base locus, adjoining that branch introduces no inverse witness. Admissibility makes A a nonsquare in $\mathbb{Q}_2$, hence $\delta_\tau \neq 0$; the norm identity in the unrefereed argument of Sun [16] is uniform in τ ; and a finite

disjunction of branch equations is represented by their product in the same witnesses. Explicit choices meet all four square classes of $\langle -1, A \rangle$, including the square branch

$$\tau^\dagger = \frac{1 + 2a^2}{1 + 4a^2}, \quad \alpha = (2a^2)^2.$$

The corrected controlled set for a prime member is

$$S = \{2, 3, 5, 7\} \cup \text{supp}(\alpha) \cup \text{supp}(\delta_\tau).$$

Outside S , the fixed content of P has even valuation; after that content is removed, the reduction is a nonzero polynomial whose degree prevents a permanent odd valuation at the remaining primes. Thus no fixed bad place is silently omitted.

More importantly, changing branches does not change the obstruction. For prime A , admissible a , and $v_A(z) \leq 0$, every branch in the relevant square-class group satisfies

$$(x, d)_A(x, d)_q = -1.$$

Hence at least one controlled symbol is negative. This branch-independent wall was checked exactly on 2,136 extended instances across all four explicit branches; the computation confirms the proved identity rather than replacing it.

Theorem 4.2 (The L11h characterization; PROVED). *For the prime- b family, an aligned class exists at the cell (w, u) , with $z = wu$, if and only if the numerator of z has a prime divisor $p \equiv 1 \pmod{4}$. On the finite grid this selects 190 of 353 cells; the other 163 cells are unreachable by this family on every branch.*

Proof sketch. For necessity, the generalized wall says that alignment is impossible unless some odd prime $p \mid A$ also has $v_p(z) > 0$. Write $a = m/n$ in lowest terms. Such a prime divides $n^2 + 4m^2$, and therefore $(2mn^{-1})^2 \equiv -1 \pmod{p}$; consequently $p \equiv 1 \pmod{4}$. The factor-by-factor telescoping argument applies also when A is composite or rational, so this necessity is PROVED for every admissible a , not merely supported by a search.

Conversely, let $p \equiv 1 \pmod{4}$ divide the numerator of z . Choose odd representatives m, n satisfying $n^2 \equiv -4m^2 \pmod{p}$ and set $a = m/n$. Then a is admissible and $v_p(A) > 0$, so the factor shared by A and z breaks the wall. Freezing the complete controlled set gives an explicit aligned progression. The construction, including its escape prime, chosen a , branch, and modulus, is replayed on all 190 predicted rows. This constructive sufficiency and the preceding necessity make the equivalence the structural centre of the class analysis. $\square$

4.3. Non-coprime escapes and complete grid coverage. The characterization concerns $b = \varepsilon q$. The only effective door through the generalized coprime wall is to share a cell prime deliberately:

$$b = \varepsilon f q, \quad f \mid z.$$

This produces aligned escape classes at 103 of the 163 prime-family-walled cells. Their frozen-side alignment is PROVED row by row; the remaining 60 walled cells with no such escape are closed instead by direct, fully soluble witnesses.

The Schinzel audit also keeps six legacy residual rational/composite- b witnesses separate as pointwise certificates. They are unconditional local witnesses, but they are not promoted into prime progressions and give no prime-value theorem. The eventual last cell, $(89, -2)$, was closed by the replayed square-branch witness $(a, b, \tau) = (3, 89/367, 19/37)$, completing cell coverage.

verified object	coverage	status
L11 prime- b aligned classes	190/190	PROVED PER ROW
non-coprime escape classes	103/103	PROVED PER ROW
all aligned classes, with a zero-bad member	293/293	PROVED PER ROW
all grid cells, with a soluble witness	353/353	PROVED PER ROW

The class total is exactly the disjoint sum of the L11 and escape families; the cell total additionally uses direct pointwise witnesses. Neither total asserts a uniform theorem beyond the verified grid.

4.4. Why no square-class shortcut remains.

Proposition 4.3 (L13c; PROVED per row). *For every one of the 706 cell-branch rows, P is irreducible of degree 8 over $\mathbb{Q}$. Consequently no constant-square or weak square-factor form collapses the emergent-place obligation on the grid.*

The boundary between theorem and experiment is sharp. The irreducibility certificates are PROVED. By contrast, the 137.5 million exact square-class tests with no admissible hit are EVIDENCE against a hidden special-value shortcut, not a universal nonexistence theorem. Likewise the observed Galois group $D_8 \wr C_2$ is EVIDENCE: its cycle types matched all 28,240 certified unramified Frobenius samples, but no group-identification proof is claimed.

Finally, the earlier zero-bad search already supplied 24 certified members across 15 distinct cells; those are PROVED INSTANCES, while the rough quarter-density inferred from the sample is only EVIDENCE. The later per-class replay strengthens the finite conclusion to the 293/293 line in the table. The uniform member property is now conditional solely on classical Schinzel H, by the L22 closure and the fixed-pair implication.

5. UNIFORM EXISTENCE OF AN ALIGNED CLASS

The prime- b characterization of Section 4 describes one family, not the availability of classes altogether. On the square branch, controlled non-coprimality gives a class for every cell. This removes the class-existence clause from the global hypothesis; it does not produce an emergent-free member.

Theorem 5.1 (Uniform class existence; PROVED). *Let w be an odd prime and let $z \in \mathbb{Q}^\times$ satisfy $v_w(z) \geq 1$. There are an odd integer a , a fixed factor $f = w$, and infinitely many positive primes q_1 for which the square branch*

$$A = 1 + 4a^2, \quad \tau^\dagger = \frac{1 + 2a^2}{A}, \quad \delta = -\frac{4a^4}{A}, \quad \alpha = (2a^2)^2,$$

with $\varepsilon = +1$ and $b_0 = fq_1$, is an aligned base class for the cell (w, z) . If

$$S = \{2, 3, 5, 7\} \cup \text{supp}(\alpha) \cup \text{supp}(\delta) \cup \{f\},$$

then the class freezes every symbol at S , the moving-prime symbol, and the real symbol at $+1$.

Proof. We give the construction, including the residue count. First choose a . For the quadratic character χ_f the standard character sum is

$$\sum_{r \bmod f} \chi_f(1 + 4r^2) = -1.$$

There are $1 + \left(\frac{-1}{f}\right)$ zero terms. Hence the number of residues with $\chi_f(1 + 4r^2) = -1$ is

$$\frac{f - \left(\frac{-1}{f}\right)}{2} = \begin{cases} (f-1)/2, & f \equiv 1 \pmod{4}, \\ (f+1)/2, & f \equiv 3 \pmod{4}, \end{cases}$$

and is always positive. Choose such a residue and lift it to an odd integer a ; adding the odd modulus f changes the parity of a lift. Thus

$$(E1) \quad \left(\frac{A}{f}\right) = -1.$$

Taking $f = w$ is always admissible. Indeed, (E1) implies $f \nmid aA$, so $v_f(\delta) = 0$. With $Z = z^3$ and $D = 1 - Z - a^2Z^2$, the cell condition gives $v_f(Z) \geq 3$ and $D \equiv 1 \pmod{f}$. Thus f is an odd numerator prime, is coprime to $aA\delta D$, and meets every fixed-factor guard. This includes $f = 5$; there is no exceptional denominator or gcd case.

Put

$$M = 4A \prod_{p \in S} p$$

and require the following reduced residue system for q_1 :

$$(E2) \quad q_1 \in (\mathbb{Z}/M\mathbb{Z})^\times, \quad \left(\frac{2fq_1}{p} \right) = +1 \quad (p \in S \text{ odd}, p \neq f).$$

At each distinct odd prime in (E2), precisely half of the unit classes satisfy the displayed character. The Chinese remainder theorem therefore gives exactly

$$(E3) \quad \frac{\varphi(M)}{2^{\#\{p \in S: p \text{ odd}, p \neq f\}}}$$

reduced classes modulo M . In particular (E2) is nonempty. Dirichlet's theorem gives infinitely many positive primes q_1 in any selected class. Removing the finitely many primes dividing the fixed cell data or producing a root or degeneracy leaves infinitely many choices.

It remains to verify the local symbols. Write $s = (a - 1)/2$ and

$$N_g(b) = 16a^4b^2 - A(b - 1)^4,$$

so the kernel polynomial is

$$(E4) \quad P(b) = 16D^2A^2b^4 - \delta a^4Z^4N_g(b)^2 - 32A^3s^2D^2b^5.$$

Since α, D^2, A^2 , and b^4 are squares, the square class of the local conic input x is that of $P(b)$.

At f , one has $v_f(b_0) = 1$, $N_g(b_0) \equiv -A \pmod{f}$, and the three terms of (E4) have valuations 4, at least 12, and at least 5. The uniquely minimal term is a square, so x is a square in $\mathbb{Q}_f$ and the f -symbol is +1.

At 2 the exact computation is needed because the least unit need not be a square. The kernel gives

$$(E5) \quad v_2(x) = 6, \quad v_2(d) = 3, \quad u_d \equiv b_0 \pmod{8}, \quad u_x \equiv 1 - 2As^2b_0 \pmod{8}.$$

If s is even then $u_x \equiv 1$. If s is odd, then $A \equiv 5 \pmod{8}$ and $u_x \equiv 1 - 2b_0 \pmod{8}$. In the standard 2-adic Hilbert formula the exponent is

$$\epsilon(u_x)\epsilon(u_d) + 6\omega(u_d) + 3\omega(u_x) \pmod{2}, \quad \epsilon(u) = \frac{u-1}{2}, \quad \omega(u) = \frac{u^2-1}{8}.$$

For $b_0 \equiv 1, 3, 5, 7 \pmod{8}$, the corresponding u_x are 7, 3, 7, 3, and the exponent is respectively 0, 0, 0, 0. Thus the 2-adic symbol is always +1.

For every other odd $p \in S$, (E2) makes $2fq_1$ a square unit. Because α is a rational square, $d = 2\alpha f q_1$ is a square in $\mathbb{Q}_p$ even when $p \mid a$; hence $(x, d)_p = +1$. At infinity $d > 0$, so the real symbol is +1.

At the moving prime the kernel identity is

$$(x, d)_{q_1} = \left(\frac{A}{q_1} \right).$$

Every prime divisor of A is $1 \pmod{4}$, because $(2a)^2 \equiv -1$ modulo that prime. Reciprocity and (E2) therefore give

$$(E6) \quad \left(\frac{A}{q_1} \right) = \prod_{p^e \parallel A} \left(\frac{q_1}{p} \right)^e = \left(\frac{2f}{A} \right) = \left(\frac{2}{A} \right) \left(\frac{f}{A} \right) = (-1)(-1) = +1.$$

Here $A \equiv 5 \pmod{8}$ and $\left(\frac{f}{A}\right) = \left(\frac{A}{f}\right) = -1$. This is the compatibility identity that prevents a hidden collision between the frozen and moving places.

Finally define

$$k_p = \texttt{L10_exponent}(P, b_0, p), \quad N = \text{lcm}(8, \{p^{k_p} : p \in S\}, 4A).$$

The Taylor-exponent lemma freezes the S -symbols throughout $q_1 + N\mathbb{Z}$, while $4A \mid N$ freezes (E6). Since (E2) makes q_1 a unit at every prime dividing N , $\gcd(q_1, N) = 1$. This completes the class construction. Emergent places outside S have deliberately not been controlled; finding a member without a negative emergent symbol is the separate member-existence problem. $\square$

Remark 5.2 (The canonical parameter reproduces the 5-wall). For $a = 1$ one has $A = 5$ and $\tau^\dagger = 3/5$. The independently computed frozen and moving symbols reduce to

$$\text{Hilb}_5 = -\left(\frac{f}{5}\right)\left(\frac{q_1}{5}\right), \quad \text{Hilb}_{q_1} = \left(\frac{5}{q_1}\right) = \left(\frac{q_1}{5}\right).$$

They can both be $+1$ exactly when $\left(\frac{f}{5}\right) = -1$. For $f = w$ and the prime- b route unavailable, this fails exactly at $w \equiv 11, 19 \pmod{20}$, the independently derived canonical 5-wall of Theorem 10.12. The separate replay recovered all 60 walled grid cells and repaired all 60 after allowing a to vary. Thus the collision at $a = 1$ is exact, while class nonexistence is not.

6. THE VERIFIED QUANTITATIVE LAYER

The class construction reduces the residual local problem to a moving arithmetic progression. For an aligned class write

$$Q(k) = q_1 + kN, \quad b(k) = \varepsilon f Q(k),$$

with the controlled Hilbert symbols already fixed to be positive. A member is called *emergent-free* when the remaining odd-valuation places contribute no negative symbol. The purpose of the quantitative layer is twofold: to certify this condition on the finite grid and to separate the exact local mechanism from the statistical model suggested by the resulting population.

6.1. L14: instance verification.

Proposition 6.1 (Finite instances of the member property $\mathcal{M}$). ***PROVED (per row).** All 293/293 aligned classes in the finite grid have a certified emergent-free member; these comprise 190 L11 classes and 103 escape classes. Every stored row was replayed through the exact cofactor ladder and returned verdict zero.*

The proposition is deliberately finite. Its acceptance condition is an aligned class together with an exact zero verdict from the member ladder. A tied-conic or ramification computation may also be attempted as an auxiliary cross-check, but a budget refusal there is accepted and logged: it neither invalidates a ladder certificate nor supplies positive evidence. In particular, no refusal is promoted to a primality or solubility assertion. This distinction is essential because the general all-cell member property is conditional on classical Schinzel H, even though the displayed grid instance is complete.

6.2. L15: the exact remainder and counting laws. For a certified member, let R be the remainder of $P(b)$ after the frozen support, the fixed shell, and the smooth-emergent primes have been stripped. An independent replay gives the following exhaustive shape table.

Thus R is squarefree on every audited row and $R = 1$ never occurs. The prime rung is not a probable-prime shortcut: its singleton cofactor is accepted only after a proof of primality. On a factorint rung the integer e counts distinct emergent factors, and the exact factorization allows every corresponding Hilbert symbol to be checked. The two shapes therefore reflect two proof routes through the same local criterion, not two definitions of closure.

Ladder shape	Rows	Exact form of R	Size data
proved-prime	197	one prime, never pC^2	median 48 digits; maximum 119
factorint	96	squarefree product, $e \in \{2, 3, 4\}$	multiplicities 76/15/5

TABLE 1. **PROVED (finite audit)**. Shapes of the stripped remainder on all 293 closure rows.

There is also an exact finite counting statement. The first certified member occurs at $k = 0$ in 207/293 classes, or 70.65%. This is **PROVED** as a statement about deterministic playback of the stored progressions. By contrast, interpreting the searches as samples from a stable member process is **EVIDENCE**: the measured per-prime-member emergent-free rate is approximately 0.040, and the observed waiting time is approximately 15.7 times the pure Bateman–Horn [1] first-prime prediction for the same progressions. The exact histogram may therefore be used without qualification inside the finite audit, while the rate and the comparison are empirical laws, not uniform density theorems.

6.3. L16: why the small layer has sieve shape. Fix an emergent odd prime p . On an aligned class, $b(k)$ is affine in k , and the sign at p is

$$(x_0, d_0)_p = \left(\frac{2\alpha b(k)}{p} \right).$$

On every checked recurring-prime row, this sign is determined by $k \bmod p$. The obstruction at p is consequently not the whole root set of $P(b(k))$ modulo p : it is the union of those root classes having the bad signature, further restricted to odd valuation,

$$\mathcal{B}_p = \{k : v_p(P(b(k))) \text{ is odd}\} \cap \{k \bmod p : \left(\frac{2\alpha b(k)}{p} \right) = -1\}.$$

For a simple bad root, Hensel lifting gives an exact Haar mass

$$\sum_{j \geq 0} \frac{p-1}{p^{2j+2}} = \frac{1}{p+1}.$$

This is the valuation-parity correction to a naive exclusion sieve: one must retain the odd levels of the Hensel tower rather than merely delete a residue class modulo p . A singular root has no such one-line factor. Its descendants are lifted recursively until the odd and even masses are certified with a residual bound; if that cannot be completed, the corresponding local product is explicitly labeled as an exclusion or partial result. Accordingly, the residue and valuation checks are **PROVED per row**; the claim that this small layer supplies the useful global layering model is **EVIDENCE**. No independence theorem is hidden in the product notation.

6.4. L17: matched validation. The full-grid residue test strengthens the preliminary structural probe. Across all 293 classes and the window $k \in [0, 119]$, it found zero residue-determination counterexamples on the identical 8,760-member sample used for evaluation. This is **PROVED** as a finite statement. The predictive use of the local masses is **EVIDENCE**; it was assessed only after the predictions and observations had been put on exactly the same member support.

The marginal statistic counts odd bad-root hits, so a member can contribute more than once; the union statistic asks whether a member is hit at least once. Their simultaneous agreement, together with the marginal-preserving permutation null, shows no detectable overlap or dependence discrepancy on this window. It does not prove independence outside the evaluated support.

A superficially different class-weighted comparison has estimand about +0.0195, at 3.63 standard deviations. That quantity compares Haar-uniform local masses in $\mathbb{Z}_p$ with the finite $k \in [0, 119]$ window, rather than comparing predictions and observations member by member on matched

Matched statistic	Model	Observed	Diagnostic
marginal odd hits	4481.3	4478	ratio 0.9993
hit-conditioned union	3477.9	3475	residual -2.9
permutation null	—	3475	$z = -0.09$; tails 0.61/0.53

TABLE 2. **EVIDENCE.** Matched small-layer evaluation on the same 8,760 members used on both sides.

support. It is therefore identified as the Haar-window gap, not as a failure of the hit-conditioned model.

6.5. L17 cofactor layer: reciprocity and its limit. The large-cofactor layer has a narrower theorem than the small layer. **PROVED:** on the prime rungs, reciprocity forces the singleton cofactor sign to be $+1$ in all 197/197 cases; on the complete closure set, the cofactor-sign product is $+1$ on 293/293 rows. These are exactly the singleton and parity-product statements. The individual positive signs in factorint closure rows are selected by the closure condition, so their residue tables are descriptive data and cannot support an unconditional sign-distribution claim.

The attempted nonclosure measurement remains **INCONCLUSIVE:** only 11/400 sampled rows were resolved, and the predeclared power gate failed. The unresolved rows are refusals, not observations of either sign. Consequently the cofactor layer is proved at reciprocity level and no further; its nonclosure sign distribution remains measurement-limited. This is also why the matched small-layer success must not be silently upgraded into a complete rate theorem for emergent-free members.

7. THE CONDITIONAL RECORD AND THE SCHINZEL AUDIT

7.1. The statement, with its hypothesis exposed. The member property below records the output required by L6; it is no longer an additional conjectural premise.

Definition 7.1 (Member property $\mathcal{M}$). For every odd prime w and every rational z with $v_w(z) \geq 1$, there is a verified aligned class for (w, z) and a prime member $Q \equiv q \pmod{N}$ outside its finite excluded set whose bad emergent set is empty. Explicitly, for every odd prime $\ell \notin S \cup \{Q\}$ with $v_\ell(P(\varepsilon fQ))$ odd, the local Hilbert symbol $(x_0, d_0)_\ell$ is $+1$, where S is the frozen set of the class.

Theorem 7.2 (CONDITIONAL ON CLASSICAL SCHINZEL H: Theorem C). *Assume classical Schinzel's Hypothesis H. For every odd cell, use the fixed canonical branch*

$$\tau^\dagger = \frac{1 + 2a^2}{1 + 4a^2}, \quad \delta_{\tau^\dagger} = -\frac{4a^4}{1 + 4a^2}.$$

Then the member property $\mathcal{M}$ holds and $\mathbb{Q} \setminus \mathbb{Z}$ is Diophantine in $\mathbb{Q}$ with six unknowns, realized by the L6 formula

$$F(z) := \exists b, s \left[(1 + 2s, b) \in \Phi_1^{\{2\}} \wedge \Theta_*(1 + 2s, b, h(1 + 2s, b, z^3), s) \right],$$

where, with $A = 1 + 4a^2$, $B = 2b$, and $\tau^\dagger = (1 + 2a^2)/A$,

$$\begin{aligned} \Theta_*(a, b, c, s) := \exists y, r \quad & [c^2 - Ay^2 - 16Br^2 = 16 - 16ABs^2] \\ & \vee [c^2 - Ay^2 - 16ABr^2 = 16A - 16A^2Bs^2] \\ & \vee [a^4(c^2 - Ay^2) + 4ABr^2 = 4A^2Bs^2 - 4A]. \end{aligned}$$

The three disjuncts remain the cleared equations for $\tau = 0$, $\tau = 2a/A$, and $\tau^\dagger$, respectively, in the same witnesses (y, r) . They preserve the finite soundness menu and the six-count; completeness in this theorem uses only the third, fixed canonical disjunct. Consequently $\mathbb{Z}$ is definable in $\mathbb{Q}$ by a formula with six universal quantifiers, $\mathbb{Z} = \mathbb{Q} \setminus (F \vee \mathbf{m}_2)$, and $\text{efd}_{\mathbb{Q}}(\mathbb{Q} \setminus \mathbb{Z}) \leq 5$.

This is a conditional definition, not an unconditional solution of Hilbert's Tenth Problem over $\mathbb{Q}$. Soundness of the finite branch menu consumes no hypothesis. For completeness, fixed- Z vertical irreducibility and Hilbert selection give a good canonical-branch a in the existing admissible progression; L20 supplies the class and all Schinzel-pair conditions; Theorem 8.1 turns classical Schinzel H into an emergent-free member; and L6 gives the displayed definition. No per-cell irreducibility certificate is assumed.

The quantifier saving is attributed precisely to Nicolas Daans, Philip Dittmann, and Arno Fehm, *Existential rank and essential dimension of diophantine sets*, arXiv:2102.06941v5, Theorem 1.4 (equivalently Corollary 5.11) [6]. Their intersection theorem turns the ranks three and two on the common base into $3 + 2 - 1 = 4$ witnesses; the two base coordinates then give six unknowns. The recorded rank-essential-dimension identity gives the bound five. These counts remain **CONDITIONAL** because the set equality to $\mathbb{Q} \setminus \mathbb{Z}$ uses classical Schinzel H.

7.2. What is proved, and what Schinzel must supply. PROVED (uniform class side and linear side). Theorem 5.1 constructs an aligned class for every cell and freezes all controlled Hilbert symbols while restricting the moving prime to one reduced progression

$$q_1(t) = q_{1,0} + Nt, \quad \gcd(q_{1,0}, N) = 1.$$

The residue construction gives infinitely many proven-prime choices of the base $q_{1,0}$, and Dirichlet's theorem supplies infinitely many prime values of the progression. This is the linear, unconditional half of the member-existence problem; no probable-prime test or refusal is used as evidence.

CONDITIONAL (the sole conjectural cofactor side). Put

$$F_{\text{cell}}(t) = P(\varepsilon f q_1(t)),$$

using the primitive integer representative after removal of the fixed rational scale recorded in the audit. The L22 theorem and quantitative Hilbert irreducibility make F_{cell} irreducible for a selected admissible a . Requiring it to be prime simultaneously with $q_1(t)$ is the two-polynomial instance of classical Schinzel's Hypothesis H [14]. The polynomial has degree eight. A prime cofactor gives an emergent-free member because the class has frozen the controlled places and reciprocity forces the remaining single symbol. After using the progression prime as variable, the pair is $\{q_1 + Nt, F_{\text{cell}}(t)\}$.

7.3. PROVED finite audit of the Schinzel pair. The exact audit contains 293 canonical records, split into 190 L11 classes and 103 escape classes. The fields `F_irreducible`, `F_value_gcd`, and `pair_product_gcd` respectively certify irreducibility on 293/293 rows, value gcd one on 293/293 rows, and pair-product gcd one on 293/293 rows. Every irreducibility certificate is a degree-eight Frobenius certificate modulo a certified prime; both gcds were computed exactly on $0 \leq t \leq 2000$. The pair-product computation is already a certificate that no prime divides the product for every integer t : any such fixed divisor would divide the computed gcd. Thus all canonical pairs pass the fixed-divisor and irreducibility conditions. This is **PROVED** for the stored pairs, not a prime-value theorem.

The finite audit does not by itself extrapolate to arbitrary rational cells. The L20 audit proves normalization, coprimality, absence of a fixed divisor, frozen-unit and symbol constancy, and positive leading coefficient uniformly. Theorem 9.1 now supplies the missing uniform input directly: for every fixed nonzero rational Z , $P(a, Z)$ is irreducible in $\mathbb{Q}(a)[b]$. Corollary 9.2 then selects a good a inside the character progression. Thus the 353 recorded target certificates remain exact finite corroboration, but they are no longer a premise for a new cell.

For a prime p dividing N , the corrected residue count is

$$\nu_p = \#\{t \in \mathbb{F}_p : q_1(t) \not\equiv 0 \pmod{p}, F_{\text{cell}}(t) \not\equiv 0 \pmod{p}\}.$$

The first condition is essential. Since $p \mid N$ and $\gcd(q_{1,0}, N) = 1$, the linear polynomial is the nonzero constant $q_{1,0}$ modulo p ; it must not be replaced by the parameter t . Consequently the

count is p , rather than $p - 1$, whenever the cofactor has no root modulo p . The audit enumerated every $p \leq 1000$ dividing the row's modulus and obtained exactly this full count in every occurrence.

primes p	records containing p N	range of ν_p	mean density ν_p/p
2, 3, 5, 7	293 each	p	1
11, 37, 89, 97	13 each	p	1
13, 53	26 each	p	1
17	39	p	1
23	38	p	1
29, 41, 61, 73	28 each	p	1
43, 47, 67, 83	10 each	p	1

TABLE 3. Corrected admissible-residue counts for primes dividing the progression modulus.

These local counts neither assume independence nor estimate a density of simultaneous prime values. Their role is narrower and exact: together with the pair-product gcd, they rule out the elementary local obstruction for each stored Schinzel pair.

7.4. Residual certificates and the literature boundary. The six residual rows are deliberately segregated. They are rational or composite- b witnesses verified by exact branch reconstruction and tied status; they are not coerced into the prime family and are not evidence for Schinzel's hypothesis.

cell (w, z)	(a, b)	τ	status
(67, 2)	(1, 4757)	3/5	PROVED pointwise
(41, 1/7)	(25, 4985)	1251/2501	PROVED pointwise
(61, -2/7)	(25, 55571)	1251/2501	PROVED pointwise
(61, 2)	(25, -71431)	1251/2501	PROVED pointwise
(53, 1/3)	(-15, 9)	451/901	PROVED pointwise
(29, -2)	(-15, -29/5)	4055/2703	PROVED pointwise

TABLE 4. The six residual witnesses: unconditional certificates for their individual cells only.

OPEN (the unconditional published boundary). The exact square-class reference is David Krumm, *Squarefree parts of polynomial values*, Journal de Théorie des Nombres de Bordeaux **28** (2016), no. 3, 699–724, arXiv:1407.4890 [11]. Its Theorem 1.3 and Propositions 3.4–3.5 give the unconditional degree-one and degree-two cases; Proposition 3.8 makes the degree-three case conditional on the Parity Conjecture for elliptic curves over $\mathbb{Q}$.

At prime arguments, the nearest unconditional results reach the cubic boundary without prescribing a square class: Helfgott proves square-free values of $f(p)$ for separable cubic f [8], while Reuss proves $(d - 1)$ -free values rather than prime or prescribed-square-class values [12]. Neither yields the required degree-eight conclusion.

The conic-bundle literature does not remove the gap. The unconditional Harpaz–Skorobogatov–Wittenberg theorem requires abelian constant fields in the bad fibres [7, Theorem 3.1 and Corollary 3.4]. The general Schinzel reference in *J. reine angew. Math.* **495** (1998), 1–28 is by Colliot-Thélène, Skorobogatov and Swinnerton-Dyer [3], not by Colliot-Thélène and Sansuc; the latter pair's Schinzel paper is [2]. No cited unconditional theorem supplies simultaneous prime values for the present linear–octic pair. Classical Schinzel H is therefore the sole conjectural input to the six-quantifier theorem, which remains **CONDITIONAL**; no unconditional record or solution of $H10(\mathbb{Q})$ is claimed.

8. SCHINZEL'S HYPOTHESIS H: THE FIXED-CLASS IMPLICATION

Theorem 5.1 supplies the canonical aligned-class construction for every cell. Section 9 proves, for every fixed nonzero rational Z , that its octic is irreducible in $\mathbb{Q}(a)[b]$, and quantitative Hilbert irreducibility selects an irreducible specialization inside the existing character-admissible a progression. The L20 admissibility package is therefore unconditional. For any such fixed class, classical Schinzel's Hypothesis H [14] is the sole remaining input and supplies the member.

8.1. The operational contract. A place is recorded as an obstruction only when its valuation is odd *and* its Hilbert symbol is -1 : the kernel appends a place under exactly that conjunction, and the ladder returns *zero* when every remaining place is proved with symbol $+1$. Emergent-freeness therefore means *no place of symbol -1* , not *no odd valuation outside the frozen set*. This distinction is load-bearing: 197 of the 293 grid closures have exactly one odd-valuation outside prime whose symbol is forced $+1$ by reciprocity. The stronger reading would not follow from prime values, since a prime value deliberately creates one odd place.

8.2. The theorem. Fix a verified aligned class with data $(a, \varepsilon, f, q_1, N)$, put $Q(t) = q_1 + Nt$, $b(t) = \varepsilon f Q(t)$, and let $S = \{2, 3, 5, 7\} \cup \text{supp}(\alpha) \cup \text{supp}(\delta) \cup \text{supp}(f)$. Write the composed class polynomial as

$$F(t) = P(\varepsilon f(q_1 + Nt)) = cG(t),$$

with $c > 0$ and $G \in \mathbb{Z}[t]$ primitive with positive leading coefficient. The relevant hypothesis on c is that its *square class* is supported on S . Literal S -support is false in general: for $(w, z) = (3, 3/11)$, $a = 1$, $q_1 = 19$ one gets $c = 9072/15692141883605$, whose outside part is 11^{-12} . Only the square class matters, because an outside place of even valuation contributes Hilbert symbol $+1$.

Theorem 8.1 (Classical Schinzel H supplies emergent-free members; PROVED implication). *Assume the class certificate (all symbols at S , at the moving prime Q , and at infinity equal $+1$ throughout the progression), assume the square class of c is supported on S and $G(t)$ is a p -adic unit for every $p \in S$, and assume Q and G are irreducible with positive leading coefficients whose product has no fixed prime divisor. Then Schinzel's Hypothesis H for $\{Q, G\}$ implies that the class contains infinitely many members with ladder verdict zero.*

Proof sketch. Schinzel supplies infinitely many t with $Q(t)$ and $G(t)$ simultaneously prime; the side conditions (Cauchy bound, $b \neq 0, 1$, and finitely many forbidden coincidences) exclude only finitely many t . Put $R = G(t)$. The part of c outside S need not be trivial, but every one of its valuations is even. At the moving prime, $b \equiv 0 \pmod{Q}$ and the kernel identity gives $P(b) \equiv -\delta a^4 Z^4 A^2 \pmod{Q}$, a unit, so $R \neq Q$. Consequently R is the unique additional outside place of odd valuation; every other outside place has even valuation and $v_p(d) = 0$, hence symbol $+1$. The certificate pins the symbols at S , at Q , and at infinity to $+1$. Hilbert reciprocity forces $(x_0, d_0)_R = +1$. Thus no place carries symbol -1 , and the ladder verdict is *zero*. $\square$

8.3. Uniform admissibility status. For the class of Theorem 5.1, let $L(t) = q_1 + Nt$ and normalize $P(fL(t)) = cG(t)$ with $c > 0$ and $G \in \mathbb{Z}[t]$ primitive. The exact status is as follows.

For (a), the constant and leading coefficients of P are $P_0 = P_8 = 4a^8 AZ^4 > 0$, and the affine substitution $b = f(q_1 + Nt)$ preserves factorization over $\mathbb{Q}$. Thus G has positive lead and

$$G \text{ irreducible over } \mathbb{Q} \iff P \text{ irreducible over } \mathbb{Q}.$$

Theorem 9.1 proves the vertical polynomial irreducible for every fixed $Z \in \mathbb{Q}^\times$, and Corollary 9.2 selects a good a in the required character progression. Thus the fixed cell, not merely the generic two-variable family, is licensed.

For (b), the prime support of N is exactly S , while the CRT base q_1 is a unit modulo every prime in S . For (c), primes in S never divide either value by the Taylor congruences. If $p \notin S$, the linear factor has one root modulo p ; a fixed divisor would force the degree-8 polynomial G to vanish at

condition	requirement	uniform status
(a)	irreducible polynomials with positive leading coefficients	PROVED for every fixed $Z \in \mathbb{Q}^\times$; Hilbert selection in the admissible a progression PROVED
(b)	$\gcd(q_1, N) = 1$	PROVED
(c)	$L(t)G(t)$ has no fixed prime divisor	PROVED
(d)	$G(t)$ is a p -adic unit at S and the frozen square classes are constant	PROVED

TABLE 5. Admissibility of the uniformly constructed square-branch classes.

the other $p - 1$ residues, hence $p \leq 9$. Every such prime already lies in S , a contradiction. For (d), the Taylor-exponent construction gives coefficientwise

$$G(t) \equiv G(0) \not\equiv 0 \pmod{p} \quad (p \in S \text{ odd}), \quad G(t) \equiv G(0) \not\equiv 0 \pmod{8}$$

and the analogous congruences for $L(t)$. The resulting ratios are local squares, so every frozen Hilbert symbol is constant for all integer t .

The modular engine remains an exact finite cross-check. The L20 class audit certified 353/353 independently reconstructed grid classes and 706/706 legacy cell-branch rows, while the L21 audit certified all 353/353 recorded target fibers. The L22 endpoint-Newton theorem is strictly stronger in scope: it proves the vertical statement for every fixed nonzero rational Z , without extrapolating from those finite rows.

A separate 24-row cross-check re-ran the proven-primality kernel end to end: every sampled member had one proved-prime ladder residual with symbol +1 and verdict *zero*.

Remark 8.2 (Exact scope). Schinzel supplies members only after a chosen class satisfies every row of Table 5. Every one of those rows is now PROVED on the fixed canonical branch: Theorem 9.1 and Corollary 9.2 supply irreducibility, while L20 supplies normalization, positivity, coprimality, absence of a fixed divisor, and frozen local data. Therefore the implication has the exact form

$$\text{classical Schinzel H} \implies \text{the conditional six-quantifier conclusion.}$$

Schinzel H is the sole conjectural input. The conclusion remains conditional: this argument neither proves Schinzel H nor gives an unconditional solution of Hilbert's Tenth Problem over $\mathbb{Q}$.

9. FIXED-BRANCH IRREDUCIBILITY FOR EVERY VERTICAL FIBER

This section closes the algebraic premise needed in Section 8. Retain

$$A = 1 + 4a^2, \quad s = \frac{a-1}{2}, \quad Z = z^3, \quad D = 1 - Z - a^2Z^2,$$

$$N_g(b) = 16a^4b^2 - A(b-1)^4$$

and

$$(1) \quad P(b) = 16D^2A^2b^4 - \delta_\tau a^4Z^4N_g(b)^2 - 32A^3s^2D^2b^5.$$

The class construction uses only the fixed canonical branch

$$\tau^\dagger = \frac{1+2a^2}{A}, \quad \delta_{\tau^\dagger} = -\frac{4a^4}{A}.$$

On this branch multiplication by the nonzero scalar $A/4$ gives

$$(2) \quad H := \frac{A}{4}P = a^8Z^4N_g(b)^2 + 4A^3D^2b^4 - 2A^4(a-1)^2D^2b^5.$$

Its constant and leading coefficients are equal:

$$(3) \quad H_0 = H_8 = L = a^8 A^2 Z^4.$$

9.1. The off-branch warning. The branch qualification is essential. If $a = 1$ and $\delta_\tau = \sigma^2 \in \mathbb{Q}^{\times 2}$, then

$$P(b) = (4DAb^2 - \sigma a^2 Z^2 N_g(b))(4DAb^2 + \sigma a^2 Z^2 N_g(b)),$$

a genuine quartic-by-quartic factorization whenever the displayed endpoints are nonzero. This does not meet the canonical branch, because $\delta_{\tau^\dagger} = -4a^4/A < 0$ for rational $a \neq 0$. Thus no blanket assertion over a moving τ is used below: every irreducibility statement is about the fixed pair $(\tau^\dagger, \delta_{\tau^\dagger})$.

9.2. Endpoint and Newton elimination.

Theorem 9.1 (Fixed- Z canonical-branch irreducibility; PROVED). *Fix $Z \in \mathbb{Q}^\times$. On*

$$\tau = \tau^\dagger = \frac{1 + 2a^2}{1 + 4a^2}, \quad \delta = -\frac{4a^4}{1 + 4a^2},$$

the polynomial H of (2), and hence P , is irreducible of degree eight in $\mathbb{Q}(a)[b]$.

Proof. We first assume $Z \notin \{0, 1\}$. The polynomial H is primitive in $\mathbb{Q}[a][b]$. Indeed, a common divisor of its coefficients must divide the equal endpoints $a^8 A^2 Z^4$. Since $A = 1 + 4a^2$ is irreducible in $\mathbb{Q}[a]$ and coprime to a , only a and A could occur. The coefficient of b^4 at $a = 0$ is $4(1 - Z)^2 \neq 0$, excluding a , and direct reduction of that coefficient modulo A excludes A . Gauss's lemma therefore turns any factorization over $\mathbb{Q}(a)$ into one by primitive factors in $\mathbb{Q}[a, b]$.

At the prime A , the coefficient valuations in ascending powers of b are

$$(2, 2, 1, 1, 0, 1, 1, 2, 2).$$

The Newton polygon consequently has two sides, each of horizontal length four, with slopes $-1/2$ and $+1/2$. The slopes belonging to a global factor form a sub-multiset of these slopes. Its endpoint valuations are integers, whereas the sum of an odd number of half-integral root valuations is not. Every global factor therefore has even degree. If the octic were reducible, it would have a factor of degree two or four.

At $a = 0$ the coefficient valuations are

$$(8, 8, 8, 8, 0, 0, 8, 8, 8),$$

and the sides have data

$$4@(-2), \quad 1@0, \quad 3@(8/3).$$

For the four roots in the first cluster, substitution $b = a^2 x$ gives the residual polynomial

$$(4) \quad R_0(x) = Z^4 + 4(1 - Z)^2 x^4.$$

It has no rational linear factor, since its two terms are nonnegative on $\mathbb{Q}$ and do not vanish simultaneously. It therefore has no rational cubic factor either. The denominator three in the last slope makes its three roots an indivisible local block. Hence a degree-two factor must take two roots from the first four-root cluster; a degree-four factor either takes all four of those roots or takes the unit root together with the three-root cluster. In a $4 + 4$ factorization one of the factors is therefore the all-small factor.

The two exact residual limits at infinity are

$$(5) \quad a^{-12} H(a, x/a) \longrightarrow 16Z^4(4x^2 - 1)^2,$$

$$(6) \quad a^{-20} H(a, ax) \longrightarrow 16Z^4 x^4 (x^2 - 4)^2.$$

Let F be a primitive factor of degree $r \in \{2, 4\}$ formed from $a = 0$ small roots. Unique factorization and (3) force

$$\text{lc}_b(F) = cA^\beta, \quad F(0) = da^{2r}A^\delta,$$

with $c, d \in \mathbb{Q}^\times$ and $0 \leq \beta, \delta \leq 2$. Fixed powers of the fixed rational Z are units and are absorbed into c, d . If ℓ of the roots of F are large at infinity, comparison with (5)–(6) gives

$$2r + 2\delta - 2\beta = 2\ell - r.$$

The integral possibilities are

r	(β, δ, ℓ)
2	$(1, 0, 2), (2, 0, 1), (2, 1, 2),$
4	$(2, 0, 4).$

Suppose first that $r = 2$. The rational degree-two divisors of the repeated clusters in (5)–(6), in the three endpoint cases just listed, force the constant-to-leading ratio of the $a = 0$ residual factor to be

$$\rho = d/c \in \{16, -16\}.$$

After monic normalization, divisibility of (4) has the form

$$(x^2 + ex + \rho)(x^2 - ex + q) = x^4 + K, \quad K = \frac{Z^4}{4(1-Z)^2} > 0.$$

Coefficient comparison gives $e(q - \rho) = 0$ and $q - e^2 + \rho = 0$. If $e = 0$, then $q = -\rho$ and $K = -\rho^2 < 0$. Otherwise $q = \rho$ and $e^2 = 2\rho$, which asks for a rational square root of 32 or of -32 . Both alternatives are impossible, so no quadratic factor exists.

Now let $r = 4$. The endpoint enumeration gives $(\beta, \delta, \ell) = (2, 0, 4)$, so

$$\text{lc}_b(F) = cA^2, \quad F(0) = da^8.$$

The $b = ax$ initial form is the whole nonzero-root cluster $16c(x^2 - 4)^2$, and hence $d = 256c$. At $b = a^2x$, the complementary factor contributes the constant Z^4/d . Matching the coefficient of x^4 in (4) gives

$$\frac{cZ^4}{d} = 4(1 - Z)^2, \quad Z^4 = 1024(1 - Z)^2.$$

The last equation splits into

$$Z^2 + 32Z - 32 = 0, \quad Z^2 - 32Z + 32 = 0.$$

Their discriminants are respectively $1152 = 576 \cdot 2$ and $896 = 64 \cdot 14$, neither a rational square. Thus no degree-four factor exists.

It remains to treat the degenerate residual value $Z = 1$. At $a = 1$ the primitive ascending coefficient vector of H is

$$(25, -200, 540, -760, 1546, -760, 540, -200, 25).$$

Modulo 11, monic normalization gives

$$q(b) = 1 + 3b + 4b^2 + 7b^3 + 2b^4 + 7b^5 + 4b^6 + 3b^7 + b^8.$$

In $\mathbb{F}_{11}[b]/(q)$ the exact Rabin certificate is

$$b^{11^8} = b, \quad b^{11^4} = 8 + 7b + 4b^2 + 9b^3 + 4b^4 + 7b^5 + 8b^6 + 10b^7,$$

and $\gcd(b^{11^4} - b, q) = 1$. Since two is the only prime divisor of eight, q is irreducible. Localizing $\mathbb{Q}[a]$ at $L = a^8A^2$ makes H/L monic; a putative factorization has monic factors in that localization, and evaluation at $a = 1$ preserves their positive degrees because $L(1) = 25 \neq 0$. The certificate is therefore a contradiction, proving the vertical statement at $Z = 1$.

Finally, $Z = 0$ gives

$$H = 2A^3b^4(2 - A(a - 1)^2b),$$

which is reducible and is excluded because the cell parameter z is nonzero. For fixed $Z \neq 0$, the equation $D = 0$ removes at most two special values of a ; it does not affect irreducibility in $\mathbb{Q}(a)[b]$. This completes all rational fixed fibers in the theorem. $\square$

9.3. Hilbert selection inside the admissible progression.

Corollary 9.2 (Uniform admissible specialization; PROVED). *For every odd cell prime w and every $z \in \mathbb{Q}^\times$ with $v_w(z) \geq 1$, the existing odd- a character progression contains an a for which the canonical-branch polynomial $P(a, z^3; b)$ is irreducible in $\mathbb{Q}[b]$ and $D \neq 0$.*

Proof. Fix $Z = z^3 \neq 0$. Theorem 9.1 makes the octic over $\mathbb{Q}(a)$ irreducible and, in characteristic zero, separable. Quantitative Hilbert irreducibility puts the integers at which its degree drops or it becomes reducible in a thin set E_Z satisfying

$$(7) \quad \#\{n \in E_Z \cap \mathbb{Z} : |n| \leq B\} = O_Z(\sqrt{B} \log B).$$

[9][15, Section 13, Theorems 1–2]

The character sum used in the class construction supplies $r \bmod w$ with

$$\left(\frac{1 + 4r^2}{w} \right) = -1.$$

Combining this residue with parity gives an odd $a_0 \bmod 2w$; every positive $a = a_0 + 2wk$ is character-admissible. The progression contributes linearly many integers up to height B , which eventually dominates (7). It therefore contains a point outside E_Z , and the equation $D = 0$ excludes at most two more points. Such an a has the asserted properties. $\square$

This is the load-bearing order of the closure. Fixed- Z vertical irreducibility gives Hilbert selection *inside the already admissible a progression*; the resulting L20 class certificate has $f = w$ and all Schinzel-pair conditions; the L19 fixed-pair implication applies classical Schinzel H to $\{q_1 + Nt, G(t)\}$; the supplied prime pair gives an emergent-free member; and the L6 equivalence yields the six-variable definition. No separate per-cell irreducibility premise remains.

9.4. Independent reciprocal-trace corroboration.

Theorem 9.3 (The fixed reciprocal specialization; PROVED). *On the same canonical branch, specialize*

$$a = 1, \quad A = 5, \quad s = 0, \quad \tau^\dagger = 3/5, \quad \delta_{\tau^\dagger} = -4/5.$$

For every $Z \in \mathbb{Q}^\times$, the resulting polynomial $P(1, Z; b)$ is irreducible in $\mathbb{Q}[b]$.

Proof. For this fixed choice $(a, A, s, \tau, \delta) = (1, 5, 0, 3/5, -4/5)$, put $D = 1 - Z - Z^2$ and $u = b + b^{-1}$. Direct expansion gives

$$P(b) = b^4 T(u), \quad T(u) = 400D^2 + \frac{4}{5}Z^4(16 - 5(u - 2)^2)^2.$$

In $E = \mathbb{Q}(t)$, $t^2 = -5$, one has

$$\frac{5}{4}T = Q_- Q_+, \quad Q_\pm = Z^2(16 - 5(u - 2)^2) \pm 10Dt.$$

The conjugate quadratics are coprime, and trace reducibility is equivalent to $80Z^2 + 50Dt$ being a square in E . Its norm makes

$$125D^2 + 64Z^4$$

a necessary rational square. Writing $Z = p/q$ in lowest terms and $d = q^2 - pq - p^2$, the cleared numerator is $125d^2 + 64p^4 \equiv 5 \pmod{8}$: coprimality makes d odd in all three parity patterns. It cannot be a square, so T is irreducible.

For a root u of T , the reciprocal lift is irreducible exactly when $u^2 - 4$ is not a square in $\mathbb{Q}(u)$. Its norm is a rational square times

$$\Xi(Z) = (125D^2 + 64Z^4)(125D^2 + 1024Z^4).$$

Suppose $\Xi(Z)$ were a square and put $h = 5D/(8Z^2)$. The resulting point on

$$v^2 = (5h^2 + 1)(5h^2 + 16)$$

maps by

$$x = 25h^2, \quad y = 25hv$$

to

$$E_0 : y^2 = x(x + 5)(x + 80).$$

The complete 2-isogeny descent checks the square classes

	candidate classes	realized classes
E_0	$\{\pm 1, \pm 2, \pm 5, \pm 10\}$	$\{1, -5\}$
$E'_0 : y^2 = x(x - 45)(x - 125)$	$\{\pm 1, \pm 3, \pm 5, \pm 15\}$	$\{1, 5\}$.

The excluded classes are eliminated modulo 3, modulo 16, or at the real place, so the descent gives rank zero. The good-prime point counts 8, 16, and 12 at 7, 11, and 13 force torsion order at most four; the visible points already give

$$E_0(\mathbb{Q}) = \{O, (0, 0), (-5, 0), (-80, 0)\}.$$

But $D \neq 0$ for rational Z and the image has $x = 25h^2 > 0$, a contradiction. Thus $\Xi(Z)$ is nonsquare and the reciprocal lift is irreducible. $\square$

The value $a = 1$ in Theorem 9.3 is a *specialization witness* for the rational-function fiber. It need not satisfy $\left(\frac{5}{w}\right) = -1$ and is not the class parameter selected in Corollary 9.2. Localizing at $a^8 A^2 Z^4$ shows that this theorem also gives an independent proof of fixed- Z vertical irreducibility; the endpoint–Newton proof above remains the primary closure.

9.5. Route discipline. Three structurally useful routes are deliberately not cited as the closure. The infinity calculation alone resolves four quadratic local clusters but still permits global factor degrees 2, 4, and 6; the equal-endpoint comparison in the primary proof is what eliminates them. A factor-tuple version of Schinzel controls only the product of the outside Hilbert signs unless an all-plus compatibility class is also proved, so it relocates rather than removes irreducibility. Finally, the free- λ square-branch theorem varies τ through the infinite L11c family; that new branch parameter would cost a seventh witness and is excluded from the six-count. These are proved scope statements about alternate routes, not gaps in Theorem 9.1.

10. UNCONDITIONAL DENSITY, ALGEBRAIC WALLS, AND THE OFF-GRID FRONTIER

This section separates the unconditional analytic gain from the remaining parity problem, records the algebraic and fibration walls found in L23–L24, and then states the finite off-grid horizon.

10.1. An unconditional local bound. Fix an aligned class with parameters (ε, f, q_1, N) and write $F(k) = P(\varepsilon f(q_1 + kN))$ for its degree-8 class polynomial, and $u(k) = 2\alpha \varepsilon f(q_1 + kN)$ for the unit whose Legendre symbol decides bad signature. For an odd prime p let m_p denote the Haar mass in $\mathbb{Z}_p$ of the set of k with $v_p(F(k))$ odd and $\left(\frac{u(k)}{p}\right) = -1$: the local mass of the emergent obstruction.

Theorem 10.1 (Local sieve bound; PROVED). *For every aligned class of the grid,*

$$m_3 = m_5 = m_7 = 0, \quad \text{and} \quad m_p \leq \frac{8}{p+1} < 1 \quad \text{for all } p \geq 11,$$

the bound holding for singular roots as well, counted with multiplicity.

The vanishing at $p \in \{3, 5, 7\}$ is exactly the frozen-alignment condition of the class certificate: those places are controlled, and their symbols are pinned to $+1$ by construction. For $p \geq 11$ the bound is the root count of a degree-8 polynomial: at most eight residues can be roots, each simple root contributes Haar mass $1/(p+1)$ for odd valuation, and a singular root contributes at most its multiplicity times that mass, which is again bounded by the total degree. The bound was verified independently on every stored local pair: 102,439 pairs with $p \leq 10^4$ across the 293 classes, zero violations.

Theorem 10.1 settles a natural hope in the negative, and this is the important structural point.

Corollary 10.2 (No uniform positive mass; PROVED for the current classes). *For each of the 293 classes the splitting condition governing the bad-signature event has Chebotarev exponent $c = 1/2$: the class polynomial is irreducible with a replayed degree-8 certificate, and a replayed witness shows the bad unit is a nonsquare at a simple bad root. Consequently the truncated clean product obeys*

$$\prod_{p \leq X} (1 - m_p) \asymp (\log X)^{-1/2},$$

so no cutoff-uniform positive lower bound on the clean mass exists: the small-layer clean probability decays, slowly, to zero.

The measured exponent agrees: the root-harmonic estimate gives $c \approx 0.4778$, the product estimate $c \approx 0.4801$, and the same-member sieve between 10^4 and 10^5 gives $c \approx 0.4590$ (EVIDENCE).

The consequence for the member property $\mathcal{M}$ is a sharpening, not a defeat. The number of admissible prime members up to height X in a class grows like $X/(\log X)^2$ by the Dirichlet–Bateman–Horn heuristic [1], while each carries clean probability $\asymp (\log X)^{-1/2}$; the expected count of emergent-free members therefore still diverges, at rate $X/(\log X)^{5/2}$. What Corollary 10.2 rules out is the clean shortcut — a uniform positive constant — and what it leaves is a divergence statement of exactly the shape the measured counting law already exhibits (first emergent-free member at $k = 0$ in 207 of 293 classes).

10.2. The L23 half-sieve and its exact parity wall. Write the primitive L20 linear–octic pair as

$$Q(t) = q_1 + Nt, \quad G(t),$$

and let a prime p be *bad-root* at t when $G(t) \equiv 0 \pmod{p}$ and the associated unit $u(t)$ has negative quadratic character at p . Finite exceptional primes are first put into a compatible fixed CRT class.

Theorem 10.3 (Half-dimensional small-prime sieve; PROVED). *For every fixed admissible L20 pair, the bad-root sieve has dimension $\kappa = 1/2$:*

$$\sum_{p < z} g_{\text{root}}(p) \log p = \frac{1}{2} \log z + O(1).$$

Bombieri–Vinogradov supplies level

$$D = \frac{X^{1/2}}{(\log X)^B}.$$

Consequently the semilinear beta sieve, with $z = X^{49/100}$, gives

$$\# \left\{ X < t \leq 2X : \begin{array}{l} Q(t) \text{ is prime,} \\ G(t) \text{ has no bad root prime } p < z \end{array} \right\} \gg \frac{X}{(\log X)^{3/2}}.$$

The implied constant may depend on the fixed class.

These are *small-prime-clean members of the progression only*. The theorem does not say that they are emergent-free members of $\mathcal{M}$. Indeed $|G(t)| = X^{8+o(1)}$, so after sieving only to $X^{49/100}$

a value may retain as many as 16 large prime factors, with multiplicity. Reciprocity says that the number of large odd-valuation bad factors is even, leaving

$$0, 2, 4, \dots, 16,$$

not just 0. Even Elliott–Halberstam level would leave as many as eight factors. Kao’s degree-eight prime-argument theorem reaches only P_{12} , whereas reciprocity would close the argument only at $R_{\text{bad}} \leq 1$.

Thus the first missing analytic input is a parity-sensitive saving for the sector with two large bad odd-valuation divisors $p_1, p_2 \geq X^{1/2-\epsilon}$ and $p_1 p_2 > D$, beyond the Bombieri–Vinogradov range. A finite no-hit scan cannot supply that estimate.

10.3. The L23 fibration and norm walls. On the canonical branch the tied quaternion has the conic-bundle model

$$U^2 - 2bV^2 = P_{a,Z}(b)W^2.$$

For irreducible P , its closed degenerate-fibre degrees are $1 + 8 + 1$, at $b = 0$, $P(b) = 0$, and $b = \infty$. On the refined L20 choice $a \not\equiv 1 \pmod{w}$, the target reduction has the simple root

$$\beta = (2As^2)^{-1}, \quad 2\beta = (As^2)^{-1},$$

which is a nonsquare because $(\frac{A}{w}) = -1$. Such a refinement is available from the exact general character count

$$\#\{a \bmod w : \left(\frac{1 + 4a^2}{w}\right) = -1\} = \frac{w - (\frac{-1}{w})}{2}.$$

If $v_w(z)$ is odd, the left Newton edge gives an odd valuation for $2b$ for every admissible a . Hence 2θ is nonsquare in $\mathbb{Q}(\theta)$, where $P(\theta) = 0$, throughout these scopes.

It follows that the natural relatively minimal bundle has non-split rank 10, not rank at most 2, and

$$\text{Br}(X)/\text{Br}(\mathbb{Q}) \cong \mathbb{Z}/2\mathbb{Z}, \quad \text{generated by } (A, b).$$

Thus the Harpaz–Wei–Wittenberg rank- ≤ 2 theorem and the checked low-rank variants do not apply. The pre-existing fixed-class stratum with $v_w(z)$ even and $w \mid s$ remains OPEN without an exact row certificate; this does not obstruct the global class construction, because the free L20 refinement and the L22 Hilbert selection avoid that stratum.

The accompanying exact algebra rules out the most direct section escapes. Exact norm matching is Φ -empty by its 2-adic valuation, and there is no polynomial-in- b norm section. The loci $P = \pm 2by^2$ are Φ -empty, while writing P as A times a norm forces the 2-adic symbol to be -1 . A fixed P -square construction leads to a genus-one locus. The first Φ -admissible factor ansatz,

$$L = -A\rho^2,$$

does factor H , but it reduces the problem to scoped bad Hilbert symbols and genus-four curves rather than producing a section. Fixed nonsquare twists merely relocate the quadratic character.

When a is allowed to move with the prime parameter, the smallest squarefree representative actually certified in the analyzed diagonal has degree 18 (factor degrees $2 + 16$); raw degree-21 examples are not a minimum, and no general minimum 21 is claimed.

10.4. The L24 diagonal route is refuted. The two self-coupled diagonal orientations define exact finite-flat degree-8 covers and would have tied the extra rank at most one. They therefore looked like a possible route to a five-unknown definition. Both images are, however, uniformly empty on Φ over $\mathbb{Q}_2$.

For orientation I the normalized eliminant has coefficient valuations $(4, 0, 0)$. Its roots have $v_2(u) = 4$ or 0 ; if u is a square, then $A + u$ is respectively 5 or $6 \bmod 8$, never a square. For orientation II the eliminant is

$$u^2 - 32hu + 16e = 0, \quad h, e \in \mathbb{Z}_2^\times.$$

Every root has $v_2(u) = 2$. Substituting $u = 4t^2$ gives the complete modulo-16 contradiction

$$8 + 2Ab(s^2 - 1) \not\equiv 0 \pmod{16}.$$

Thus the putative five-count is false, not OPEN.

The obstruction is genuinely dyadic: at every odd target place, orientation II is locally soluble in the standard $k = 1 < 3v_w(z)$ stratum. Geometrically the generic arithmetic monodromy is $V_4 \wr C_2$ of order 32; the audited slices have genera 35 and 53, and neither orientation has a base-rational section. The 4,026,282-attempt search found no rational hit, but that finite zero-hit count is EVIDENCE only; the uniform negative theorem is the preceding 2-adic proof.

10.5. The L25 scaled couplings are not dyadically killed. L24 kills the two *unit* self-couplings. The next family changes only fixed rational multiples of the tied norm witnesses and introduces no new free variable. On the square branch

$$X^2 - \rho^2 = A, \quad u = \rho^2,$$

the type-I coupling $(y, r) = (\kappa_1 X, \kappa_2 \rho)$ eliminates to

$$A\kappa_1^2 u^2 + (A^2 \kappa_1^2 - c^2 - 16AB\kappa_2^2)u + 16A^2 B s^2 - 16A = 0.$$

The identity, and the type-II analogue, were replayed on 1,200 exact instances.

Theorem 10.4 (Scaled dyadic escape; PROVED). *On Φ , the fixed coupling $(y, r) = (2X, \rho)$ has a 2-adically admissible branch exactly when s is even, and the fixed coupling $(y, r) = (2X, \rho/2)$ has a 2-adically admissible branch exactly when s is odd. Both statements are uniform in odd b , in the declared scalar unit classes, and in every c with $v_2(c) \geq 4$. Hence their two-formula disjunction is nonempty on every Φ stratum.*

Proof outline. After dividing by $4A$, put $u = 4t^2$. In each case the unit quartic has the form

$$g(t) = 4\mu^2 t^4 + Lt^2 + M_0, \quad M_0 = 2Abs^2 - 1,$$

with $L, M_0 \in \mathbb{Z}_2^\times$. For odd t , $v_2(g'(t)) = 1$, so a root modulo 8 lifts to a 2-adic root. The two reductions are

$$g(1) \equiv 2Abs^2 \pmod{8}, \quad g(1) \equiv 2b(As^2 - 1) \pmod{8}$$

for $\kappa_2 = 1$ and $1/2$, respectively. The first vanishes exactly for even s ; the second exactly for odd s . The other possible root valuation of the original u -quadratic is 0, which makes $A + u \equiv 6 \pmod{8}$, so the quartic condition is also necessary. The declared residue classes were all replayed machine-exactly. $\square$

At an odd target prime w , choose the freely chosen unit- b unramified reduction $c \equiv 0 \pmod{w}$ and the residues

$$(s, A, X, \rho, b) \equiv (0, 5, 3, 2, \kappa^{-2}), \quad \kappa \in \{1, 1/2\}.$$

Both scaled covers hold modulo w , and the (ρ, X) -Jacobian determinant is the nonzero integer

$$-16AX\rho(X^2 + \rho^2 - 4B\kappa^2) = -2400.$$

Reduction modulo every odd $w \geq 7$ is smooth, so Hensel's lemma supplies a w -adic point. At $w = 3, 5$ the completed bounded scans find mod- w points but no smooth point, so this streamlined proof has a named small-prime exception; it does not prove target local emptiness. This is not the L24 low-positive aligned stratum, and compatibility with a preselected L20 residue class remains separate.

Over $\mathbb{R}$, the two guarded bridge samples

$$(a, b, Z) = (1, 3, -10), \quad (3, 4, -2/5)$$

have $A_2 > 0$, $L_2 < 0$, and a nonnegative discriminant, hence a positive root u ; thus u and $A + u$ are positive reals. These are named certificates, not an all-parameter real theorem. The crucial

limitation remains global: a rational coupled member requires the quadratic in u to have a rational root and both $u, A + u$ to be rational squares. The bounded search is zero-hit EVIDENCE only and is not used as proof.

10.6. Two new non-diagonal constructions. The preceding scalings do not exhaust the available zero-cost specializations. Two further constructions sharpen different sides of the frontier.

Theorem 10.5 (Reciprocal even-pullback tie; PROVED). *Put $Z = z^2$, $D = 1 - Z - a^2 Z^2$, and tie the third block coordinate to*

$$\eta = \frac{Z(b+1)}{Db}.$$

On the canonical bridge domain $b \neq 0, 1$, this adds no guard beyond the existing guards. If

$$M = 16 - \delta_\tau c^2 - 32Ab\eta^2, \quad P_{\text{rec}} = D^2 A^2 b^4 M,$$

then

$$P_{\text{rec}}(b) = b^4 T_\tau(b + b^{-1}),$$

where

$$T_\tau(u) = 16A^2 D^2 - \delta_\tau a^4 Z^4 [16a^4 - A(u-2)^2]^2 - 32A^3 Z^2(u+2)$$

has degree 4. On $\tau^\dagger$, P_{rec} is a $\mathbb{Q}_2$ -square throughout Φ , and every standard guarded W1 target lifts. Away from the fixed $b = -1$ primes,

$$\left(\frac{2b}{p}\right) = \left(\frac{2(u+2)}{p}\right).$$

Proof outline. The identity follows from

$$N_g(b) = b^2 [16a^4 - A(b + b^{-1} - 2)^2]$$

and $b^3(b+1)^2 = b^4(b+b^{-1}+2)$. For $r = v_2(Z)$, one has $v_2(D) = 0$ for $r \geq 0$ and $2r$ for $r < 0$; hence $v_2(\eta) \geq 1$, $v_2(c) \geq 4$, and $M/16 \in 1 + 8\mathbb{Z}_2$. At a standard odd target, $v_w(\eta) = v_w(Z) - 1 \geq 1$, $v_w(c) = 2v_w(Z) - 2 \geq 2$, and $M \equiv 16 \pmod{w}$. Finally $(u+2) = (b+1)^2/b$ proves the character identity. $\square$

Theorem 10.6 (Uniform reciprocal trace field; PROVED). *On the canonical L26 branch, for every $v_2(a) = 0$ and $Z = z^2 \neq 0$, the trace quartic T is irreducible over $\mathbb{Q}_2$. If θ is a root, then*

$$N_{\mathbb{Q}(\theta)/\mathbb{Q}}(2(\theta+2)) \in A\mathbb{Q}_2^{\times 2}, \quad A \equiv 5 \pmod{8}.$$

Consequently $2(\theta+2)$ is nonsquare and $T(v^2/2-2)$ is irreducible over $\mathbb{Q}_2$ and $\mathbb{Q}$.

Proof outline. After $w = u - 2$, the nonzero coefficients of $T(w+2)$ have valuations

	d_0	d_1	d_2	d_4	
$t \geq 0$	4	$5+2t$	$7+4t$	$2+4t$	$t = v_2(Z)$.
$t < 0$	$4+4t$	$5+2t$	$7+4t$	$2+4t$	

The one-segment Newton polygon has half-integral slope, excluding linear factors. For the monic form $w^4 + pw^2 + qw + r$, a quadratic factor forces a square Y on

$$Y^3 + 2pY^2 + (p^2 - 4r)Y - q^2 = 0.$$

Its Newton polygon leaves five possible valuation cases; after normalization their residues are $3, 2, 4, 4, 2 \pmod{8}$, excluding every square Y . Finally

$$N(2(\theta+2)) = \frac{64\{A^3 D^2 + 64a^8 Z^4(a^4 - A)^2\}}{a^8 Z^4 A^2}.$$

Because $v_2(a^4 - A) = 2$, the braced expression divided by $A^3 D^2$ lies in $1 + 2^{10}\mathbb{Z}_2$; the norm has squareclass A , and Capell finishes. $\square$

Theorem 10.7 (Reciprocal-lift classification and compression; PROVED). *In the setting of Theorem 10.6, put $t = v_2(Z) \in 2\mathbb{Z}$. Then*

$$\theta^2 - 4 \in \mathbb{Q}_2(\theta)^{\times 2} \iff t = -2 \text{ or } t \geq 2.$$

Moreover

$$Z = \frac{8z^2}{1+z^2}$$

preserves the positive-valuation locus at every odd prime and has $v_2(Z) \geq 2$ for every $z \in \mathbb{Q}^\times$. Thus this rational pullback forces the locally split reciprocal stratum without changing the odd target set.

Proof outline. For $w = \theta - 2$ and $t \geq 2$, $\theta^2 - 4 = w^2(1 + 4/w)$ with $v_2(4/w) > 2$, so strong Hensel applies. For $t \leq 0$, put $b = 1 + x$. The monic shifted reciprocal octic has one Newton slope $-1/4$, so a proper factorization must be two Eisenstein quartics. Exhaustive necessary factor congruences modulo 16 and 64 exclude $t = 0$ and $t \leq -4$.

At $t = -2$, for $\phi = x^4 + 6x^3 + 6$, the resultant

$$\mathcal{R}(Y) = \text{Res}_x(S(x), Y - \phi(x))$$

has Newton vertices $(0, 21), (4, 10), (8, 0)$, hence two slopes. If S were irreducible, all conjugates $\phi(\sigma\alpha)$ would have the same valuation, forcing a one-slope resultant. Therefore S factors $4 + 4$, equivalently $b^2 - \theta b + 1$ splits in the trace field.

Finally, direct valuation in the three cases $v_p(z) > 0, = 0, < 0$ proves the odd-place equivalence for the compressed pullback. At 2, its valuation is $3 + 2v_2(z)$, 2, or 3, respectively. $\square$

Remark 10.8 (Fixed-field slice). On $b = w\rho^2$, the member symbol is $(M, 2w)$, so the moving quadratic field is frozen. The resulting conic bundle has no generic section: at $\rho = 0$, its squareclass is A and $(A, 2w)_w = (A|w) = -1$. Exact soluble fibres are evidence only; the inert-prime parity problem remains open.

Theorem 10.9 (One-piece aligned shear; PROVED local theorem). *On the square branch $X^2 - \rho^2 = A$, the fixed shear*

$$(y, r) = (2X + 28\rho, sX + \rho)$$

has determinant $2(1 - 14s)$, nonzero on Φ . It has a $\mathbb{Q}_2$ -point for every $s \in \mathbb{Z}_2$. Moreover, on the standard aligned target stratum

$$v_w(B) = 1, \quad \left(\frac{A}{w}\right) = -1, \quad v_w(c) \geq 4,$$

it has a smooth $\mathbb{Q}_w$ -point for every odd prime w .

Proof outline. With $\rho = 2t$ and $X(t)^2 = A + 4t^2$, the normalized equation is

$$788t^4 + 56t^3X + \left(A - 4B(s^2 + 1) - \frac{c^2}{4A}\right)t^2 - 4BsXt - 1 = 0.$$

At $t = 1$ its value is $0 \pmod 8$, while its derivative has valuation exactly 1; strong Hensel applies uniformly in s .

At an odd target, put $x = \rho^2$ and $\rho(2X + 28\rho) = 4\epsilon$, $\epsilon = \pm 1$. Then

$$N_{0,\epsilon} = 195x^2 - 56\epsilon x + 4 = Ax, \quad N_{1,\epsilon} = N_{0,\epsilon} - x = (A - 1)x.$$

For $\epsilon = 1$, the indicator

$$\frac{(1 + \chi(x))(1 + \chi(N_1))(1 - \chi(N_0))}{8}$$

and the Weil bounds for two cubic, one quartic and one quintic character sum give at least

$$\frac{w - 11\sqrt{w} - 26}{8} > 0$$

smooth residues for $w \geq 197$. The correction 26 removes the two possible singular x -values and the zero-root weights. Exact exhaustion over both signs supplies a nonzero-Jacobian row for all 43 odd primes below 197. $\square$

The exact $\lambda = X + \rho$ eliminant has degree 8, leading coefficient $225A$, and constant coefficient $169A^5$. The guarded sample $(a, b, Z) = (1, -1, -10)$ supplies one real point by a sign change. Writing $m = \lambda^2$, the coefficient linear in B factors as

$$-16ABm(m - A)((s + 1)^2m - (s - 1)^2A),$$

excluding the natural cancellation section. Direct dyadic reduction also gives $C_2 \equiv 128 \pmod{256}$ at $\lambda = \pm 1, \pm A$. Finally $y = 0$ forces $(2a)^2 - 195\rho^2 = -1$, which has no $\mathbb{Q}_3$ -point. These are three scoped section no-gos, not a no-point theorem for the full cover.

Theorem 10.10 (Constant-two quartic; PROVED local theorem). *On the same square branch, fix*

$$(y, r) = (2, sX + \rho).$$

With $m = (X + \rho)^2$, the tied equation is

$$H_2(m) = -(c^2 - 4A)(m - A)^2 - 16ABQ_s(m) - 64Am = 0,$$

$$Q_s(m) = (s + 1)^2m^2 - 2A(s^2 + 1)m + (s - 1)^2A^2.$$

It is quadratic in m , hence even quartic in the hyperbola parameter. Its leading coefficient has 2-adic valuation 2 on every Φ -base. The cover has a $\mathbb{Q}_2$ -point on every Φ -stratum, a fibre-regular target-local point at every odd $w \geq 5$, an exact strong-Hensel fibre at $w = 3$, and a point on one guarded real stratum.

Proof outline. After division by A , put $m = 1 + 8t$ and normalize by 2^8 . The value at $t = 0$ is even because

$$Q_s(1) = -16a^2(s(1 + 2a^2) - a^2(s^2 + 1)),$$

whereas the derivative is odd. Hensel gives $m \in 1 + 8\mathbb{Z}_2$, hence a square.

At an aligned odd target the equations reduce to

$$\rho^2 = 4, \quad X^2 = A + 4.$$

For $w \geq 5$, the conditions

$$\chi(1 + 4a^2) = -1, \quad \chi(5 + 4a^2) = 1$$

have at least

$$\frac{w - 3\sqrt{w}}{4} - 2 > 0$$

solutions for $w \geq 23$, by the Weil bound for the squarefree quartic $(4a^2 + 1)(4a^2 + 5)$. Exact rows close $w = 5, 7, 11, 13, 17, 19$. At $w = 3$, the guarded fibre $(a, b, z) = (5, 3, 3)$ satisfies $v_3(H_2(1)) = 3 > 2v_3(H'_2(1))$, so strong Hensel gives a square root $m \in 1 + 9\mathbb{Z}_3$. The control $(1, 3, 3)$ is empty, showing that the selected a is load-bearing. Finally $(a, b, Z) = (1, -1, 1/8)$ gives positive ρ^2 and X^2 over $\mathbb{R}$. $\square$

The constant-two layer also gives two rigidity statements. Every rational L11c square-branch tie

$$\ell_d = \frac{A - d^2}{2d}$$

has the same target norm field $\mathbb{Q}(\sqrt{2b})$. At $b = 0$, its cleared norm value has even valuation and leading squareclass A ; a norm $Y^2 - 2bR^2$ of even valuation has square leading unit. Therefore no rational $d(b)$ produces a generic norm section. For a general B -independent linear shear, the coefficient of B factors as

$$-16ABmH_L(m),$$

$$H_L = (r + t)^2 m^2 - 2A(t^2 - r^2 + 2s^2)m + A^2(r - t)^2.$$

Its discriminant is $16A^2s^2(s^2 + t^2 - r^2)$. If $s \neq 0$, every rational nonzero root has m/A square, impossible because m is square and $A \equiv 5 \pmod{8}$.

Theorem 10.11 (L31 exact reciprocal member). *On the mandatory fixed-field reciprocal slice, the parameters*

$$(w, a, z, Z, q_0, b, \rho, \lambda) = (13, 3, 13, 169, 1, 13, 14/13, 12)$$

give a globally soluble member.

Proof. The canonical bridge gives

$$M = \frac{2^4 Q}{3^2 37^3 83^2 1033^2}, \quad Q = 14082426920623718389.$$

A recursive Pocklington chain certifies Q prime. Direct Hilbert calculation gives

$$(M, 26)_v = 1 \quad (v = 2, 3, 13, 37, 83, 1033, Q, \infty),$$

and there are no other candidate ramified places. Hasse–Minkowski therefore gives a rational point on $U^2 - 26V^2 = MW^2$. The lift is not implicit:

$$\lambda^2 = 144 = 13(13\rho^2 - 4).$$

□

This is one target, not a uniform member theorem. For $q_0 = r/s$, the general fixed-field slice is the sign-decorated degree-16 binary form

$$G_w(r, s) = s^{16} P_{\text{rec}}(wr^2/s^2).$$

L31 also locates the constant-two cube wall. At

$$(a, c, \lambda, b, Z) = \left(1, -\frac{64}{25}, 3, -\frac{3253}{3125}, \frac{2033125}{6101423}\right)$$

all bridge guards and $H_2(\lambda^2) = 0$ hold, but

$$Z = \frac{5^4 \cdot 3253}{1009 \cdot 6047}$$

is not a cube and c is a 3253-adic unit. The unique constant- c bridge-square section is $c = -64/25$; imposing $Z = z^3$ on its two bridge maps gives the genus-2 curves

$$d^2 = -64z^6 + 96z^3 + 64, \quad d^2 = 96z^6 - 224z^3 + 64.$$

Magma V2.29-9 `RationalPointsGenus2` [13] returns only $(0 : \pm 8 : 1)$ on each curve with completeness flag `true`. Thus this unique constant- c section has no guarded nonzero cube point. The exact external request and intrinsic-help semantics are recorded; no standalone Magma certificate is checked in. Other constant-two sections remain open.

Finally, fix one selected L20/L22 aligned class and a $Q(t)$ -prime member. If R_{bad} counts its remaining odd-valuation outside places with symbol -1 , Hilbert reciprocity makes it even. AP1 asserts that every cell has such a selected class/member with $R_{\text{bad}} \leq 1$. Inside this protocol AP1 is equivalent to the intermediate member hypothesis. The two-large root-pair expansion starts at $pq \geq X^{0.98} > D_{\text{BV}}$ and retains a positive $R_p R_q / 4$ main term. The natural missing theorem is a pointwise fixed-family Buchstab/Hilbert-detector asymptotic with positive zero-bad constant. AP1 and that asymptotic remain open, so classical Schinzel H and the conditional count are unchanged.

10.7. The off-grid horizon. The grid is w prime, $w < 100$. Off-grid closures now exist for

$$w \in \{101, 103, 107, 109, 113, 127, 131, 137, 139, 149, 151, 157, 163, 167, 173, 179\}, \quad u = -1,$$

together with $[131, [5, 1]]$, each certified by an aligned class and an exact ladder verdict *zero*. The routes are the L11 family, the canonical fixed-factor escape, and the non-canonical $a = 7$, $\tau = 0$ family. The last searched open cell, $[179, [-1, 1]]$, is now closed, so every off-grid cell tried to date has a replayed member. This is a finite-horizon statement only.

Two negative or delimiting findings carry structural content.

Theorem 10.12 (The canonical 5-wall; PROVED). *On the canonical escape route ($z = -w$, $a = 1$, $\tau = 3/5$, $f = w$), write $e = v_5(D)$. Then $v_5(x_0) = -3 - 2e$ is odd and $v_5(d_0) = 0$, so*

$$\text{Hilb}_5 = -\left(\frac{w}{5}\right)\left(\frac{q_1}{5}\right), \quad \text{Hilb}_{q_1} = \left(\frac{5}{q_1}\right),$$

and both equal $+1$ for some admissible q_1 if and only if $\left(\frac{w}{5}\right) = -1$. Since the L11 route requires a root of $1 + 4r^2 \equiv 0 \pmod{w}$, i.e. $w \equiv 1 \pmod{4}$, the canonical protocol is obstructed exactly for

$$w \equiv 3 \pmod{4} \quad \text{and} \quad \left(\frac{w}{5}\right) = +1 \quad \Longleftrightarrow \quad w \equiv 11, 19 \pmod{20}.$$

The criterion was audited on all 53 primes $w \in [101, 400]$ with both character branches of q_1 covered: 26 cells have the L11 route, 13 the canonical escape route, 14 are obstructed, and there were zero identity or classification mismatches.

The wall is a statement about the canonical parameter, not about the cells. In fact it is always breakable on the class side.

Theorem 10.13 (Every canonical 5-wall is breakable; PROVED). *Let $w \equiv 11, 19 \pmod{20}$ be prime and set $z = -w$, $f = w$. There is an odd integer a such that the $\tau = 0$ fixed-factor family has an aligned class for infinitely many positive primes q_1 . This theorem concerns class existence; it does not assert an emergent-free member. If a is fixed to 7, the class-side break exists exactly when $\left(\frac{w}{197}\right) = -1$.*

Proof. Because $w \equiv 3 \pmod{4}$, the character sum $\sum_{r \bmod w} \left(\frac{1+4r^2}{w}\right) = -1$ has no zero terms. Exactly $(w+1)/2$ residues therefore give character -1 ; choose one for $a \bmod w$. By CRT also impose $a \equiv 0 \pmod{\text{rad}(w^3+2)}$ and choose an odd lift. For $A = 1 + 4a^2$, $Z = -w^3$, and $D = 1 - Z - a^2Z^2$, the identity

$$4D \equiv (Z-2)^2 \pmod{p} \quad (p \mid A)$$

then gives $\gcd(A, D) = 1$ while preserving $\left(\frac{A}{w}\right) = -1$.

At every prime occurring to odd order in A , prescribe $\left(\frac{-2\varepsilon w q_1}{p}\right) = +1$; impose the exact 2-adic class $\varepsilon w q_1 \equiv a \pmod{4}$; and make $-2A\varepsilon w q_1$ a square at each of 3, 5, 7 not dividing A . These independent local conditions define a reduced CRT class, so Dirichlet gives infinitely many primes q_1 in it. After finitely many exclusions and a real-sign threshold, every frozen, moving, fixed-factor, and infinite symbol is $+1$, and the exponent lemma freezes the resulting progression. The factor-by-factor identity

$$\prod_{p \mid A} (x, d)_p \left(\frac{A}{q_1}\right) = \left(\frac{-2\varepsilon}{A}\right) \left(\frac{w}{A}\right)$$

shows that for odd a the break criterion is $\left(\frac{w}{A}\right) = -1$. Putting $a = 7$ gives $A = 197$ and the stated fixed-parameter criterion. $\square$

Proposition 10.14 (Finite member closures; PROVED per row). *The non-canonical $\tau = 0$ family closes the obstructed cells $w = 131, 139, 151$ at $a = 7$, $q_1 = Q = 41$, $k = 0$. It also closes $[179, [-1, 1]]$ at $a = 7$, $\varepsilon = -1$, $f = 179$, $q_1 = Q = 251$, $k = 0$, with prime rung, tied status exact, and empty ramified set.*

These member rows are stronger than the class theorem only at their four individual cells. Their fixed factor $f = w$ divides $\text{num}(z)$, so they pass through the controlled non-coprimality door excluded by the generalized coprime wall. The kernel verdict uses the same tied-status, ramification, alignment, and complete cofactor ladder as the canonical rows; refusals in the surrounding scans are not evidence.

Remark 10.15 (Composite w). Composite w is not admissible in the architecture: w indexes a place, and the residue-field construction has no meaning for composite modulus — the break occurs already at the field constructor. The downstream arithmetic does generalize once w is decomposed into genuine prime places; as an illustration the pseudo-cell $[21, [-1, 1]]$ closes with the escape route at the honest prime $f = 3$ (member $k = 0$, $Q = 11$, factorint rung, tied and ramified checks exact). This is a clarification of scope, not an extension of the horizon.

11. CONCLUSION AND OUTLOOK

For every nonzero rational Z , the fixed canonical branch $\tau^\dagger = (1+2a^2)/(1+4a^2)$, $\delta_{\tau^\dagger} = -4a^4/(1+4a^2)$ is vertically irreducible. The endpoint–Newton proof is uniform in the fixed fiber, including the exact $Z = 1$ certificate; quantitative Hilbert irreducibility then chooses a good a inside the character progression already required by the class construction.

The logical chain is therefore complete up to one named conjecture: fixed- Z irreducibility gives Hilbert selection in the admissible a progression; the L20 class certificate has $f = w$ and all Schinzel-pair conditions; the L19 fixed-pair implication turns classical Schinzel H into infinitely many emergent-free members; and L6 yields

$$\mathbb{Z} \text{ universally definable in } \mathbb{Q} \text{ with six quantifiers,} \quad \text{efd}_{\mathbb{Q}}(\mathbb{Q} \setminus \mathbb{Z}) \leq 5.$$

The independent reciprocal proof specializes to

$$(a, A, s, \tau^\dagger, \delta) = (1, 5, 0, 3/5, -4/5),$$

factors through a trace quartic, and closes the reciprocal lift by a complete 2-isogeny descent. Its $a = 1$ is only a specialization witness for the rational-function fiber; it need not satisfy the target character condition and is not the class parameter chosen by Hilbert irreducibility.

The alternate routes retain their honest scope without weakening the primary theorem. Infinity alone permits factor degrees 2, 4, and 6; factor tuples control only the product of outside signs without an all-plus compatibility theorem; and a free λ moves τ through an infinite L11c family and would add a seventh witness. None is used as the six-count closure.

L23 sharpens the unconditional frontier but does not cross it. Its half-dimensional beta sieve proves a positive lower bound $\gg X/(\log X)^{3/2}$ for candidates clean at every bad prime below $X^{49/100}$. It proves no positive lower bound for fully clean members: the degree-eight cofactor can retain up to 16 large factors and parity only forces the bad count to be even. The first missing estimate is the two-large-bad-divisor sector beyond Bombieri–Vinogradov. Known prime-argument almost-prime theorems, even Kao’s P_{12} , do not reach the required $R_{\text{bad}} \leq 1$.

The same audit closes several algebraic escape routes. On the refined class, the conic bundle has non-split rank 10 and $\text{Br}(X)/\text{Br}(\mathbb{Q}) \cong \mathbb{Z}/2\mathbb{Z}$, generated by (A, b) ; the rank- ≤ 2 fibration theorem does not apply. Exact norm matching and the loci $P = \pm 2b$ times a square miss Φ , no polynomial-in- b norm section exists, and fixed twists only move the character obligation. The factor branch $L = -A\rho^2$ leads to scoped Hilbert symbols and genus-four curves, not a global section.

L24 decisively refutes the unscaled diagonal five-count. Both exact finite-flat degree-eight unit orientations have empty image on $\Phi(\mathbb{Q}_2)$: orientation I has root valuations 0 or 4, incompatible with the two required squares, while orientation II reduces to a complete contradiction modulo 16. Odd target places are locally soluble in the standard $k = 1$ stratum, so this is a genuinely dyadic obstruction. The 4,026,282 zero-hit search is corroborating EVIDENCE only, not the proof.

L25 separates that fact from a universal coupling wall. Its two scaled type-I formulas cover complementary dyadic parities, while their smooth target formula is on a freely chosen unit- b stratum rather than the aligned L20 stratum. L26 changes the tie: $Z = z^2$ and $\eta = Z(b+1)/(Db)$ make the tied polynomial reciprocal with a quartic trace character and remove its dyadic symbol. L28 proves uniformly that this trace quartic is irreducible over $\mathbb{Q}_2$, and that

$$N(2(\theta+2)) \in A\mathbb{Q}_2^{\times 2}, \quad A \equiv 5 \pmod{8}.$$

Thus the bad-sign extension is always nontrivial. L29 further proves

$$\theta^2 - 4 \in \mathbb{Q}_2(\theta)^{\times 2} \iff v_2(Z) = -2 \text{ or } v_2(Z) \geq 2$$

on the square- Z family, and the pullback $Z = 8z^2/(1+z^2)$ preserves odd targets while forcing the split stratum. L27 instead uses the single shear

$$(y, r) = (2X + 28\rho, sX + \rho).$$

It covers all dyadic parities and all aligned odd targets. L30 improves the algebraic route by fixing

$$(y, r) = (2, sX + \rho).$$

The resulting equation is quadratic in $m = \lambda^2$, hence even quartic in λ . A normalized Hensel argument solves every dyadic stratum; a two-quadratic character count supplies fibre-regular rows for every $w \geq 5$; and $(a, b, z) = (5, 3, 3)$ is an exact strong-Hensel $w = 3$ fibre. L30 also rules out a generic norm section for every rational square-branch tie and every B -independent linear-shear cancellation.

L31 proves the exact fixed-field member

$$(w, a, Z, q_0, b, \rho, \lambda) = (13, 3, 169, 1, 13, 14/13, 12).$$

Its norm cofactor is prime by a recursive Pocklington certificate and every Hilbert symbol is $+1$. On the quartic side, L31 gives an exact rational point at

$$(a, c, \lambda, b, Z) = (1, -64/25, 3, -3253/3125, 2033125/6101423),$$

but Z is not a cube and its shared target prime leaves c a unit. A complete external Magma calculation [13] proves that the two genus-2 cube curves on this unique constant- c section have only $z = 0$. Thus this section is closed and the controlled quartic point remains open on other sections.

Accordingly, classical Schinzel's Hypothesis H remains the sole conjectural input to the established six-universal-quantifier definition. The exact next frontiers are a uniform reciprocal member theorem, a cube-compatible controlled point on L30's quartic, or AP1 via a pointwise fixed-family Buchstab/Hilbert-detector asymptotic. The theorem remains conditional: we make no new unconditional quantifier-record claim and do not claim that H10($\mathbb{Q}$) is solved.

REPRODUCIBILITY

All checked-in producer certificates in this paper use a stdlib-only kernel with a proven-primality discipline: deterministic Miller–Rabin below 3.3×10^{24} , Pocklington with the Brillhart–Lehmer–Selfridge relaxation above, and explicit refusal labels that never count as evidence. The sole external-CAS result is the scoped L31 genus-2 calculation; its exact Magma V2.29-9 request, complete output, and intrinsic-help semantics are recorded, while the absence of a standalone checked-in Magma certificate is explicit. The companion paper documents the engine, artifact schemas, exact reproduction commands, and precise regeneration scope.

REFERENCES

- [1] P. T. Bateman and R. A. Horn, *A heuristic asymptotic formula concerning the distribution of prime numbers*, Mathematics of Computation **16** (1962), 363–367.
- [2] J.-L. Colliot-Thélène and J.-J. Sansuc, *Sur le principe de Hasse et l’approximation faible, et sur une hypothèse de Schinzel*, Acta Arithmetica **41** (1982), no. 1, 33–53.
- [3] J.-L. Colliot-Thélène, A. N. Skorobogatov and Sir P. Swinnerton-Dyer, *Rational points and zero-cycles on fibred varieties: Schinzel’s hypothesis and Salberger’s device*, Journal für die reine und angewandte Mathematik **495** (1998), 1–28.
- [4] G. Cornelissen and K. Zahidi, *Topology of Diophantine sets: remarks on Mazur’s conjectures*, in: Hilbert’s Tenth Problem: Relations with Arithmetic and Algebraic Geometry (Ghent, 1999), Contemporary Mathematics **270**, American Mathematical Society, Providence RI, 2000, pp. 253–260; arXiv:math/0006140.
- [5] N. Daans, *Universally defining $\mathbb{Z}$ in $\mathbb{Q}$ with 10 quantifiers*, Journal of the London Mathematical Society (2) **109** (2024), no. 2, e12864, 20 pp.; doi:10.1112/jlms.12864; arXiv:2301.02107.
- [6] N. Daans, P. Dittmann and A. Fehm, *Existential rank and essential dimension of diophantine sets*, preprint, arXiv:2102.06941v5 (2021–2024).
- [7] Y. Harpaz, A. N. Skorobogatov and O. Wittenberg, *The Hardy–Littlewood conjecture and rational points*, Compositio Mathematica **150** (2014), no. 12, 2095–2111; doi:10.1112/S0010437X14007568; arXiv:1304.3333.
- [8] H. A. Helfgott, *Square-free values of $f(p)$, f cubic*, Acta Mathematica **213** (2014), no. 1, 107–135; doi:10.1007/s11511-014-0117-2; arXiv:1112.3820.
- [9] S. D. Cohen, *The distribution of Galois groups and Hilbert’s irreducibility theorem*, Proceedings of the London Mathematical Society (3) **43** (1981), no. 2, 227–250; doi:10.1112/plms/s3-43.2.227.
- [10] J. Koenigsmann, *Defining $\mathbb{Z}$ in $\mathbb{Q}$* , Annals of Mathematics (2) **183** (2016), no. 1, 73–93; doi:10.4007/annals.2016.183.1.2; arXiv:1011.3424.
- [11] D. Krumm, *Squarefree parts of polynomial values*, Journal de Théorie des Nombres de Bordeaux **28** (2016), no. 3, 699–724; doi:10.5802/jtnb.959; arXiv:1407.4890.
- [12] T. Reuss, *Power-free values of polynomials*, Bulletin of the London Mathematical Society **47** (2015), no. 2, 270–284; doi:10.1112/blms/bdv007; arXiv:1307.2802.
- [13] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system I: The user language*, Journal of Symbolic Computation **24** (1997), nos. 3–4, 235–265; computation replayed with Magma V2.29-9.
- [14] A. Schinzel and W. Sierpiński, *Sur certaines hypothèses concernant les nombres premiers*, Acta Arithmetica **4** (1958), 185–208; erratum, ibid. **5** (1958), 259.
- [15] J.-P. Serre, *Lectures on the Mordell–Weil Theorem*, 3rd ed., Aspects of Mathematics **E15**, Friedr. Vieweg & Sohn, Braunschweig, 1997, x+218 pp.; doi:10.1007/978-3-663-10632-6.
- [16] Z.-W. Sun, *$\mathbb{Q} \setminus \mathbb{Z}$ is diophantine over $\mathbb{Q}$ with 7 unknowns*, preprint, arXiv:2607.28606 (unrefereed).