



量子少体模型在核反应研究中的挑战与前沿进展

金磊
同济大学

- ◆ STARS
- ◆ Three body Forces in Reaction Model
- ◆ Trojan Horse Mechanism
- ◆ Quantum Four Body Model

Shanghai Tongji Advanced Reaction Solver (STARS)

3

For a two-body structured projectile $a = b + x$ interacting with a target A , in the three-body model space the scattering wave function can be solved by

$$(E - H)\Psi = 0, \text{ with } H = H_{bx} + T_{aA} + U_b + U_x$$

For a two-body structured projectile $a = b + x$ interacting with a target A , in the three-body model space the scattering wave function can be solved by

$$(E - H)\Psi = 0, \text{ with } H = H_{bx} + T_{aA} + U_b + U_x$$

The unit operator in this three body model space is given by

$$1 = \sum_{\alpha n_0} \int R^2 dR |\phi_{bx}^{n_0} R \alpha\rangle \langle \phi_{bx}^{n_0} R \alpha| + (2\pi)^{-3} \sum_{\alpha} \int R^2 dR \int d\vec{k} |\phi_{bx}^{\vec{k}(+)} R \alpha\rangle \langle \phi_{bx}^{\vec{k}(+)} R \alpha| \xrightarrow[\text{CD}]{\text{continuum-discretized}} 1 \approx \sum_{\alpha n} \int R^2 dR |\phi_{bx}^n R \alpha\rangle \langle \phi_{bx}^n R \alpha|$$

For a two-body structured projectile $a = b + x$ interacting with a target A , in the three-body model space the scattering wave function can be solved by

$$(E - H)\Psi = 0, \text{ with } H = H_{bx} + T_{aA} + U_b + U_x$$

The unit operator in this three body model space is given by

$$1 = \sum_{\alpha n_0} \int R^2 dR |\phi_{bx}^{n_0} R \alpha\rangle \langle \phi_{bx}^{n_0} R \alpha| + (2\pi)^{-3} \sum_{\alpha} \int R^2 dR \int d\vec{k} |\phi_{bx}^{\vec{k}(+)} R \alpha\rangle \langle \phi_{bx}^{\vec{k}(+)} R \alpha| \xrightarrow[\text{CD}]{\text{continuum-discretized}} 1 \approx \sum_{\alpha n} \int R^2 dR |\phi_{bx}^n R \alpha\rangle \langle \phi_{bx}^n R \alpha|$$

Insert the unit operator, and then the Schrödinger equation becomes:

$$(E - \varepsilon_n - T_{aA}(R\alpha)) \langle \phi_{bx}^n R \alpha | \Psi^{j_{\text{in}}(+)} \rangle - \sum_{\tilde{\alpha}'} U_{\tilde{\alpha}, \tilde{\alpha}'}(R) \langle \phi_{bx}^{n'} R \alpha' | \Psi^{j_{\text{in}}(+)} \rangle = 0, \text{ coupled-channels equations}$$

For a two-body structured projectile $a = b + x$ interacting with a target A , in the three-body model space the scattering wave function can be solved by

$$(E - H)\Psi = 0, \text{ with } H = H_{bx} + T_{aA} + U_b + U_x$$

The unit operator in this three body model space is given by

$$1 = \sum_{\alpha n_0} \int R^2 dR |\phi_{bx}^{n_0} R \alpha\rangle \langle \phi_{bx}^{n_0} R \alpha| + (2\pi)^{-3} \sum_{\alpha} \int R^2 dR \int d\vec{k} |\phi_{bx}^{\vec{k}(+)} R \alpha\rangle \langle \phi_{bx}^{\vec{k}(+)} R \alpha| \xrightarrow[\text{CD}]{\text{continuum-discretized}} 1 \approx \sum_{\alpha n} \int R^2 dR |\phi_{bx}^n R \alpha\rangle \langle \phi_{bx}^n R \alpha|$$

Insert the unit operator, and then the Schrödinger equation becomes:

$$(E - \varepsilon_n - T_{aA}(R\alpha)) \langle \phi_{bx}^n R \alpha | \Psi^{j_{\text{in}}(+)} \rangle - \sum_{\tilde{\alpha}'} U_{\tilde{\alpha}, \tilde{\alpha}'}(R) \langle \phi_{bx}^{n'} R \alpha' | \Psi^{j_{\text{in}}(+)} \rangle = 0, \text{ coupled-channels equations}$$

Using the R-matrix method transforms the differential equations into linear equations.

For a two-body structured projectile $a = b + x$ interacting with a target A , in the three-body model space the scattering wave function can be solved by

$$(E - H)\Psi = 0, \text{ with } H = H_{bx} + T_{aA} + U_b + U_x$$

The unit operator in this three body model space is given by

$$1 = \sum_{\alpha n_0} \int R^2 dR |\phi_{bx}^{n_0} R \alpha\rangle \langle \phi_{bx}^{n_0} R \alpha| + (2\pi)^{-3} \sum_{\alpha} \int R^2 dR \int d\vec{k} |\phi_{bx}^{\vec{k}(+)} R \alpha\rangle \langle \phi_{bx}^{\vec{k}(+)} R \alpha| \xrightarrow[\text{CD}]{\text{continuum-discretized}} 1 \approx \sum_{\alpha n} \int R^2 dR |\phi_{bx}^n R \alpha\rangle \langle \phi_{bx}^n R \alpha|$$

Insert the unit operator, and then the Schrödinger equation becomes:

$$(E - \varepsilon_n - T_{aA}(R\alpha)) \langle \phi_{bx}^n R \alpha | \Psi^{j_{\text{in}}(+)} \rangle - \sum_{\tilde{\alpha}'} U_{\tilde{\alpha}, \tilde{\alpha}'}(R) \langle \phi_{bx}^{n'} R \alpha' | \Psi^{j_{\text{in}}(+)} \rangle = 0, \text{ coupled-channels equations}$$

Using the R-matrix method transforms the differential equations into linear equations.

STARS uses modern techniques such as emulator, Bayesian Analysis, and machine learning and to solve the linear equations of the quantum three-body problem.

Three body Forces in Reaction Model

A nuclear reaction is a complicated many-body problem. However, for some reactions, it can be solved using a three-body model space.

$$H^{\text{many}} \xrightarrow{\text{projection}} H^3 \approx H^{\text{model}}$$

Three body Forces in Reaction Model

A nuclear reaction is a complicated many-body problem. However, for some reactions, it can be solved using a three-body model space.

$$H^{\text{many}} \xrightarrow{\text{projection}} H^3 \approx H^{\text{model}}$$

Take $d + A$ reaction as an example

$$H^{\text{many}} = H_0 + V_{np} + U_{nA} + U_{pA} + H_A + \Delta v_{nA} + \Delta v_{pA}$$

Three body Forces in Reaction Model

A nuclear reaction is a complicated many-body problem. However, for some reactions, it can be solved using a three-body model space.

$$H^{\text{many}} \xrightarrow{\text{projection}} H^3 \approx H^{\text{model}}$$

Take $d + A$ reaction as an example

$$H^{\text{many}} = H_0 + V_{np} + U_{nA} + U_{pA} + H_A + \Delta v_{nA} + \Delta v_{pA}$$

By projection, keep the three-body observables unchanged

$$H^3 = H_0 + V_{np} + U_{nA} + U_{pA} + U_{npA}$$

Three body Forces in Reaction Model

A nuclear reaction is a complicated many-body problem. However, for some reactions, it can be solved using a three-body model space.

$$H^{\text{many}} \xrightarrow{\text{projection}} H^3 \approx H^{\text{model}}$$

Take $d + A$ reaction as an example

$$H^{\text{many}} = H_0 + V_{np} + U_{nA} + U_{pA} + H_A + \Delta v_{nA} + \Delta v_{pA}$$

By projection, keep the three-body observables unchanged

$$H^3 = H_0 + V_{np} + U_{nA} + U_{pA} + U_{npA}$$

Take the approximation:

$$H^{\text{model}} = H_0 + V_{np} + U_{nA} + U_{pA}$$

Three body Forces in Reaction Model

A nuclear reaction is a complicated many-body problem. However, for some reactions, it can be solved using a three-body model space.

$$H^{\text{many}} \xrightarrow{\text{projection}} H^3 \approx H^{\text{model}}$$

Take $d + A$ reaction as an example

$$H^{\text{many}} = H_0 + V_{np} + U_{nA} + U_{pA} + H_A + \Delta v_{nA} + \Delta v_{pA}$$

By projection, keep the three-body observables unchanged

$$H^3 = H_0 + V_{np} + U_{nA} + U_{pA} + U_{npA}$$

Take the approximation:

$$H^{\text{model}} = H_0 + V_{np} + U_{nA} + U_{pA}$$

$$U_{npA} = \langle \phi_A | \left(\Delta v_{nA} + \Delta v_{pA} \right) + \left(\Delta v_{nA} + \Delta v_{pA} \right) Q_A \frac{1}{E + i\epsilon - Q_A H Q_A} Q_A \left(\Delta v_{nA} + \Delta v_{pA} \right) | \phi_A \rangle$$

In weakly bound systems, three-body forces are minimal;
in tightly bound systems, they are significant.

Three body Forces in Reaction Model

A nuclear reaction is a complicated many-body problem. However, for some reactions, it can be solved using a three-body model space.

$$H^{\text{many}} \xrightarrow{\text{projection}} H^3 \approx H^{\text{model}}$$

Take $d + A$ reaction as an example

$$H^{\text{many}} = H_0 + V_{np} + U_{nA} + U_{pA} + H_A + \Delta v_{nA} + \Delta v_{pA}$$

By projection, keep the three-body observables unchanged


$$H^3 = H_0 + V_{np} + U_{nA} + U_{pA} + U_{npA}$$

Take the approximation:


$$H^{\text{model}} = H_0 + V_{np} + U_{nA} + U_{pA}$$

$$U_{npA} = \langle \phi_A | \left(\Delta v_{nA} + \Delta v_{pA} \right) + \left(\Delta v_{nA} + \Delta v_{pA} \right) Q_A \frac{1}{E + i\epsilon - Q_A H Q_A} Q_A \left(\Delta v_{nA} + \Delta v_{pA} \right) | \phi_A \rangle$$


In weakly bound systems, three-body forces are minimal;
in tightly bound systems, they are significant.



Trojan Horse Mechanism




Trojan Horse Mechanism



For the reaction of $a(=b+x) + A \rightarrow b + (x+A)^*$


Trojan Horse Mechanism



For the reaction of $a(=b+x) + A \rightarrow b + (x+A)^*$

In energy conservation,
 $E_{bx} + E_{aA} = E_b + E_{xA}$. If E_{aA}
 overcomes the Coulomb
 barrier, E_b takes most of the
 energy, making E_{xA} relatively
 small.


Trojan Horse Mechanism




The Trojan Horse mechanism may offer a potential way to synthesize neutron-rich superheavy isotopes.

For the reaction of $a(=b+x) + A \rightarrow b + (x+A)^*$

In energy conservation, $E_{bx} + E_{aA} = E_b + E_{xA}$. If E_{aA} overcomes the Coulomb barrier, E_b takes most of the energy, making E_{xA} relatively small.





Trojan Horse Mechanism



The Trojan Horse mechanism may offer a potential way to synthesize neutron-rich superheavy isotopes.


For the reaction of $a(=b+x)+A\rightarrow b+(x+A)^*$

In energy conservation, $E_{bx} + E_{aA} = E_b + E_{xA}$. If E_{aA} overcomes the Coulomb barrier, E_b takes most of the energy, making E_{xA} relatively small.



Quantum Four Body Model


Study the reaction mechanism of three-body structured projectiles, such as ${}^9\text{C} = {}^7\text{Be} + p + p$ induced reactions



Quantum Four Body Model

Study the reaction mechanism of three-body structured projectiles, such as ${}^9\text{C} = {}^7\text{Be} + p + p$ induced reactions


Even in the simplest DWBA form, it requires the three-body bound state wave function in coordinate space



Quantum Four Body Model

Study the reaction mechanism of three-body structured projectiles, such as ${}^9\text{C} = {}^7\text{Be} + p + p$ induced reactions

Even in the simplest DWBA form, it requires the three-body bound state wave function in coordinate space



Need to solve Faddeev–Merkuriev Equations

$$\left(E - H_0 - V_{ij} - W_k\right) \psi_{ij} = V_{ij}^S \left(\psi_{jk} + \psi_{ki}\right),$$


$$\left(E - H_0 - V_{jk} - W_i\right) \psi_{jk} = V_{jk}^S \left(\psi_{ij} + \psi_{ki}\right),$$

$$\left(E - H_0 - V_{ki} - W_j\right) \psi_{ki} = V_{ki}^S \left(\psi_{ij} + \psi_{jk}\right),$$

Quantum Four Body Model

Study the reaction mechanism of three-body structured projectiles, such as ${}^9\text{C} = {}^7\text{Be} + p + p$ induced reactions

Even in the simplest DWBA form, it requires the three-body bound state wave function in coordinate space



Need to solve Faddeev–Merkuriev Equations

$$\left(E - H_0 - V_{ij} - W_k\right) \psi_{ij} = V_{ij}^S \left(\psi_{jk} + \psi_{ki}\right),$$

$$\left(E - H_0 - V_{jk} - W_i\right) \psi_{jk} = V_{jk}^S \left(\psi_{ij} + \psi_{ki}\right),$$

$$\left(E - H_0 - V_{ki} - W_j\right) \psi_{ki} = V_{ki}^S \left(\psi_{ij} + \psi_{jk}\right),$$

Using the complex scaling technique, one can treat the three-body **bound state**, **scattering states**, and **resonance states** simultaneously