

# Notes of "Group"

Jinxin Wang

## 1 Overview

- Group and subgroup
  - Def: A group
    - \* Rmk: The uniqueness of the identity element in a group
    - \* Rmk: The uniqueness of the inverse element of an element in a group
    - \* Rmk: The definition of a group does not specify the uniqueness of the identity element and the inverse element of each element
    - \* Rmk: The default notation of an abstract group is with multiplication, such as the operator and the identity element
  - Def: A subgroup of a group
    - \* Rmk: Trivial subgroups and proper subgroups
    - \* Rmk: Is the identity element of a subgroup always the same as the one of its parent group?
  - Examples of groups and subgroups
    - \* Eg:  $(\mathbb{Q}, +)$  and  $(\mathbb{Z}, +)$
    - \* Eg:  $(\mathbb{R}/\{0\}, \cdot)$  and  $(\mathbb{R}^+, \cdot)$
    - \* Eg:  $S_n$  and the set of even permutations of order  $n$
    - \* Eg:  $GL_n(\mathbb{R})$  and  $SL_n(\mathbb{R})$
    - \* Eg: The set of invertible elements in a monoid
    - \* Eg:  $(\{1, -1\}, \cdot)$
  - Def: An abelian group
  - Examples of abelian groups
    - \* Eg:  $(\mathbb{Z}/n\mathbb{Z}, +)$
    - \* Eg:  $A(X, Y) = \{f : X \rightarrow Y\}$  where  $Y$  is an abelian group
    - \* Eg:  $L(X, Y) = \{f : X \rightarrow Y \mid f \text{ is an additive map}\}$
  - Def: The order (cardinality) of a group, a finite group and an infinite group
  - Examples of finite groups and infinite groups
    - \* Examples of finite groups:  $S_n, (\{1, -1\}, \cdot)$
    - \* Examples of infinite groups:  $(\mathbb{Q}, +)$

- Cyclic groups
  - Rmk: An element in a group generates a subgroup of the group
  - Def: A cyclic group and its generator(s)
    - \* Rmk: The uniqueness of the generator(s) of a cyclic group
  - Examples of cyclic groups
    - \* Eg:  $(\mathbb{Z}, +)$  can be generated by 1 or  $-1$
    - \* Eg:  $(\{1, -1\}, \cdot)$  can be generated by  $-1$
  - Rmk: The order of a generated cyclic group by an element in a finite group
- The order of an element in a group
  - Def: An element of infinite order or finite order in a group
  - Examples of elements of infinite order and elements of finite order in groups
    - \* Examples of elements of finite order in groups: A permutation in  $S_n$
    - \* Examples of elements of infinite order in groups: 1 and  $-1$  in  $(\mathbb{Z}, +)$
  - Prop: The relationship between the order of an element in a group and the order of the cyclic group generated by it
- Subgroups of a cyclic group
  - Prop: The form of subgroups of a cyclic group
    - \* Rmk: For a subgroup of a finite cyclic group, the factor  $k$  is not unique?
  - Prop: The relationship between different generators of a finite cyclic group
  - Eg: An application of the form of subgroups of a cyclic group to  $(\mathbb{Z}, +)$
- Homomorphisms and isomorphisms
  - Def: A group homomorphism
  - Def: A group isomorphism
    - \* Rmk: Both a homomorphism and an isomorphism refer to a mapping rather than a relation between two algebraic structures
  - Prop: Some basic properties of a group homomorphism
  - Prop: Some basic properties of a group isomorphism
- Examples and conclusions of group homomorphisms and group isomorphisms
  - Prop: A necessary and sufficient condition of two cyclic groups to be isomorphic in terms of the orders of them

## 2 Group and subgroup

**Definition 1** (A group). *A set  $G$  is called a group if a binary operation  $\cdot$  is defined on it, and for any  $a, b, c \in G$ , it holds that*

**D1** *Associative law:  $a \cdot (b \cdot c) = (a \cdot b) \cdot c$*

**D2** *Identity element: There exists  $e \in G$  such that  $e \cdot a = a \cdot e = a$  for each  $a \in G$*

**D3** *Inverse element: For each  $a \in G$ , there exists  $b \in G$  such that  $ab = ba = e$*

**Remark 1** (The uniqueness of the identity element in a group). *Suppose there are  $e \in G$  and  $e' \in G$  which satisfy the definition of the identity element, then*

$$e = e \cdot e' = e'$$

**Remark 2** (The uniqueness of the inverse element of an element in a group). *For each  $a \in G$ , suppose there are two inverse elements  $b \in G$  and  $b' \in G$ , then*

$$b = b \cdot e = b \cdot (a \cdot b') = (b \cdot a) \cdot b' = e \cdot b' = b'$$

**Remark 3** (The definition of a group does not specify the uniqueness of the identity element and the inverse element of each element). *As we can see, the definition of a group does not require the uniqueness of the identity element and the inverse element of each element in a group. The reason is that the uniqueness is a natural property of the identity element and the inverse element of each element once a set satisfies the definition of a group. Hence, we don't need and aren't supposed to add such conditions to the definition.*

**Remark 4** (The default notation of an abstract group is with multiplication, such as the operator and the identity element).

**Definition 2** (A subgroup of a group).

**Remark 5** (Is the identity element of a subgroup always the same as the one of its parent group?). *The identity element of a group is also the identity of its every subgroup.*

*Proof:* Suppose  $U$  is a group,  $V \subset U$  is a subgroup of  $U$ , and  $e_U$  and  $e_V$  are the identity elements of them respectively. For  $e_V$ , we have  $e_V^2 = e_V$ . Since  $e_V \in U$ , the equation also holds in  $U$ . Then we have  $e_V^{-1}e_V^2 = e_V^{-1}e_V$ , which is  $e_V = e_U$ .

*Recall that the similar conclusion doesn't hold for monoids. We can see the changes brought by the additional axioms of a group compared with a monoid.*

**Example 1** ( $(\mathbb{Q}, +)$  and  $(\mathbb{Z}, +)$ ).  *$(\mathbb{Q}, +)$  is a group, and  $(\mathbb{Z}, +)$  is a subgroup of it.*

**Example 2** ( $(\mathbb{R}/\{0\}, \cdot)$  and  $(\mathbb{R}^+, \cdot)$ ).  *$(\mathbb{R}/\{0\}, \cdot)$  is a group, and  $(\mathbb{R}^+, \cdot)$  is a subgroup of it.*

**Example 3** ( $S_n$  and the set of even permutations of order  $n$ ). *The set of permutations of order  $n$   $S_n$  is a group, and the set of even permutations of order  $n$  is a subgroup of  $S_n$ .*

**Example 4** ( $GL_n(\mathbb{R})$  and  $SL_n(\mathbb{R})$ ). *The set of invertible real-valued matrices of order  $n$ , denoted by  $GL_n(\mathbb{R})$ , forms a group. The set of invertible real-valued matrices of order  $n$  whose determinant is 1, denoted by  $SL_n(\mathbb{R})$ , is a subgroup of  $GL_n(\mathbb{R})$ .*

### 3 Cyclic groups

#### 4 The order of an element in a group

**Definition 3** (An element of infinite order or finite order in a group). *Given an element  $a$  in a group, we check the sequence of its integer exponentiations. If all of them are different, we say that the order of the element  $a$  is infinite, and  $a$  is called an element of infinite order in the group. If there are same elements, then there exists integers  $k_i$  such that  $a^{k_i} = e$ . The minimal integer in the set  $\{k_i\}$ , denoted by  $q$ , is called the order of the element  $a$  in the group, and  $a$  is called an element of finite order in the group, or an element of order  $q$  in the group.*

#### 5 Subgroups of a cyclic group

#### 6 Homomorphisms and isomorphisms

#### 7 Examples and conclusions of group homomorphisms and group isomorphisms