

Lecture 9: Max-Min and Least Squares

1 Application of Partial Derivatives: Optimization Problem

1.1 Max-Min

Optimization problems refer to the tasks of finding the maximum or minimum point of a function. Here we focus on finding the maximum or minimum point of multivariable functions, such as $f(x, y)$.

Another concept is local max/min point, which means the function value of this point is greater/less than the function values of adjacent points, but it is not necessarily the greatest or least function value in the function domain.

Theorem. *If a point in a function f is a local max/min point, then all of the partial derivatives at this point are equal to 0.*

Proof:

We will prove it by contradiction. Suppose there is a local max/min point in a function $f(x_1, x_2, \dots, x_n)$, and one of its partial derivative $\frac{\partial f}{\partial x_i}$ is not 0. According to the definition of partial derivatives, in that direction the function values of the two adjacent point are

$$\begin{aligned} f(x_1, x_2, \dots, x_i + \Delta x_i, \dots, x_n) &= f(x_1, x_2, \dots, x_n) + \frac{\partial f}{\partial x_i} \Delta x_i \\ f(x_1, x_2, \dots, x_i - \Delta x_i, \dots, x_n) &= f(x_1, x_2, \dots, x_n) - \frac{\partial f}{\partial x_i} \Delta x_i \end{aligned}$$

Therefore, one of them is less than the local max/min point and the other is greater than the local max/min point. Hence, it is not a local max/min point. There is a contradiction.

Therefore, it is proved that if a point in a function f is a local max/min point, then all of the partial derivatives at this point are equal to 0.

Therefore, A point is a local max/min point

\Rightarrow All partial derivatives at that point are 0

\iff The tangent plane at that point is horizontal

\iff The point is a critical point.

1.2 Critical Points

Definition. A point is a critical point of a function f if and only if all the partial derivatives on this point are equal to 0.

For a multivariable function with two independent variables $f(x, y)$, a point (x_0, y_0) is a critical point if $\frac{\partial f}{\partial x}(x_0, y_0) = 0$ and $\frac{\partial f}{\partial y}(x_0, y_0) = 0$.

Example 1. Find all the critical points of the function $f(x, y) = x^2 - 2xy + 3y^2 + 2x - 2y$.

Solution:

$$\begin{aligned}\frac{\partial f}{\partial x} &= 2x - 2y + 2 \\ \frac{\partial f}{\partial y} &= -2x + 6y - 2\end{aligned}$$

For critical points, they should satisfy

$$\begin{cases} \frac{\partial f}{\partial x} = 2x - 2y + 2 = 0 \\ \frac{\partial f}{\partial y} = -2x + 6y - 2 = 0 \end{cases}$$

Solving this linear system, the result is $(-1, 0)$. Therefore, there are only one critical point for this function, which is $(-1, 0)$.

1.3 Types of Critical Points

A critical point of a function f can be

- local maximum point
- local minimum point
- saddle point

Question 1. Is there any other possibilities for a critical point of a multivariable function? If not, how to prove it?

How to determine the type of a critical point?

- Test the second derivative (cover in the next lecture).
- Find the codomain of the function and compare the function value on the critical point with the codomain.

Example 2. Find the type of the critical point $(-1, 0)$ of the function $f(x, y) = x^2 - 2xy + 3y^2 + 2x - 2y$.

Solution:

The function value of the critical point $(-1, 0)$ is

$$\begin{aligned} f(-1, 0) &= (-1)^2 - 2 \times (-1) \times 0 + 3 \times 0^2 + 2 \times (-1) - 2 \times 0 \\ &= 1 - 2 \\ &= -1 \end{aligned}$$

To find the codomain of the function, we can transform its formula

$$\begin{aligned} f(x, y) &= x^2 - 2xy + 3y^2 + 2x - 2y \\ &= (x - y)^2 + 2y^2 + 2x - 2y \\ &= ((x - y)^2 + 2(x - y) + 1) + 2y^2 - 1 \\ &= (x - y + 1)^2 + 2y^2 - 1 \end{aligned}$$

Therefore, it is apparent that the codomain of the function is $[-1, \infty]$. Hence, the critical point $(-1, 0)$ is the minimum point of the function.

2 Example of Optimization Problem: Least Squares