

PS3

Wednesday, 9 December 2020


13:57

Q1, 3, 5, 2, 5, 3, 6, 7

$$1a) \llbracket \log(-1) \rrbracket = \{ \log|-1| + i\theta : \theta \in \llbracket \arg(-1) \rrbracket \}$$

$$= \{ i \frac{3\pi}{2} k : k \in \mathbb{Z} \} \quad \log(-1) = -i \frac{\pi}{2}$$

$$b) \llbracket \log(1-i) \rrbracket = \{ \log|1-i| + i\theta : \theta \in \llbracket \arg(1-i) \rrbracket \}$$

$$= \{ \log\sqrt{2} - i \frac{\pi}{4} k : k \in \mathbb{Z} \} \quad \log(1-i) = \frac{1}{2} \log 2 - i \frac{\pi}{4}$$


$$c) \llbracket \log(e^{-\frac{2\pi i}{3}}) \rrbracket = \{ \log|e^{-\frac{2\pi i}{3}}| + i\theta : \theta \in \llbracket \arg(e^{-\frac{2\pi i}{3}}) \rrbracket \}$$

$$= \{ -i \frac{2\pi}{3} k : k \in \mathbb{Z} \} \quad \log e^{-\frac{2\pi i}{3}} = -i \frac{2\pi}{3}$$

$$3.1) f(z) = \operatorname{Re}(z)$$

$$\gamma(t) = t - it^2 \quad (t \in [0, 1])$$

$$\int_{\gamma} f(z) dz = \int_0^1 \operatorname{Re}(t - it^2) (1 - 2it) dt$$

$$= \int_0^1 (t - 2it^2) dt$$

$$= \left[\frac{1}{2} t^2 - \frac{2}{3} i t^3 \right]_0^1$$

$$= \frac{1}{2} - \frac{2}{3} i$$

$$2) f(z) = \frac{1}{z}$$

γ = unit circle clockwise starting at 1 traversing unit circle 4 times

$$\gamma = e^{it} \quad (t \in [0, -8\pi])$$

$$\int_{\gamma} f(z) dz = \int_0^{-8\pi} \frac{1}{e^{it}} i e^{it} dt = \int_0^{-8\pi} i dt = i t \Big|_0^{-8\pi} = -8\pi i$$

$$3) f(z) = |z|^4$$

γ = straight line from $-1+i$ to $1+i$



$$\gamma(t) = t + i \quad (t \in [-1, 1])$$

$$|t + i|^4 = (\sqrt{t^2 + 1})^4 = (t^2 + 1)^2 = t^4 + 2t^2 + 1$$

$$\int_{\gamma} f(z) dz = \int_{-1}^1 (t^4 + 2t^2 + 1) dt = \left[\frac{1}{5} t^5 + \frac{2}{3} t^3 + t \right]_{-1}^1 = \frac{56}{15}$$

$$5.2) \Gamma_R = \text{half circle in upper half of plane radius } R \text{ centred at the origin traversed CCW.}$$

$$= R e^{it} \quad (t \in [0, \pi])$$

Let $f(z) = z^{-4} e^{iz}$. So $|f(z)| = |z|^{-4}$. On Γ_R we have

Let $f(z) = z^{-4} e^{iz}$. So $|f(z)| = |z|^{-4}$. On Γ_R we have

$$|f(\Gamma_R(t))| = |Re^{it}|^{-4} = R^{-4} |e^{it}|^{-4} = R^{-4}$$

So for $t \in [0, \pi]$, $|f(z)| \leq R$. Further we have that $\text{length } \Gamma_R = \pi R$ so

$$\left| \int_{\Gamma} f(z) dz \right| \leq R^{-4} \pi R = \pi R^{-3}$$

by the estimation lemma.

3) $\gamma(z_0; r)$ = circle radius r centred at z_0 traversed CCW.

$$\gamma(0; R) = Re^{it} \quad (t \in [0, 2\pi])$$

Let $f(z) = \frac{z-1}{z+1}$. So $|f(z)| = \left| \frac{z-1}{z+1} \right|$. On γ we have

$$|f(\gamma(0; R))| = \left| \frac{Re^{it}-1}{Re^{it}+1} \right| = \frac{|Re^{it}-1|}{|Re^{it}+1|}$$


Clearly the maximum will occur when $e^{it} = -1$ so we have

$$|f(z)| \leq \frac{|-1-1-R|}{|-1-1+R|} = \frac{1+R}{1-R}$$

provided $R \neq 1$. Further $\text{length } \gamma(0; R) = 2\pi R$ so by the estimation lemma

$$\left| \int_{\gamma} f(z) dz \right| \leq \frac{1+R}{1-R} 2\pi R = \frac{2\pi R(R+1)}{|R-1|}$$

6a)



$$\gamma_1(t) = t + (1-t)i \quad t \in [0, 1]$$

$$\gamma_2(t) = (1-t) - it \quad t \in [0, 1]$$

$$\gamma_3(t) = it \quad t \in [0, 1]$$

$\int_{\gamma} z dz = 0$ by Cauchy's theorem since γ is closed and z is holomorphic everywhere.

$$\begin{aligned} \text{b) } \int_{\gamma} \bar{z} dz &= \int_0^1 ((1-t) + ti)(1-i) + (1-t)i(-1) + t(i) dt \\ &= \int_0^1 ((1-t)(1-i) + t(1-i) - i + ti + it) dt \\ &= \int_0^1 t(-1 + i + i + 2i) + 1 - i - i dt \\ &= \int_0^1 4it + 1 - 2i dt \\ &= 2it^2 + (1-2i)t \Big|_0^1 \end{aligned}$$

$$\begin{aligned}
 &= \int_0^1 4it + (1-2i) dt \\
 &= 2it^2 + (1-2i)t \Big|_0^1 \\
 &= 1
 \end{aligned}$$

7) Cauchy's theorem states that if f is holomorphic on a simply connected region then the path integral of f is 0 for any closed path in that region. Now γ is the circle of radius $\frac{3}{2}$ centered at the origin. We have:

(i) $\frac{1}{z-1}$ not holomorphic at $z=1 \in \gamma$ so path integral not equal to 0.

(ii) $z^5 e^{4z}$ holomorphic everywhere so path integral equal to 0.

(iii) $\sec^2(z) = \frac{1}{\cos^2(z)}$ not holomorphic if $\cos z = 0$.
However $\cos z \neq 0$ for any $z \in \gamma$ (0)