

# PS7

Sunday, 22 November 2020

14:39

- a) Consider random variables  $X_i$  for  $i=1, \dots, 12$  representing the 12 bulbs. We have  $X_i \stackrel{iid}{\sim} \text{Bernoulli}(P)$  such that  $X_i = 1$  means the  $i$ th bulb is red and  $X_i = 0$  means the  $i$ th bulb is yellow. Then  $P \in \Theta = \{0.6, 0.25\}$ . We have hypotheses

$$H_0: P = 0.25$$

$$H_1: P = 0.6$$

and critical region

$$R = \{x: T(x) \geq 8\}$$

where our critical value is 8 and  $T(x)$  is the number of red bulbs which is the sum  $\sum_{i=1}^{12} X_i$

- b) We have the power function

$$\begin{aligned} \pi(P; T, c) &= P(T(x) \geq c; P) \\ &= P\left(\sum_{i=1}^{12} X_i \geq 8; P\right). \end{aligned}$$

By the definition of the binomial distribution  $Y = \sum_{i=1}^{12} X_i \sim \text{Binomial}(12, P)$ . So

$$\begin{aligned} \pi(P; t, c) &= P(Y \geq 8; P) \\ &= 1 - P(Y \leq 7; P) \\ &= 1 - \text{pbinom}(7, 12, P). \end{aligned}$$

The significance level is given by

$$\alpha = \sup_{P \in \Theta} \pi(P; t, c) = \pi(0.25) = 0.00278151.$$

The type II error is given by

$$1 - \pi(0.6) = 0.8618218$$

- c)  $\ln R$

- d) We have

$$T_{NP}(x) = f_n(x; 0.6) / f_n(x; 0.25) = \prod_{i=1}^n f(x_i; 0.6) / f(x_i; 0.25)$$

as  $X_i \stackrel{iid}{\sim} f_n(x; P)$ . Let  $t(x_i) = f(x_i; 0.6) / f(x_i; 0.25)$ . So

$$\begin{aligned} f(x_i) &= 0.6^{x_i} 0.4^{1-x_i} / 0.25^{x_i} 0.75^{1-x_i} \\ &= \left(\frac{0.6}{0.25}\right)^{x_i} \left(\frac{0.4}{0.75}\right)^{1-x_i} \\ &= \left(\frac{12}{5}\right)^{x_i} \left(\frac{8}{15}\right)^{1-x_i} \\ &= \left(\frac{9}{2}\right)^{x_i} \left(\frac{8}{15}\right)^{x_i} \left(\frac{8}{15}\right)^{1-x_i} \end{aligned}$$

$$= \left(\frac{9}{2}\right)^{x_i} \left(\frac{8}{15}\right)^{x_i} \left(\frac{8}{15}\right)^{1-x_i} \\ = \frac{8}{15} \left(\frac{9}{2}\right)^{x_i}$$

So

$$T_{NP}(x) = \prod_{i=1}^{15} \frac{8}{15} \left(\frac{9}{2}\right)^{x_i} = \left(\frac{8}{15}\right)^{15} \left(\frac{9}{2}\right)^{\sum_{i=1}^{15} x_i}$$

NOT SURE WHAT TO DO FROM HERE

e) Let  $T(x) = \sum_{i=1}^n x_i$  and  $\pi(p, T, c) = P(T(x) \geq c; p)$ . We know  $X = \sum_{i=1}^n x_i \sim \text{Binomial}(n, p)$ . So we need

$$\pi(0.25) = 0.05, [1 - \pi(0.6) = 0.1 \Leftrightarrow \pi(0.6) = 0.9]$$

or

$$\pi(0.6) - \pi(0.25) \approx 0.85$$

$$\Rightarrow g(n, c) := \pi(0.6) - \pi(0.25) - 0.85 \approx 0$$

Using R we find our solution to be  $n=15, c=7$ .

$$2a) f(x) = \frac{\lambda_1^x e^{-\lambda_1}}{x!} / \frac{\lambda_0^x e^{-\lambda_0}}{x!} = \left(\frac{\lambda_1}{\lambda_0}\right)^x e^{-\frac{\lambda_1}{\lambda_0}}$$

$$T(x) = \prod_{i=1}^n f(x) = \prod_{i=1}^n \left(\frac{\lambda_1}{\lambda_0}\right)^{x_i} e^{-\frac{\lambda_1}{\lambda_0}} = \left(\frac{\lambda_1}{\lambda_0}\right)^{\sum x_i} e^{-n \frac{\lambda_1}{\lambda_0}}$$

$$b) f(x) = \lambda_1 e^{-\lambda_1} x_i / \lambda_0 e^{-\lambda_0} x_i = \frac{\lambda_1}{\lambda_0} e^{-\frac{\lambda_1}{\lambda_0}}$$

$$T(x) = \prod_{i=1}^n f(x) = \prod_{i=1}^n \frac{\lambda_1}{\lambda_0} e^{-\frac{\lambda_1}{\lambda_0}} = \left(\frac{\lambda_1}{\lambda_0} e^{-\frac{\lambda_1}{\lambda_0}}\right)^n$$

$$c) f(x) = \frac{1}{\sigma_1 \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu_1}{\sigma_1}\right)^2} / \frac{1}{\sigma_0 \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu_0}{\sigma_0}\right)^2} = e^{\left(\frac{x-\mu_1}{\sigma_1}\right)^2}$$

$$T(x) = \prod_{i=1}^n f(x) = \prod_{i=1}^n e^{\left(\frac{x_i-\mu_1}{\sigma_1}\right)^2}$$

$$d) f(x) = \frac{1}{\sigma_1 \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu_1}{\sigma_1}\right)^2} / \frac{1}{\sigma_0 \sqrt{2\pi}} e^{-\frac{1}{2} \left(\frac{x-\mu_0}{\sigma_0}\right)^2} = \frac{\sigma_0}{\sigma_1} e^{\left(\frac{\sigma_0}{\sigma_1}\right)^2}$$

$$T(x) = \prod_{i=1}^n f(x) = \prod_{i=1}^n \frac{\sigma_0}{\sigma_1} e^{\left(\frac{\sigma_0}{\sigma_1}\right)^2} = \left(\frac{\sigma_0}{\sigma_1} e^{\left(\frac{\sigma_0}{\sigma_1}\right)^2}\right)^n$$

## Problem Sheet 7

### Problem Sheet 7

#### Question 1b

```
power <- function (p) 1-pbinom(7,12,p)
```

```
alpha <- power(0.25)
```

```
prob.type.II.error <- 1 - power(0.6)
```

```
alpha
```

```
## [1] 0.00278151
```

```
prob.type.II.error
```

```
## [1] 0.5618218
```

#### Question 1c


```
ps <- seq(0,1,0.001)
```

```
powers <- power(ps)
```

```
plot(ps, powers, type="l")
```

```
abline(v=c(0.25, 0.6), col=c("red", "blue"))
```

```
legend(x=0.01, y=0.99, legend=c("H0", "H1"),  
       col=c("red", "blue"), lty=1, cex=0.8)
```



### Question 1e

```
power <- function (p, n, c) 1-pbinom(c-1,n,p)
g <- function (n, c) power(0.6, n, c) - power(0.25, n, c) - 0.85

n.final <- 0
c.final <- 0
e.final <- Inf

for (n in seq(1, 50, 1)) {
  for (c in seq(1, 50, 1)) {
    e <- abs(g(n, c))
    if (e < e.final) {
      n.final <- n
      c.final <- c
      e.final <- e
    }
  }
}

c(n.final, c.final, e.final)
```

```
## [1] 15.000000000 7.000000000 0.001667718
```