To my arrow...

Jake Ross New Mexico Institute of Mining and Technology April, 2014

Geochronology of Southern McMurdo Sound and development of a micro laser furnace

by

Jake Ross

Submitted in Partial Fulfillment
of the Requirements for the Degree of
Doctor of Philosophy

New Mexico Institute of Mining and Technology Socorro, New Mexico April, 2014

ABSTRACT

Minna Bluff has been a significant topographic barrier to the flow of the Ross Ice Shelf since the mid-Miocene. Detailed Ar-Ar analyses of kaersutite and sanidine phenocrysts, and groundmass concentrates from volcanic units indicate an overall west to east progression of volcanic activity. Eruptions of basaltic to intermediate lavas, domes, and scoria cones started at 12 Ma in at what is now the eastern most point of Minna Bluff, Minna Hook. Activity was centered in this area for 4 Ma, constructing a pre-Minna Bluff island. Multiple glacial unconformities found at Minna Hook suggest repeated interaction with large warmbased, erosive ice sheets. Activity migrated westward from Minna Bluff Island at 7-8 Ma closing the gap created by the island and the mainland. Significant edifice construction continued until 4-5 Ma with sporadic and parasitic scoria cone eruptions, possibly associated with Mt. Discovery activity, continuing until 2 Ma.

The orientations of Minna Bluffs two major axes are strongly controlled by regional tectonic features. Minna Bluffs E-W axis, McIntosh Cliffs, is sub-parallel to the Radial Lineament and the N-S axis, Minna Hook, appears as extension of faulting bounding the Terror Rift. The constructional evolution of the 70km long volcanic complex has an important role in interpreting the climate signals recovered by the ANDRILL Project. Minna Bluff influenced the material delivered to the AND-1B drill site (ANDRILL MIS 2006-2007) in three critical ways: 1) Minna Bluff diverted upstream material, 2) provided

a pinning and stabilizing point for the Ross Ice Shelf, possible controlling

the calving line prior to the emergence of Ross Island, and 3) was a significant

source of fresh volcanic material throughout much of the period recovered by

ANDRILL MIS. For example, a kaersutite-bearing clast recovered from 822.78

mbsf in AND-1B yielded an age of 8.530.51 Ma, and was likely derived from

Minna Bluff. The results from this study can be incorporated into detailed glacier

and ice-sheet models of the McMurdo Sound region, a critical area in the Ross Ice

Sheet and global climate system. Jourdan et al. (2007)

Keywords: A; B; C

ACKNOWLEDGMENTS

Foo

This dissertation was typeset with LATEX¹ by the author.

¹The LATEX document preparation system was developed by Leslie Lamport as a special version of Donald Knuth's TeX program for computer typesetting. TeX is a trademark of the American Mathematical Society. The LATEX macro package for the New Mexico Institute of Mining and Technology dissertation format was written for the Tech Computer Center by John W. Shipman.

CONTENTS

LI	ST O	F TABLES	V
LI	ST O	F FIGURES v	7 i
ΡF	REFA	CE v	ii
1.	ANI	DRILL	1
2.	GEO	OCHRONOLOGY OF MINNA BLUFF, SOUTHERN MCMURDO	
	sou	JND, ANTARTICA	2
	2.1	Introduction	2
	2.2	Geology	2
		2.2.1 Volcanic	2
		2.2.2 Glacial	2
	2.3	Methods	3
		2.3.1 Ar-Ar	3
	2.4	Results	3
		2.4.1 Ar-Ar Laser Fusion	3
		2.4.2 Ar-Ar Step Heating	3
	2.5	Discussion	3

3.	DEV	DEVELOPMENT AND TESTING OF A LASER MICRO FURNACE FOR						
	AR-	AR ANALYSIS	4					
	3.1	Design	4					
	3.2	Testing	4					
	3.3	Preliminary Results	4					
4.		HRON: NOBLE GAS DATA ACQUISITION AND PROCESSING MEWORK	5					
Α.	AR-	AR DATA	6					
В.	ELE	CTRON MICROPROBE DATA	7					
RE	REFERENCES							

LIST OF TABLES

LIST OF FIGURES

This dissertation is a	accepted on behalf of the faculty of the Institute by the
lowing committee:	
William C. McIntosh	, Advisor
I release this docume	ent to the New Mexico Institute of Mining and Techno

PREFACE

Foo

ANDRILL

GEOCHRONOLOGY OF MINNA BLUFF, SOUTHERN MCMURDO SOUND, ANTARTICA

2.1 Introduction

Minna Bluff is a large volcanic penisula 70km south of Ross Island, Antarctica

2.2 Geology

2.2.1 Volcanic

This section describes the glacial geology of Minna Bluff. Here is the text wrapping around and a displaying a the outer indentation

2.2.2 Glacial

This section describes the volcanic geology of Minna Bluff. Here is the text wrapping around and a displaying a the outer indentation

- 2.3 Methods
 - 2.3.1 Ar-Ar
- 2.4 Results
 - 2.4.1 Ar-Ar Laser Fusion
 - 2.4.2 Ar-Ar Step Heating
- 2.5 Discussion

DEVELOPMENT AND TESTING OF A LASER MICRO FURNACE FOR AR-AR ANALYSIS

- 3.1 Design
- 3.2 Testing
- 3.3 Preliminary Results

PYCHRON: NOBLE GAS DATA ACQUISITION AND PROCESSING FRAMEWORK

APPENDIX A

AR-AR DATA

APPENDIX B ELECTRON MICROPROBE DATA

REFERENCES

Bachmann, O. and Dungan, M. A. (2002). Temperature-induced Al-zoning in hornblendes of the Fish Canyon magma, Colorado. *American Mineralogist*, 87(8-9):1062–1076.

Bachmann, O., Dungan, M. A., and Bussy, F. (2005). Insights into shallow magmatic processes in large silicic magma bodies: the trace element record in the Fish Canyon magma body, Colorado. *Contributions to Mineralogy and Petrology*, 149(3):338–349.

Bachmann, O., Dungan, M. A., and Lipman, P. W. (2002). The Fish Canyon magma body, San Juan volcanic field, Colorado: rejuvenation and eruption of an upper-crustal batholith. *Journal of Petrology*, 43(8):1469–1503.

Bachmann, O., OBERLI, F., DUNGAN, M., MEIER, M., Mundil, R., and FISCHER, H. (2007). 40Ar/39Ar and U–Pb dating of the Fish Canyon magmatic system, San Juan Volcanic field, Colorado: Evidence for an extended crystallization history. *Chemical Geology*, 236(1-2):134–166.

Charlier, B. L. A., Bachmann, O., Davidson, J. P., Dungan, M. A., and Morgan, D. J. (2007). The Upper Crustal Evolution of a Large Silicic Magma Body: Evidence from Crystal-scale Rb Sr Isotopic Heterogeneities in the Fish Canyon Magmatic System, Colorado. *Journal of Petrology*, 48(10):1875–1894.

Jourdan, F., Matzel, J., and Renne, P. R. (2007). 39Ar and 37Ar recoil loss during neutron irradiation of sanidine and plagioclase. *Geochimica et Cosmochimica Acta*, 71(11):2791–2808.

Jourdan, F. and Renne, P. R. (2007). Age calibration of the Fish Canyon sanidine 40Ar/39Ar dating standard using primary K–Ar standards. *Geochimica et Cosmochimica Acta*, 71(2):387–402.

Lipman, P., Dungan, M., and Bachmann, O. (1997). Comagmatic granophyric granite in the Fish Canyon Tuff, Colorado: implications for magma-chamber processes during a large ash-flow eruption. *Geology*, 25(10):915.

Lipman, P. W. and McIntosh, W. C. (2008). Eruptive and noneruptive calderas, northeastern San Juan Mountains, Colorado: Where did the ignimbrites come from? *Geological Society of America Bulletin*, 120(7-8):771–795.

Min, K., Mundil, R., Renne, P. R., and Ludwig, K. R. (2000). A test for systematic errors in 40Ar/39Ar geochronology through comparison with U/Pb analysis of a 1.1-Ga rhyolite. *Geochimica et Cosmochimica Acta*, 64(1):73–98.

Renne, P. R., Mundil, R., Balco, G., Min, K., and Ludwig, K. R. (2010). Joint determination of 40K decay constants and 40Ar*/40K for the Fish Canyon sanidine standard, and improved accuracy for 40Ar/39Ar geochronology. *Geochimica et Cosmochimica Acta*, 74(18):5349–5367.

Schmitz, M. D. and Bowring, S. A. (2001). U-Pb zircon and titanite systematics of the Fish Canyon Tuff: an assessment of high-precision U-Pb geochronology and its application to young volcanic rocks. *Geochimica et Cosmochimica Acta*, 65(15):2571–2587.

Geochronology of Southern McMurdo Sound and development of a micro laser furnace

by

Jake Ross

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the last page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and may require a fee.