

Homework 3

Jiří Klepl

(10 points) One of the preprocessing steps can be to eliminate some of the variables using so-called DP-elimination (or DP-resolution). In particular, assume we have a CNF φ and a variable x which we want to eliminate. Denote

$$\varphi_0 = \{C \in \varphi \mid \neg x \in C\}$$

$$\varphi_1 = \{C \in \varphi \mid x \in C\}$$

$$\varphi_r = \{C \in \varphi \mid C \cap \{x, \neg x\} = \emptyset\}$$

Namely, φ_0 consists of the clauses containing negative literal $\neg x$, φ_1 consists of the clauses containing positive literal x , φ_r contains the rest of the clauses. Let us now define

$$\varphi_{dp} = \{Res(C_0, C_1) \mid C_0 \in \varphi_0, C_1 \in \varphi_1\}$$

where $Res(C_0, C_1)$ denotes the clauses originating from C_0 and C_1 by resolution. Show that φ is equisatisfiable with $\varphi_0 = \varphi_r \wedge \varphi_{dp}$.

First, we define: $\varphi'_0 = \{C \setminus \{\neg x\} \mid C \in \varphi_0\}$, $\varphi'_1 = \{C \setminus \{x\} \mid C \in \varphi_1\}$

It follows that: $\varphi_0 \equiv \{x \rightarrow C \mid C \in \varphi'_0\}$, $\varphi_1 \equiv \{\neg x \rightarrow C \mid C \in \varphi'_1\}$ (1)

and from that: $\varphi_0 \equiv x \rightarrow \{C \mid C \in \varphi'_0\}$, $\varphi_1 \equiv \neg x \rightarrow \{C \mid C \in \varphi'_1\}$

and $\varphi_0 \wedge \varphi_1 \equiv x \rightarrow \{C \mid C \in \varphi'_0\} \wedge \neg x \rightarrow \{C \mid C \in \varphi'_1\}$ and (since those are the only two occurrences of variable x):

$$\varphi_0 \wedge \varphi_1 \equiv \{C \mid C \in \varphi'_0\} \vee \{C \mid C \in \varphi'_1\}$$

From distributivity of \wedge and \vee and the previous: $\varphi_0 \wedge \varphi_1 \equiv \{C_0 \vee C_1 \mid C_0 \in \varphi'_0, C_1 \in \varphi'_1\}$

We can finalize this as follows:

$$\varphi_0 \wedge \varphi_1 \equiv \{C_0 \vee C_1 \mid C_0 \in \varphi'_0, C_1 \in \varphi'_1\} \stackrel{Def}{=} \{Res(\{\neg x\} \cup C_0, \{x\} \cup C_1) \mid C_0 \in \varphi'_0, C_1 \in \varphi'_1\} \stackrel{(1)}{=}$$

$$\{Res(C_0, C_1) \mid C_0 \in \varphi_0, C_1 \in \varphi_1\} = \varphi_{dp}$$