

01204211 Discrete Mathematics

Lecture 8b: Modular arithmetic

Jittat Fakcharoenphol

October 1, 2024

Quick check 1

$$\begin{array}{l} 2/100 \quad 5/100 \\ \downarrow \\ 10/100 \end{array}$$

$$6/90 \quad 9/90$$

$$\underline{54/90}$$

X

If $a|m$ and $b|m$, can we say that $ab|m$? Prove this fact or provide a counter example.

$$\begin{array}{l} 2 \cdot 2 \cdot 5 \cdot 5 \\ \hline (2) \end{array}$$

$$\begin{array}{l} 2 \cdot 2 \cdot 5 \cdot 5 \\ \hline (5) \end{array}$$

If $\gcd(a, b) = 1,$



$$\begin{array}{l} 3 \cdot 3 \cdot 2 \cdot 5 \\ \hline (3 \cdot 2) \end{array}$$

$$\begin{array}{l} 3 \cdot 3 \cdot 2 \cdot 5 \\ \hline (3 \cdot 3) \end{array}$$

Quick check 2

If $a|m$, $b|m$, and $a \neq b$ are both prime, can we say that $ab|m$? Prove this fact or provide a counter example.

Prime factorization

One useful fact that we use over and over again is the following.

Unique Factorization (or Fundamental Theorem of Arithmetic)

Every integer greater than 1 can be written *uniquely* as a product of prime numbers (up to the order of factors).

Examples:

▶ $10 = 2 \cdot 5$

▶ $13 = 13$

▶ $112 = 2 \cdot 2 \cdot 2 \cdot 2 \cdot 7 = 2^4 \cdot 7$

There are 3 clocks. At this moment, all three clocks ring at the same time. The first clock rings every 3 hours, the second clock rings every 4 hours, and the third clock rings every 10 hours. How long do you have to wait until you would hear all clocks ring a the same time again?

You have a large water container and two smaller buckets. The first bucket carries 3 litres of water and the second bucket carries 5 litres of water.
Can you put exactly 1 litre of water in the water container?

You have a large water container and two smaller buckets. The first bucket carries 6 litres of water and the second bucket carries 15 litres of water.

What is the minimum volume of water you can exactly put in the water container?



$$a \cdot x + b \cdot y$$

You have a large water container and two smaller buckets. The first bucket carries 6 litres of water and the second bucket carries 15 litres of water.

What is the minimum volume of water you can exactly put in the water container?

In general if you have two buckets of volumes x and y , the amount that you can exactly make must be in the form of

$$ax + by,$$

for some integers x and y . (Note that x and y may be negative.)

You have a large water container and two smaller buckets. The first bucket carries 6 litres of water and the second bucket carries 15 litres of water.

What is the minimum volume of water you can exactly put in the water container?

In general if you have two buckets of volumes x and y , the amount that you can exactly make must be in the form of

$$ax + by,$$

for some integers x and y . (Note that x and y may be negative.)

Do you see why the sum must be divisible by any common divisor of x and y ?

Useful fact

Given integer x & y , can find ^{integers} a & b

$$\boxed{ax + by = \gcd(x, y)}$$

For any integer x and y , consider the term

$$a \cdot \underline{x} + b \cdot \underline{y},$$

for some integer a and b .

Useful fact

$$2 \cdot 4 + (-1) \cdot 6 = 2$$

$$\underline{3(7)} + \underline{(-5)(4)} = \underline{1}$$

For any integer x and y , consider the term

$$\underline{a \cdot x} + \underline{b \cdot y},$$

for some integer a and b .

When the term is ~~non zero~~, it must be divisible by $\underline{\gcd(x, y)}$, so it has to be at least positive $\gcd(x, y)$.

It turns out that you can actually attain that value, i.e., there exist a pair of integer a and b such that

$$\boxed{a} \cdot \overset{\downarrow}{x} + \boxed{b} \cdot \overset{\downarrow}{y} = \underline{\gcd(x, y)}.$$

Finding a and b : Extended Euclid Algorithm

We will modify the Euclid algorithm so that it also returns a and b together with $\gcd(x, y)$.

$$0 \cdot x + 1 \cdot y = y$$

Know:

$$a'y + b'(x \bmod y) = g$$

Goal:

$$a \cdot x + b \cdot y = g$$

Algorithm Euclid(x, y):

if $x \bmod y == 0$:

return y , 0 , 1

else:

$g, a', b' = \text{Euclid}(y, x \bmod y)$

$$a = b'$$

$$b = a' - b' \left\lfloor \frac{x}{y} \right\rfloor$$

return g, a, b

$\gcd(x, y)$

x	y	g	a	b
113	110	1	37	-38
110	3	1	-1	37
3	2	1	1	-1
2	1	1	0	1

$37 \cdot 113 + (-38) \cdot 110 = 1$
 $11 - \left\lfloor \frac{11}{3} \right\rfloor \cdot 3 = 2$
 $0 \cdot 2 + 1 \cdot 1 = 1$

Euclid(x, y) returns (g, a, b)
 $- g = \gcd(x, y)$

$$a \cdot x + b \cdot y = g$$

$$(x - \left\lfloor \frac{x}{y} \right\rfloor \cdot y) = x \bmod y$$

$$11 - \left\lfloor \frac{11}{3} \right\rfloor \cdot 3 = 2$$

Notes:

$$(g, a', b') = \text{Euclid}(y, x \bmod y)$$

We have a' and b' such that

$$a' \cdot y + b' \cdot (x \bmod y) = g.$$



$$y = a' \cdot y + b' \left(x - \left\lfloor \frac{x}{y} \right\rfloor \cdot y \right) = a' \cdot y + b' \cdot x - b' \cdot \left\lfloor \frac{x}{y} \right\rfloor \cdot y$$

$$= \underbrace{b' \cdot x}_a + \underbrace{\left(a' - b' \left\lfloor \frac{x}{y} \right\rfloor \right) \cdot y}_b$$

<u>x</u>	<u>y</u>	g	a = b'	b = a' - b' ⌊x/y⌋
y	x mod y	g	a'	b'

$$(a \cdot x + b \cdot y = 1)$$

Secret sharing




$k=2$

distribute S
to n people

- any set of k people
can recover S
- any set of $< k$ people
know nothing.

Secret sharing scheme based on straight lines



Days

What day is it today?

Days

What day is it today? Thursday.

Days

What day is it today? Thursday.
What day is 3 days after today?

Days

What day is it today? Thursday.

What day is 3 days after today? Sunday.

Days

What day is it today? Thursday.

What day is 3 days after today? Sunday.

What day is 20 days after today?

Days

Sat	Su	M	Tu	W	Th	F	Sa	Su	M	T	W	Th	F	Sa	Su
			29	30	31	1	2	3	4	5	6	7	8	9	10
								4	5	6	0	1	2	3	4

What day is it today? Thursday.

What day is 3 days after today? Sunday.

What day is 20 days after today? Wednesday.

Days

What day is it today? Thursday.

What day is 3 days after today? Sunday.

What day is 20 days after today? Wednesday.

What day is 10 days before today?

Days

What day is it today? Thursday.

What day is 3 days after today? Sunday.

What day is 20 days after today? Wednesday.

What day is 10 days before today? Monday.

Clocks

Suppose that it is 1 o'clock.

Clocks

Suppose that it is 1 o'clock.
What time is the next 5 hours?

Clocks

Suppose that it is 1 o'clock.
What time is the next 5 hours? 6 o'clock.

Clocks

Suppose that it is 1 o'clock.

What time is the next 5 hours? 6 o'clock.

What time is the next 10 hours?

Clocks

Suppose that it is 1 o'clock.

What time is the next 5 hours? 6 o'clock.

What time is the next 10 hours? 11 o'clock.

Clocks

Suppose that it is 1 o'clock.

What time is the next 5 hours? 6 o'clock.

What time is the next 10 hours? 11 o'clock.

What time is the next 20 hours?

Clocks

Suppose that it is 1 o'clock.

What time is the next 5 hours? 6 o'clock.

What time is the next 10 hours? 11 o'clock.

What time is the next 20 hours? 9 o'clock.

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus**

m .

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

$+$ $-$ \times

We can then define all arithmetic operations **modulo** m .

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

We can then define all arithmetic operations **modulo** m .

Suppose that $m = 7$. We would like to say that

$$4 + 5 = 9 \bmod m = 2.$$

Or

$$3 \cdot 4 = 5$$

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

We can then define all arithmetic operations **modulo** m .

Suppose that $m = 7$. We would like to say that

$$4 + 5 = 9 \bmod m = 2.$$

Or

$$3 \cdot 4 = 12 \bmod m =$$

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

We can then define all arithmetic operations **modulo** m .

Suppose that $m = 7$. We would like to say that

$$4 + 5 = 9 \bmod m = 2.$$

Or

$$3 \cdot 4 = 12 \bmod m = 5.$$

Or

$$2 - 6 = -4$$

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

We can then define all arithmetic operations **modulo** m .

Suppose that $m = 7$. We would like to say that

$$4 + 5 = 9 \bmod m = 2.$$

Or

$$3 \cdot 4 = 12 \bmod m = 5.$$

Or

$$2 - 6 = -4 \bmod 7 =$$

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

We can then define all arithmetic operations **modulo** m .

Suppose that $m = 7$. We would like to say that

$$4 + 5 = 9 \bmod m = 2.$$

Or

$$3 \cdot 4 = 12 \bmod m = 5.$$

Or

$$2 - 6 = \underline{-4} \bmod \underline{7} = \underline{3} \bmod 7 = \underline{3}.$$

Modular arithmetic

As in the days of weeks and clocks examples (and also as the modulo in RSA algorithm in our experiment), when working under modular arithmetic, we start with a **modulus** m .

We can then define all arithmetic operations **modulo** m .

Suppose that $m = 7$. We would like to say that

$$4 + 5 = 9 \bmod m = 2.$$

Or

$$3 \cdot 4 = 12 \bmod m = 5.$$

Or

$$2 - 6 = -4 \bmod 7 = 3 \bmod 7 = 3.$$

Note that when you view integers under the lense of modulus 7, these numbers


$$\dots, -19, -12, -5, 2, 9, 16, 23, \dots$$

are essentially **the same**.

Properties (1)

$$a \equiv b \pmod{m}$$

$a \bmod m = b \bmod m$, if and only if $m \mid a - b$.



Properties (1)

$a \bmod m = b \bmod m$, if and only if $m \mid a - b$.

Proof.

(\Rightarrow) Let $r = a \bmod m$. We can write

$$a = qm + r,$$

and

$$b = pm + r,$$

for some integers q and p . Thus, we have

$$a - b = qm + r - pm - r = (q - p)m.$$

Therefore $m \mid a - b$.

(\Leftarrow) Exercise.



Properties (2)

- ▶ $(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$
- ▶ $(a - b) \bmod m = ((a \bmod m) - (b \bmod m)) \bmod m$
- ▶ $(a \cdot b) \bmod m = ((a \bmod m) \cdot (b \bmod m)) \bmod m$

Congruences

$$5 \bmod 7 = 12 \bmod 7 = -2 \bmod 7$$

$$5 \equiv 12 \equiv -2 \pmod{7}$$

Definition (congruences)

For an integer $m > 0$, if integers a and b are such that

$$a \bmod m = b \bmod m,$$

we write

$$a \equiv b \pmod{m}.$$

Congruences

Definition (congruences)

For an integer $m > 0$, if integers a and b are such that

$$a \bmod m = b \bmod m,$$

we write

$$a \equiv b \pmod{m}.$$

We also have that

$$a \equiv b \pmod{m} \quad \Leftrightarrow \quad m \mid (a - b)$$

Congruences: properties (1)

- ▶ (reflexivity)

$$a \equiv a \pmod{m}.$$

- ▶ (symmetry)

$$a \equiv b \pmod{m} \text{ implies } b \equiv a \pmod{m}.$$

- ▶ (transitivity)

$$a \equiv b \pmod{m} \text{ and } b \equiv c \pmod{m} \text{ implies } a \equiv c \pmod{m}.$$

Congruences: properties (2) – operations

If we have that


$$a \equiv b \pmod{m},$$

and

$$c \equiv d \pmod{m},$$

then

- ▶ $a + c \equiv b + d \pmod{m}$
- ▶ $a - c \equiv b - d \pmod{m}$
- ▶ $ac \equiv bd \pmod{m}$



Congruences: properties (2) – operations

If we have that

$$a \equiv b \pmod{m},$$

and

$$c \equiv d \pmod{m},$$

then

- ▶ $a + c \equiv b + d \pmod{m}$
- ▶ $a - c \equiv b - d \pmod{m}$
- ▶ $ac \equiv bd \pmod{m}$

We can pretty much think of this “congruence” as a normal equation.

Congruences: properties (2) – operations

If we have that

$$a \equiv b \pmod{m},$$

and

$$c \equiv d \pmod{m},$$

then

- ▶ $a + c \equiv b + d \pmod{m}$
- ▶ $a - c \equiv b - d \pmod{m}$
- ▶ $ac \equiv bd \pmod{m}$

We can pretty much think of this “congruence” as a normal equation.

What is missing here?

Congruences: properties (2) – operations

If we have that

$$a \equiv b \pmod{m},$$

and

$$c \equiv d \pmod{m},$$

then

- ▶ $a + c \equiv b + d \pmod{m}$
- ▶ $a - c \equiv b - d \pmod{m}$
- ▶ $ac \equiv bd \pmod{m}$

We can pretty much think of this “congruence” as a normal equation.

What is missing here?

Division!

$$8x = 30 = 5 \cdot 6$$

$$\underline{x = 6}$$

Also, we wish we can do “cancellation”, i.e., if

$$xa \equiv xb \pmod{m},$$

then $a \equiv b \pmod{m}$. **BUT THIS IS NOT ALWAYS TRUE.**

Also, we wish we can do “cancellation”, i.e., if

$$xa \equiv xb \pmod{m},$$

then $a \equiv b \pmod{m}$. **BUT THIS IS NOT ALWAYS TRUE.**

Let's see the following example:

$$2 \cdot 1 \equiv 2 \cdot 3 \pmod{4},$$

but


$$1 \not\equiv 3 \pmod{4}.$$

Multiplications as functions

Let's view multiplication by 2 as a function, i.e., let $f(x) = 2 \cdot x \bmod 4$.

Multiplications as functions

Let's view multiplication by 2 as a function, i.e., let $f(x) = 2 \cdot x \bmod 4$.



$$\underline{2 \cdot x} \equiv \underline{2 \cdot y} \pmod{4}$$


$$\cancel{\downarrow}$$
$$\cancel{x \equiv y \pmod{4}}$$

$$3 - a \equiv 3 - b \pmod{4}$$

$$\downarrow \checkmark$$

$$a \equiv b \pmod{4}$$

Let's also see $g(x) = 3 \cdot x \bmod 4$.



Multiplications as functions

Let's view multiplication by 2 as a function, i.e., let $f(x) = 2 \cdot x \bmod 4$.

Let's also see $g(x) = 3 \cdot x \bmod 4$.

Which functions have inverses?



Multiplicative inverses (standard arithmetic)

In standard arithmetic, what is $2/5$?

Multiplicative inverses (standard arithmetic)

In standard arithmetic, what is $2/5$?

We are looking for a number x such that $2 = 5x$. How can we do that?



$$5x = 2$$

$$x \rightarrow 5^{-1} \cdot 5x = 5^{-1} \cdot 2$$

$$f(x) = 5x$$

$$f(x) = 2$$

$$f^{-1}(y)$$

Multiplicative inverses (standard arithmetic)

In standard arithmetic, what is $2/5$?

We are looking to a number x such that $2 = 5x$. How can we do that?

By dividing on both sides with 5:

$$2/5 = 5x/5 = x,$$

Multiplicative inverses (standard arithmetic)

$$(5^{-1}) 5 = 1$$

or $0.2 = \frac{1}{5}$

In standard arithmetic, what is $2/5$?

We are looking to a number x such that $2 = 5x$. How can we do that?

By dividing on both sides with 5:

$$2/5 = 5x/5 = x,$$

or equivalently, by multiplying with $(1/5) = 5^{-1}$:

$$2 \cdot 5^{-1} = 5x \cdot 5^{-1} = x \cdot 5 \cdot 5^{-1} = x \cdot 1 = x.$$

Here 5^{-1} is a multiplicative inverse of 5.

Multiplicative inverses (modular arithmetic)

You can do the same thing in modular arithmetic. Let the modulus be $m = 7$. Note that

$$5 \cdot 3 \equiv 15 \equiv 1 \pmod{7}.$$

$$\textcircled{2/5} \rightarrow \underline{5}x \equiv 2 \pmod{7}$$

Therefore, $5^{-1} \equiv \underline{3} \pmod{7}$.

$$\underline{x} \equiv 5^{-1} \cdot 5x \equiv 2 \cdot 5^{-1} \pmod{7}$$

$$\equiv 2 \cdot 3 \equiv \textcircled{6} \pmod{7}$$

$$5 \cdot 6 \equiv 30 \equiv \textcircled{2} \pmod{7}$$

Multiplicative inverses (modular arithmetic)

You can do the same thing in modular arithmetic. Let the modulus be $m = 7$. Note that

$$5 \cdot 3 \equiv 15 \equiv 1 \pmod{7}.$$

Therefore, $5^{-1} \equiv 3 \pmod{7}$.

To find $2/5$, we can view our goal as to find the value of x such that

$$2 \equiv 5x \pmod{7}.$$

We can multiply both sides with $5^{-1} \equiv 3$ to get

$$2 \cdot 5^{-1} \equiv 2 \cdot 3 \equiv 6 \equiv 5^{-1} \cdot 5x \equiv x \pmod{7}.$$

Multiplicative inverses (modular arithmetic)

You can do the same thing in modular arithmetic. Let the modulus be $m = 7$. Note that

$$5 \cdot 3 \equiv 15 \equiv 1 \pmod{7}.$$

Therefore, $5^{-1} \equiv 3 \pmod{7}$.

To find $2/5$, we can view our goal as to find the value of x such that

$$2 \equiv 5x \pmod{7}.$$

We can multiply both sides with $5^{-1} \equiv 3$ to get

$$2 \cdot 5^{-1} \equiv 2 \cdot 3 \equiv 6 \equiv 5^{-1} \cdot 5x \equiv x \pmod{7}.$$

Let's check:

$$5 \cdot 6 \equiv 30 \equiv 2 \pmod{7},$$

as required.

Multiplicative inverse modulo m

$$m \bmod 11 = m$$

$$\underline{3} \cdot \underline{4} = 12 \equiv 1 \pmod{11}$$

$$\begin{aligned} \underline{x} &\equiv 12x \equiv 3 \cdot 4 \cdot x \equiv 5 \cdot 3 \equiv 15 \pmod{11} \\ &\equiv 4 \pmod{11} \end{aligned}$$

Definition

The multiplicative inverse modulo \underline{m} of \underline{a} , denoted by \underline{a}^{-1} , is an integer such that

$$a \cdot \underline{a^{-1}} \equiv \underline{1} \pmod{m}.$$

$$4x \equiv 5 \pmod{11}$$

$$x \equiv 4 \pmod{11}$$

Multiplicative inverse modulo (11)

Let's try to figure out multiplicative inverse of every integer modulo 11.

a	$a^{-1} \pmod{11}$
1	1
2	6
3	4
4	3
5	9
6	2
7	8
8	7
9	5
10	10

$$12 \pmod{11} = 1$$

$$12 \text{ mu } 11 = 1$$

mod 13

1	1	7	2
2	7	8	5
3	9	9	3
4	10	10	4
5	8	11	6
6	11	12	12

13
26
39
52
65
78
91
104
117
130
143
156
169
182
195
208
221
234
247
260
273
286
299
312
325
338
351
364
377
390
403
416
429
442
455
468
481
494
507
520
533
546
559
572
585
598
611
624
637
650
663
676
689
702
715
728
741
754
767
780
793
806
819
832
845
858
871
884
897
910
923
936
949
962
975
988
1001

Example: secret sharing

- ▶ Think of a secret number $m \in \{0, 1, \dots, 10\}$.
- ▶ Pick a random number $a \in \{1, 2, \dots, 10\}$.
- ▶ Your straight line function $f(x) = (ax + m) \bmod 11$.
- ▶ We will generate 3 points from f and give them to 3 of your friends, each with only 1 point. Pick 3 numbers x_1, x_2, x_3 from $\{1, 2, \dots, 10\}$.
- ▶ Let's compute

$$(x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3)).$$

- ▶ Give them to 3 of your friends and challenge them to form a group of 2 people and figure out your number m .

Theorem 1



An integer a has a multiplicative inverse modulo m iff $\gcd(a, m) = 1$.

Proof.

(\Rightarrow) If $\gcd(a, m) > 1$, then there is no multiplicative inverse of a modulo m . We prove by contradiction. Assume that x is $a^{-1} \pmod{m}$ i.e.,

$$x \cdot a \equiv 1 \pmod{m}$$

This means $m \mid x \cdot a - 1$, this implies that there is an integer b s.t.

$$bm = x \cdot a - 1, \text{ or } 1 = x \cdot a + b(-m).$$

This means that $\gcd(a, m) \mid x \cdot a + b(-m)$, i.e., $\gcd(a, m) \mid 1$.

But this is impossible because $\gcd(a, m) > 1$. That's a contradiction.

Theorem 1

An integer a has a multiplicative inverse modulo m iff $\gcd(a, m) = 1$.

Proof.

(\Leftarrow) Recall that there exist integers x and y such that

$$x \cdot a + y \cdot m = \gcd(a, m) = 1.$$

Thus, $(x \cdot a + y \cdot m) \bmod m = x \cdot a \bmod m = 1 \bmod m$, i.e., $x \cdot a \equiv 1 \pmod{m}$. Therefore x is the inverse.

Theorem 1

$$\exists x, y \text{ integer } b \text{ s.t. } x \cdot a + y \cdot m = 1$$

An integer a has a multiplicative inverse modulo m iff $\gcd(a, m) = 1$.

$$b \cdot a \equiv 1 \pmod{m}$$

Proof.

(\Leftarrow) Recall that there exist integers x and y such that

$$x \cdot 113 + y \cdot d = 1$$

$$x \cdot a + y \cdot m = \gcd(a, m) = 1.$$

Thus, $(x \cdot a + y \cdot m) \bmod m = x \cdot a \bmod m = 1 \bmod m$, i.e., $x \cdot a \equiv 1 \pmod{m}$. Therefore x is the inverse.

(\Rightarrow) Let $r = \gcd(a, m)$. Suppose that b is the multiplicative inverse of a modulo m , i.e., we have that

$$b \cdot a \equiv 1 \pmod{m},$$

Thus, $ba \bmod m = 1 \bmod m = 1$, i.e., there exists an integer q such that

$$ba = qm + 1,$$

or $ba - qm = 1$. However, r since $r|a$ and $r|m$, r also divides $ba - qm$ and 1. But it $r \nmid 1$ because $r > 1$ and we have the contradiction. \square

Examples: division in modular arithmetic

Since the requirement for an existence of a^{-1} modulo m is that $\gcd(a, m) = 1$, if we let m be a prime number, every a which is not a multiple of m has an inverse.

Can you solve this equation?

$$4x + 9 \equiv 0 \pmod{11}.$$

Examples: division in modular arithmetic

Since the requirement for an existence of a^{-1} modulo m is that $\gcd(a, m) = 1$, if we let m be a prime number, every a which is not a multiple of m has an inverse.


Can you solve this equation?

$$4x + 9 \equiv 0 \pmod{11}.$$


We can even perform gaussian elimination (*which is very useful later*):

$$\begin{array}{rcl} 2x + y & \equiv & 3 \pmod{7} \\ x + 3y & \equiv & 5 \pmod{7} \end{array}$$

Public-key cryptography



public key



Digital signature



RSA

Quick recap: RSA

- ▶ Private key: (d, n) , Public key: (e, n)
- ▶ Encryption $E(m) = m^e \bmod n$, Decryption: $D(w) = w^d \bmod n$.
- ▶ Goal: Select e, d, n such that $D(E(m)) = m^{ed} \bmod n = m$.

Quick recap: RSA

$$d \equiv e^{-1} \pmod{(p-1)(q-1)}$$

$$\begin{aligned} (m^e)^d &\equiv m^{1+k(p-1)(q-1)} \pmod{pq} \\ &= m \cdot \cancel{m^{k(p-1)(q-1)}} \pmod{pq} \\ &\equiv m \pmod{pq} \end{aligned}$$

Euler Thm

- ▶ Private key: (d, n) , Public key: (e, n)
- ▶ Encryption $E(m) = m^e \pmod{n}$, Decryption: $D(w) = w^d \pmod{n}$.
- ▶ Goal: Select e, d, n such that $D(E(m)) = m^{ed} \pmod{n} = m$.
- ▶ Pick two primes p and q . Let $n = pq$. (Note: 65537 is circled in pink)
- ▶ Pick e (usually a small number)
- ▶ Pick d such that $d \equiv e^{-1} \pmod{(p-1)(q-1)}$, i.e., $ed \equiv 1 \pmod{(p-1)(q-1)}$, or
$$ed = k \cdot (p-1)(q-1) + 1,$$

for some integer k .

- ▶ What is $m^{ed} \pmod{n}$?

Polynomials & secret sharing.

What's next?

- ▶ We will prove Fermat's Little Theorem and show how to efficiently test if a number is prime.
- ▶ We will also use Fermat's Little Theorem to prove the correctness of RSA.
- ▶ Modular arithmetic is also key to our usage of polynomials to perform secret sharing and error correcting codes, because now we can do Gaussian elimination using only integers.