

01204211 Discrete Mathematics

Lecture 2c: Terminologies

Jittat Fakcharoenphol

July 4, 2021

Terminologies

These are terminologies used when showing mathematical facts.

- ▶ A **theorem** is a statement that can be argued to be true.
- ▶ A **proof** is the sequence of statements forming that mathematical argument.

Terminologies

These are terminologies used when showing mathematical facts.

- ▶ A **theorem** is a statement that can be argued to be true.
- ▶ A **proof** is the sequence of statements forming that mathematical argument.
- ▶ An **axiom** is a statement that is assumed to be true. (Note that we do not prove an axiom; therefore, the validity of a theorem proved using an axiom relies on the validity of the axiom.)

Terminologies

These are terminologies used when showing mathematical facts.

- ▶ A **theorem** is a statement that can be argued to be true.
- ▶ A **proof** is the sequence of statements forming that mathematical argument.
- ▶ An **axiom** is a statement that is assumed to be true. (Note that we do not prove an axiom; therefore, the validity of a theorem proved using an axiom relies on the validity of the axiom.)
- ▶ To prove a theorem, we may prove many simple lemmas to make our argument. A **lemma**, in this sense, is a smaller theorem (or a supportive one).

Terminologies

These are terminologies used when showing mathematical facts.

- ▶ A **theorem** is a statement that can be argued to be true.
- ▶ A **proof** is the sequence of statements forming that mathematical argument.
- ▶ An **axiom** is a statement that is assumed to be true. (Note that we do not prove an axiom; therefore, the validity of a theorem proved using an axiom relies on the validity of the axiom.)
- ▶ To prove a theorem, we may prove many simple lemmas to make our argument. A **lemma**, in this sense, is a smaller theorem (or a supportive one).
- ▶ A **corollary** is a theorem which is a “fairly” direct result of other theorems.

Terminologies

These are terminologies used when showing mathematical facts.

- ▶ A **theorem** is a statement that can be argued to be true.
- ▶ A **proof** is the sequence of statements forming that mathematical argument.
- ▶ An **axiom** is a statement that is assumed to be true. (Note that we do not prove an axiom; therefore, the validity of a theorem proved using an axiom relies on the validity of the axiom.)
- ▶ To prove a theorem, we may prove many simple lemmas to make our argument. A **lemma**, in this sense, is a smaller theorem (or a supportive one).
- ▶ A **corollary** is a theorem which is a “fairly” direct result of other theorems.
- ▶ A **conjecture** is a statement which we do not know if it is true or false.

Fermat's Last Theorem

Theorem: No three positive integers a , b , and c can satisfy the equation $a^n + b^n = c^n$ when $n > 2$.

This theorem has been conjectured by Pierre de Fermat in 1637. It remained a conjecture until Andrew Wiles proved it in 1994.

Goldbach's conjecture

Conjecture: Every even integer greater than 2 can be expressed as the sum of two primes.

In 1742, Christian Goldbach proposed this conjecture to Leonhard Euler. It remains unsolved.

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

1. A straight line segment can be drawn joining any two points.

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.
3. Given any straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.
3. Given any straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.
4. All right angles are congruent.

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.
3. Given any straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.
4. All right angles are congruent.
5. (The parallel postulate) If two lines are drawn which intersect a third in such a way that the sum of the inner angles on one side is less than two right angles, then the two lines inevitably must intersect each other on that side if extended far enough.

Euclid's axioms

Euclidean geometry is defined by the following 5 postulates (axioms).

1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.
3. Given any straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.
4. All right angles are congruent.
5. (The parallel postulate) If two lines are drawn which intersect a third in such a way that the sum of the inner angles on one side is less than two right angles, then the two lines inevitably must intersect each other on that side if extended far enough.

References: Weisstein, Eric W. "Euclid's Postulates." From MathWorld—A Wolfram Web Resource.

<http://mathworld.wolfram.com/EuclidsPostulates.html>

The triangle postulate

The following statement is called the triangle postulate.

The sum of the angles in every triangle is 180° .

The only way to prove this in Euclidean geometry is to use the parallel postulate. (Exercise: try to prove it.)

The triangle postulate

The following statement is called the triangle postulate.

The sum of the angles in every triangle is 180° .

The only way to prove this in Euclidean geometry is to use the parallel postulate. (Exercise: try to prove it.)

Is this statement always true everywhere in the world (or in the universe)?

The triangle postulate

The following statement is called the triangle postulate.

The sum of the angles in every triangle is 180° .

The only way to prove this in Euclidean geometry is to use the parallel postulate. (Exercise: try to prove it.)

Is this statement always true everywhere in the world (or in the universe)?

There are other geometries where Euclid's 5th postulate is not true; then the triangle postulate may not be true in those cases. Can you imagine one?