# 01204211 Discrete Mathematics Lecture 10a: Applications

Jittat Fakcharoenphol

September 6, 2022

A food factory can produce 2 products: a sleeping candy bar (S) and an energy bar (E).

- ▶ There are two ingredients: A and B. The factory has 120 units of A and 100 units of B.
- ➤ To produce 1 unit of candy bar S, you need 15 units of A and 10 units of B.
- ▶ To produce 1 unit of energy bar E, you need 10 units of A and 20 units of B.








A food factory can produce 2 products: a sleeping candy bar (S) and an energy bar (E).

- ▶ There are two ingredients: A and B. The factory has 120 units of A and 100 units of B.
- ▶ To produce 1 unit of candy bar S, you need 15 units of A and 10 units of B.
- ➤ To produce 1 unit of energy bar E, you need 10 units of A and 20 units of B.

How can we visualize the problem?

A food factory can produce 2 products: a sleeping candy bar (C) and an energy bar (E).

- There are two ingredients: A and B. The factory has 120 units of A and 100 units of B.
- ▶ To produce 1 unit of candy bar C, you need 15 units of A and 10 units of B.
- To produce 1 unit of energy bar E, you need 10 units of A and 20 units of B.




How can we choose the amount of C and E to produce?

How can we choose the amount of C and E to produce? It depends on the <u>prices</u> per unit of C and E.

MIC


How can we choose the amount of C and E to produce? It depends on the prices per unit of C and E. What if 1 unit of C is 1 baht and 1 unit of E is also 1 baht?




How can we choose the amount of C and E to produce?

How can we choose the amount of C and E to produce? It depends on the prices per unit of C and E.


How can we choose the amount of C and E to produce? It depends on the prices per unit of C and E. What if 1 unit of C is 1 baht and 1 unit of E is also 1 baht?



What if 1 unit of C is 10 baht and 1 unit of E is also 1 baht?



What if 1 unit of C is 1 baht and 1 unit of E is also 10 baht?



If we produce  $x_1$  units of C and  $x_2$  units of E, we would make

If we produce  $x_1$  units of C and  $x_2$  units of E, we would make

$$\overbrace{p_C \cdot x_1 + p_E \cdot x_2,}$$

where  $p_C$  and  $p_E$  are unit prices for C and E.


If we produce  $x_1$  units of C and  $x_2$  units of E, we would make

$$p_C \cdot x_1 + p_E \cdot x_2$$
,

where  $p_C$  and  $p_E$  are unit prices for C and E. This is called an <u>objective function</u>. So far, we tried 3 objective functions:


- $x_1 + x_2$
- $ightharpoonup 10 \cdot x_1 + x_2$
- $x_1 + 10 \cdot x_2$

Let's see what they look like again.





#### Vertices

Interesting things happen at the intersections. What are they?





## Simplex algorithm






# Simplex algorithm - How do you move?






A quick history of machine learning


- Perceptrons
- ► Neural networks
- Convolutional neural networks




#### Perceptrons


Invented in 1943 by McCulloch and Pitts.






# Perceptrons: XOR limit and multilayer perceptrons





XOR



## Inspiration from visual cortex study

