

01204211 Discrete Mathematics

Lecture 9b: Polynomials (1)¹

Jittat Fakcharoenphol

October 2, 2023

¹This section is from Berkeley CS70 lecture notes.

Quick exercise

For any integer $a \neq 1$, $a - 1 \mid a^2 - 1$.

Quick exercise

For any integer $a \neq 1$, $a - 1 \mid a^2 - 1$.

For any integer $a \neq 1$ and $n \geq 1$, $a - 1 \mid a^n - 1$.

Polynomials

A **single-variable polynomial** is a function $p(x)$ of the form

$$p(x) = \underline{a_d}x^{\underline{d}} + a_{d-1}x^{d-1} + \cdots + a_1x + a_0.$$


We call a_i 's *coefficients*. Usually, variable x and coefficients a_i 's are real numbers. The **degree** of a polynomial is the largest exponent of the terms with non-zero coefficients.

Examples

- ▶ $x^3 - 3x + 1$
- ▶ $x + 10$
- ▶ 10
- ▶ 0 ← zero polynomial

Folklore


degree = 1



know 2 points \Rightarrow
get $f(x)$


know < 2 points
 \Rightarrow know nothing

degree = 2



know 3 points
 \rightarrow get $f(x)$

degree = 3





know 4
points.

degree = d



$d+1$ points

Applications




► Secret sharing

Secret: S

Share S among n people
so that any group of
 k people can recover
the secret but any set
of $< k$ people know nothing.

$$f(x) = a_{k-1}x^{k-1} + a_{k-2}x^{k-2} + \dots + S$$




Share a secret
to n people,
where any k people
can recover the secret.



$(x, f(x)), \dots (-)$



Applications



- ▶ Secret sharing
- ▶ Error-correcting codes



Basic facts



Definition

a is a **root** of polynomial $f(x)$ if $f(a) = 0$.

Properties

- **Property 1:** A non-zero polynomial of degree d has at most d roots.
- **Property 2:** Given $d + 1$ pairs $(x_1, y_1), \dots, (x_{d+1}, y_{d+1})$ with distinct x_i 's, there is a *unique* polynomial $p(x)$ of degree at most d such that $p(x_i) = y_i$ for $1 \leq i \leq d + 1$.

Lemma 1 *

If two polynomials $f(x)$ and $g(x)$ of degree at most d that share $d+1$ points $(x_1, y_1), \dots, (x_{d+1}, y_{d+1})$, where all x_i 's are distinct, i.e., $f(x_i) = g(x_i) = y_i$, then $f(x) = g(x)$.



Proof.

Suppose that $f(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0$ and $g(x) = b_d x^d + b_{d-1} x^{d-1} + \dots + b_0$.

Let $h(x) = f(x) - g(x)$, i.e., let $h(x) = c_d x^d + c_{d-1} x^{d-1} + \dots + c_0$, where $c_i = a_i - b_i$. Note that $h(x)$ is also a polynomial of degree (at most) d .


We claim that $h(x)$ has $d+1$ roots. Note that since $f(x_i) = g(x_i) = y_i$, we have that

$$h(x_i) = f(x_i) - g(x_i) = y_i - y_i = 0,$$

i.e., every x_i is a root of $h(x)$.

From **Property 1**, if $h(x)$ is non-zero it has at most d roots; therefore, $h(x)$ must be zero, i.e.,


$f(x) - g(x) = 0$ or $f(x) = g(x)$ as required.



Polynomial interpolation - ideas

- polynomial degree (2)

$$\underline{y_1 \cdot \Delta_1(x) + y_2 \cdot \Delta_2(x) + y_3 \cdot \Delta_3(x)}$$



$$\Delta_1(x) = \frac{(x-x_2)(x-x_3)}{(x_1-x_2)(x_1-x_3)}$$

$(1, 0, 0)$

$$\Delta_2(x) = \frac{(x-x_1)(x-x_3)}{(x_2-x_1)(x_2-x_3)}$$

$$\Delta_3(x) = \frac{(x-x_1)(x-x_2)}{(x_3-x_1)(x_3-x_2)}$$

$(0, 0, 1)$

Lagrange polynomial


For $d+1$ points $(x_1, y_1), (x_2, y_2), \dots, (x_{d+1}, y_{d+1})$ where all x_i 's are distinct, let

$$\Delta_i(x) = \frac{(x - x_1)(x - x_2) \cdots (x - x_{i-1})(x - x_{i+1}) \cdots (x - x_{d+1})}{(x_i - x_1)(x_i - x_2) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_{d+1})}.$$

Note that $\Delta_i(x)$ is a polynomial of degree d

↳ is 1 at x_i -

↳ is 0 at other x_j $j \neq i$ -



Lagrange polynomial

For $d + 1$ points $(x_1, y_1), (x_2, y_2), \dots, (x_{d+1}, y_{d+1})$ where all x_i 's are distinct, let

$$\Delta_i(x) = \frac{(x - x_1)(x - x_2) \cdots (x - x_{i-1})(x - x_{i+1}) \cdots (x - x_{d+1})}{(x_i - x_1)(x_i - x_2) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_{d+1})}.$$

Note that $\Delta_i(x)$ is a polynomial of degree d . Also we have that

► For $j \neq i$, $\Delta_i(x_j) =$

Lagrange polynomial

For $d + 1$ points $(x_1, y_1), (x_2, y_2), \dots, (x_{d+1}, y_{d+1})$ where all x_i 's are distinct, let

$$\Delta_i(x) = \frac{(x - x_1)(x - x_2) \cdots (x - x_{i-1})(x - x_{i+1}) \cdots (x - x_{d+1})}{(x_i - x_1)(x_i - x_2) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_{d+1})}.$$

Note that $\Delta_i(x)$ is a polynomial of degree d . Also we have that

- ▶ For $j \neq i$, $\Delta_i(x_j) = 0$, and
- ▶ $\Delta_i(x_i) =$

Lagrange polynomial

For $d + 1$ points $(x_1, y_1), (x_2, y_2), \dots, (x_{d+1}, y_{d+1})$ where all x_i 's are distinct, let

$$\Delta_i(x) = \frac{(x - x_1)(x - x_2) \cdots (x - x_{i-1})(x - x_{i+1}) \cdots (x - x_{d+1})}{(x_i - x_1)(x_i - x_2) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_{d+1})}.$$

Note that $\Delta_i(x)$ is a polynomial of degree d . Also we have that

- ▶ For $j \neq i$, $\Delta_i(x_j) = 0$, and
- ▶ $\Delta_i(x_i) = 1$.

Lagrange polynomial

For $d + 1$ points $(x_1, y_1), (x_2, y_2), \dots, (x_{d+1}, y_{d+1})$ where all x_i 's are distinct, let

$$\Delta_i(x) = \frac{(x - x_1)(x - x_2) \cdots (x - x_{i-1})(x - x_{i+1}) \cdots (x - x_{d+1})}{(x_i - x_1)(x_i - x_2) \cdots (x_i - x_{i-1})(x_i - x_{i+1}) \cdots (x_i - x_{d+1})}.$$

Note that $\Delta_i(x)$ is a polynomial of degree d . Also we have that

- ▶ For $j \neq i$, $\Delta_i(x_j) = 0$, and
- ▶ $\Delta_i(x_i) = 1$.

We can use $\Delta_i(x)$ to construct a degree- d polynomial

$$p(x) = y_1 \cdot \Delta_1(x) + y_2 \cdot \Delta_2(x) + \cdots y_{d+1} \cdot \Delta_{d+1}(x).$$

What can you say about $p(x_i)$?

Property 2 *

Given $d+1$ pairs $(x_1, y_1), \dots, (x_{d+1}, y_{d+1})$ with distinct x_i 's, there is a **unique** polynomial $p(x)$ of degree at most d such that $p(x_i) = y_i$ for $1 \leq i \leq d+1$.

Proof of Property 2.

Using Lagrange interpolation, we know that there exists a polynomial $p(x)$ of degree d such that $p(x_i) = y_i$ for all $1 \leq i \leq d+1$.

For uniqueness, assume that there exists another polynomial $g(x)$ of degree d also satisfying the condition. Since $p(x)$ and $g(x)$ agree on more than d points, $p(x)$ and $g(x)$ must be equal from Lemma 1. □