

01204211 Discrete Mathematics

Lecture 13b: Eigenvalues and Eigenvectors

Jittat Fakcharoenphol

September 22, 2022

Review: Hamming codes (1)

Review: Hamming codes (1)

The code is defined by the generator matrix

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix}$$

Consider the encoding function $e : GF(2)^4 \rightarrow GF(2)^7$. Let $e(\mathbf{x}) = G\mathbf{x}$. What is $\text{Ker } e$?

Review: Hamming codes (1)

The code is defined by the generator matrix

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix}$$

Consider the encoding function $e : GF(2)^4 \rightarrow GF(2)^7$. Let $e(\mathbf{x}) = G\mathbf{x}$. What is $\text{Ker } e$?

What is $\dim \text{Im } e$?

Review: Hamming codes (2)

Review: Hamming codes (2)

The code is defined by the generator matrix $G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix}$

What can you say about the minimum “distance”?

Examples: random walks

Examples: differential equations (1)

Let's start with a simple system with one variable.

$$\frac{du}{dt} = au,$$

with $u = u(0)$ when $t = 0$.

Examples: differential equations (2)

Now consider a system with two variables v and w :

$$\begin{aligned}\frac{dv}{dt} &= 4v - 5w \\ \frac{dw}{dt} &= 2v - 3w\end{aligned}$$

with $v = 5$ and $w = 4$ when $t = 0$,

Examples: differential equations (2)

Now consider a system with two variables v and w :

$$\begin{aligned}\frac{dv}{dt} &= 4v - 5w \\ \frac{dw}{dt} &= 2v - 3w\end{aligned}$$

with $v = 5$ and $w = 4$ when $t = 0$, or if we let $u(t) = \begin{bmatrix} v(t) \\ w(t) \end{bmatrix}$ and

$$A = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix},$$

we have

$$\frac{du}{dt} = Au,$$

with $u(0) = \begin{bmatrix} 5 \\ 4 \end{bmatrix}$

Examples: differential equations (3)

Eigenvalues and eigenvectors

Definition

For an n -by- n matrix A , a vector v is an **eigenvector** of A if

$$Av = \lambda v,$$

and $v \neq \mathbf{0}$. The scalar λ is called an **eigenvalue** associated with v .

Example

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$.

If we let $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, we have

$$A\mathbf{v}_1 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

Example

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$.

If we let $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, we have

$$A\mathbf{v}_1 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 + 7 \\ -5 + 3 \end{bmatrix} =$$

Example

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$.

If we let $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, we have

$$A\mathbf{v}_1 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 + 7 \\ -5 + 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix} = (-2) \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

If we let $\mathbf{v}_2 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}$, we have

$$A\mathbf{v}_2 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 7 \\ 5 \end{bmatrix} =$$

Example

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$.

If we let $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, we have

$$A\mathbf{v}_1 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 + 7 \\ -5 + 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix} = (-2) \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

If we let $\mathbf{v}_2 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}$, we have

$$A\mathbf{v}_2 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 7 \\ 5 \end{bmatrix} = \begin{bmatrix} 35 + 35 \\ 35 + 15 \end{bmatrix} =$$

Example

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$.

If we let $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, we have

$$A\mathbf{v}_1 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 + 7 \\ -5 + 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix} = (-2) \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

If we let $\mathbf{v}_2 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}$, we have

$$A\mathbf{v}_2 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 7 \\ 5 \end{bmatrix} = \begin{bmatrix} 35 + 35 \\ 35 + 15 \end{bmatrix} = \begin{bmatrix} 70 \\ 50 \end{bmatrix} = 10 \cdot \begin{bmatrix} 7 \\ 5 \end{bmatrix}.$$

Example

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$.

If we let $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, we have

$$A\mathbf{v}_1 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 + 7 \\ -5 + 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix} = (-2) \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

If we let $\mathbf{v}_2 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}$, we have

$$A\mathbf{v}_2 = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 7 \\ 5 \end{bmatrix} = \begin{bmatrix} 35 + 35 \\ 35 + 15 \end{bmatrix} = \begin{bmatrix} 70 \\ 50 \end{bmatrix} = 10 \cdot \begin{bmatrix} 7 \\ 5 \end{bmatrix}.$$

See demo in colab.

Invariant subspace

Definition (invariant subspace)

For an n -by- n matrix A , subspace $\mathcal{V} \subseteq \mathbb{R}^n$ is called an **invariant subspace** under linear map $f(x) = Ax$ if for all $u \in \mathcal{V}$, $f(u) = Au \in \mathcal{V}$.

Invariant subspace

Definition (invariant subspace)

For an n -by- n matrix A , subspace $\mathcal{V} \subseteq \mathbb{R}^n$ is called an **invariant subspace** under linear map $f(x) = Ax$ if for all $u \in \mathcal{V}$, $f(u) = Au \in \mathcal{V}$.

Eigenvector

If v is an eigenvector of matrix A , then

$$\text{Span } \{v\}$$

is a 1-dimensional invariant subspace under linear map defined by A .

Finding eigenvalues and eigenvectors

Given A , we want to find an eigenvalue λ and a vector $\mathbf{u} \neq \mathbf{0}$ such that

$$A\mathbf{u} = \lambda\mathbf{u}.$$

Finding eigenvalues and eigenvectors

Given A , we want to find an eigenvalue λ and a vector $\mathbf{u} \neq \mathbf{0}$ such that

$$A\mathbf{u} = \lambda\mathbf{u}.$$

After some writing, we want to solve this equation

$$(A - \lambda I)\mathbf{u} = \mathbf{0},$$

where $\mathbf{u} \neq \mathbf{0}$.

Review: ranks and invertible matrices

Consider an n -by- n matrix A and the following linear system of equations

$$A\mathbf{x} = \mathbf{0}.$$

Review: ranks and invertible matrices

Consider an n -by- n matrix A and the following linear system of equations

$$Ax = \mathbf{0}.$$

Suppose that there exists $x \neq \mathbf{0}$ that satisfies the equation, what can you say about A ?

Review: ranks and invertible matrices

Consider an n -by- n matrix A and the following linear system of equations

$$Ax = \mathbf{0}.$$

Suppose that there exists $x \neq \mathbf{0}$ that satisfies the equation, what can you say about A ?

Clearly, A cannot have an inverse because no matrix B can bring x back from $Ax = 0$.

Review: ranks and invertible matrices

Consider an n -by- n matrix A and the following linear system of equations

$$Ax = \mathbf{0}.$$

Suppose that there exists $x \neq \mathbf{0}$ that satisfies the equation, what can you say about A ?

Clearly, A cannot have an inverse because no matrix B can bring x back from $Ax = 0$. In this case, we say that A is **singular**.

Review: ranks and invertible matrices

Consider an n -by- n matrix A and the following linear system of equations

$$Ax = \mathbf{0}.$$

Suppose that there exists $x \neq \mathbf{0}$ that satisfies the equation, what can you say about A ?

Clearly, A cannot have an inverse because no matrix B can bring x back from $Ax = 0$. In this case, we say that A is **singular**.

Review: ranks and invertible matrices

Consider an n -by- n matrix A and the following linear system of equations

$$Ax = \mathbf{0}.$$

Suppose that there exists $x \neq \mathbf{0}$ that satisfies the equation, what can you say about A ?

Clearly, A cannot have an inverse because no matrix B can bring x back from $Ax = 0$. In this case, we say that A is **singular**.

Equivalent conditions:

- ▶ The rank of A is less than n .
- ▶ Rows of A are not linearly independent.
- ▶ The linear function $f(x) = Ax$ is not injective.
- ▶ $\text{Ker } f \neq \{\mathbf{0}\}$.
- ▶ $\dim \text{Ker } f \neq 0$.

Finding λ

From this equation

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

Since we want it to have nonzero solution \mathbf{x} . Our goal is to find λ so that $A - \lambda I$ becomes singular.

Finding λ

From this equation

$$(A - \lambda I)x = \mathbf{0}.$$

Since we want it to have nonzero solution x . Our goal is to find λ so that $A - \lambda I$ becomes singular.

Typically, the tool to use is the **determinant**. However, we do not cover this topic in this class. We will look at small examples and consider an iterative method instead.

Example: 2×2 matrix

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We want to find λ such that

$$\begin{bmatrix} 5 - \lambda & 7 \\ 5 & 3 - \lambda \end{bmatrix}$$

is singular.

Example: 2×2 matrix

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We want to find λ such that

$$\begin{bmatrix} 5 - \lambda & 7 \\ 5 & 3 - \lambda \end{bmatrix}$$

is singular. This amounts to solving

$$\frac{5 - \lambda}{5} = \frac{7}{3 - \lambda},$$

Example: 2×2 matrix

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We want to find λ such that

$$\begin{bmatrix} 5 - \lambda & 7 \\ 5 & 3 - \lambda \end{bmatrix}$$

is singular. This amounts to solving

$$\frac{5 - \lambda}{5} = \frac{7}{3 - \lambda},$$

i.e.,

$$\lambda^2 - 8\lambda - 20 = 0.$$

Example: 2×2 matrix

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We want to find λ such that

$$\begin{bmatrix} 5 - \lambda & 7 \\ 5 & 3 - \lambda \end{bmatrix}$$

is singular. This amounts to solving

$$\frac{5 - \lambda}{5} = \frac{7}{3 - \lambda},$$

i.e.,

$$\lambda^2 - 8\lambda - 20 = 0.$$

The equation can be re-written as $(\lambda - 10)(\lambda + 2) = 0$; thus, it has 2 roots: 10 and -2 .

Example: 2×2 matrix

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We want to find λ such that

$$\begin{bmatrix} 5 - \lambda & 7 \\ 5 & 3 - \lambda \end{bmatrix}$$

is singular. This amounts to solving

$$\frac{5 - \lambda}{5} = \frac{7}{3 - \lambda},$$

i.e.,

$$\lambda^2 - 8\lambda - 20 = 0.$$

The equation can be re-written as $(\lambda - 10)(\lambda + 2) = 0$; thus, it has 2 roots: 10 and -2 .

You can find associated eigenvectors by solving corresponding $(A - \lambda I)x = \mathbf{0}$ equations.

Matrix multiplication (again)

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We know that A has two eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

with corresponding eigenvalues $\lambda_1 = 10$ and $\lambda_2 = -2$.

Matrix multiplication (again)

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We know that A has two eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

with corresponding eigenvalues $\lambda_1 = 10$ and $\lambda_2 = -2$.

What can we say about

$$\begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

Matrix multiplication (again)

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We know that A has two eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

with corresponding eigenvalues $\lambda_1 = 10$ and $\lambda_2 = -2$.

What can we say about

$$\begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 5 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 7-2 \\ 5+2 \end{bmatrix} =$$

Matrix multiplication (again)

Consider matrix $A = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix}$. We know that A has two eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 7 \\ 5 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

with corresponding eigenvalues $\lambda_1 = 10$ and $\lambda_2 = -2$.

What can we say about

$$\begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 5 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 7-2 \\ 5+2 \end{bmatrix} = \begin{bmatrix} 5 & 7 \\ 5 & 3 \end{bmatrix} \left(\begin{bmatrix} 7 \\ 5 \end{bmatrix} + 2 \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right)$$

Matrix multiplication (again and again)

Fact: An n -by- n matrix A has n linearly independent eigenvectors v_1, \dots, v_n with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$. (They might not be real vectors.)

Matrix multiplication (again and again)

Fact: An n -by- n matrix A has n linearly independent eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$. (They might not be real vectors.)

Since $\mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis, for any vector \mathbf{x} there exist $\alpha_1, \alpha_2, \dots, \alpha_n$ such that

$$\mathbf{x} = \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n.$$

Matrix multiplication (again and again)

Fact: An n -by- n matrix A has n linearly independent eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$. (They might not be real vectors.)

Since $\mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis, for any vector \mathbf{x} there exist $\alpha_1, \alpha_2, \dots, \alpha_n$ such that

$$\mathbf{x} = \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n.$$

Let's multiply \mathbf{x} with A :

$$\begin{aligned} A\mathbf{x} &= A(\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n) \\ &= A\alpha_1 \mathbf{v}_1 + A\alpha_2 \mathbf{v}_2 + \dots + A\alpha_n \mathbf{v}_n \\ &= \end{aligned}$$

Matrix multiplication (again and again)

Fact: An n -by- n matrix A has n linearly independent eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$. (They might not be real vectors.)

Since $\mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis, for any vector \mathbf{x} there exist $\alpha_1, \alpha_2, \dots, \alpha_n$ such that

$$\mathbf{x} = \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n.$$

Let's multiply \mathbf{x} with A :

$$\begin{aligned} A\mathbf{x} &= A(\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n) \\ &= A\alpha_1 \mathbf{v}_1 + A\alpha_2 \mathbf{v}_2 + \dots + A\alpha_n \mathbf{v}_n \\ &= \lambda_1 \alpha_1 \mathbf{v}_1 + \lambda_2 \alpha_2 \mathbf{v}_2 + \dots + \lambda_n \alpha_n \mathbf{v}_n. \end{aligned}$$

Matrix multiplication (again and again)

Fact: An n -by- n matrix A has n linearly independent eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$. (They might not be real vectors.)

Since $\mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis, for any vector \mathbf{x} there exist $\alpha_1, \alpha_2, \dots, \alpha_n$ such that

$$\mathbf{x} = \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n.$$

Let's multiply \mathbf{x} with A :

$$\begin{aligned} A\mathbf{x} &= A(\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n) \\ &= A\alpha_1 \mathbf{v}_1 + A\alpha_2 \mathbf{v}_2 + \dots + A\alpha_n \mathbf{v}_n \\ &= \lambda_1 \alpha_1 \mathbf{v}_1 + \lambda_2 \alpha_2 \mathbf{v}_2 + \dots + \lambda_n \alpha_n \mathbf{v}_n. \end{aligned}$$

We can keep multiplying with A many times:

$$A^k \mathbf{x} =$$

Matrix multiplication (again and again)

Fact: An n -by- n matrix A has n linearly independent eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ with corresponding eigenvalues $\lambda_1, \dots, \lambda_n$. (They might not be real vectors.)

Since $\mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis, for any vector \mathbf{x} there exist $\alpha_1, \alpha_2, \dots, \alpha_n$ such that

$$\mathbf{x} = \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n.$$

Let's multiply \mathbf{x} with A :

$$\begin{aligned} A\mathbf{x} &= A(\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n) \\ &= A\alpha_1 \mathbf{v}_1 + A\alpha_2 \mathbf{v}_2 + \dots + A\alpha_n \mathbf{v}_n \\ &= \lambda_1 \alpha_1 \mathbf{v}_1 + \lambda_2 \alpha_2 \mathbf{v}_2 + \dots + \lambda_n \alpha_n \mathbf{v}_n. \end{aligned}$$

We can keep multiplying with A many times:

$$A^k \mathbf{x} = \lambda_1^k \alpha_1 \mathbf{v}_1 + \lambda_2^k \alpha_2 \mathbf{v}_2 + \dots + \lambda_n^k \alpha_n \mathbf{v}_n.$$

The power method

If A has eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ such that

$$|\lambda_1| > |\lambda_i|,$$

for $i \neq 1$. We call λ_1 the **dominant eigenvalue**. We also call the eigenvectors corresponding to λ_1 **dominant eigenvectors**.

The power method (or power iteration)

- ▶ Start with a random vector x_0 .
- ▶ For $i = 0, 1, \dots, k$,
Let $x_{i+1} = Ax_i$, with probably some scaling.