

# 01204211 Discrete Mathematics

## Lecture 11b: Four fundamental subspaces (preview)

Jittat Fakcharoenphol


November 4, 2024

# What is a matrix?


Matrices arise in many places. We will see that there are essentially two ways to look at matrices.

$$\left[ \begin{array}{c|c|c} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 10 & 11 & 12 \end{array} \right] = \left[ \begin{array}{ccc} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 10 & 11 & 12 \end{array} \right] = \left[ \begin{array}{ccc} 1 & 2 & 3 \\ \hline 4 & 5 & 6 \\ \hline 7 & 8 & 9 \\ \hline 10 & 11 & 12 \end{array} \right]$$

# Matrices and elimination



row echelon form




row operation

$$R_2 \leftarrow R_2 + \alpha R_1$$

## Matrices and elimination

$$\begin{bmatrix} 1 & 2 & 0 & 1 \\ 1 & 5 & 1 & 1 \\ 2 & 10 & 2 & 4 \\ 2 & 7 & 1 & 10 \end{bmatrix}$$

# Matrices and elimination



$$Ax = b$$

if  $A$  is in  
row echelon form.

→ can find  
solution  
quickly.

$$\begin{bmatrix} 1 & 2 & 0 & 1 \\ 1 & 5 & 1 & 1 \\ 2 & 10 & 2 & 4 \\ 2 & 7 & 1 & 10 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & 3 & 1 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \phantom{0} \\ \phantom{0} \\ 1 \\ 0 \end{bmatrix}$$

$$3x_2 + x_3 = 1$$

is  $u$  a  
linear comb  
of rows?


$$u = \alpha_1 v + \alpha_2 w$$

$$\begin{cases} x_1 + 2x_2 + x_4 = 3 \\ 3x_2 + x_3 = 1 \\ 2x_4 = 2 \end{cases}$$


Back  
substitution.

Are there  $\alpha_1, \alpha_2$  such that  $u = \alpha_1 v + \alpha_2 w$

# Row echelon form



row echelon form



## Definition

A set of vectors  $\{u_1, \dots, u_k\} = U$

- ①  $U$  spans vector space  $V$
- ②  $u_1, u_2, \dots$  are linearly independent.

$\Rightarrow u_1, u_2, \dots, u_k$  is a basis of  $V$

Row Space of  $A$

$\triangleright v_1, v_2, \dots, v_k$  are linearly independent.

$\triangleright \text{Span} \{v_1, v_2, \dots, v_k\} = \text{Span} \{v'_1, v'_2, \dots, v'_m\} = C(A^T)$

$C(A^T) = \text{Span} \{v'_1, v'_2, \dots, v'_m\}$

recreate.  
from  $B \rightarrow A$

$v_1, v_2, \dots, v_k$  is a basis of  $C(A^T)$

## Linearly independent rows

## Vector spaces related to a matrix

Consider an  $m$ -by- $n$  matrix  $A$  over  $\mathbb{R}$ .



# Vector spaces related to a matrix

Consider an  $m$ -by- $n$  matrix  $A$  over  $\mathbb{R}$ .

We can view  $A$  as


- ▶  $n$  columns of  $m$ -vectors:  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$

## Vector spaces related to a matrix

Consider an  $m$ -by- $n$  matrix  $A$  over  $\mathbb{R}$ .

We can view  $A$  as

- ▶  $n$  columns of  $m$ -vectors:  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$
- ▶  $m$  rows of  $n$ -vectors:  $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m$



# Vector spaces related to a matrix

Consider an  $m$ -by- $n$  matrix  $A$  over  $\mathbb{R}$ .

We can view  $A$  as

- ▶  $n$  columns of  $m$ -vectors:  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$
- ▶  $m$  rows of  $n$ -vectors:  $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m$

When we have a set of vectors, recall that its span forms a vector space.

We have

# Vector spaces related to a matrix

Consider an  $m$ -by- $n$  matrix  $A$  over  $\mathbb{R}$ .

We can view  $A$  as

- ▶  $n$  columns of  $m$ -vectors:  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$
- ▶  $m$  rows of  $n$ -vectors:  $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m$

When we have a set of vectors, recall that its span forms a vector space.

We have

- ▶ Column space:  $\text{Span} \{ \mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n \} \subseteq \mathbb{R}^m$

# Vector spaces related to a matrix

Consider an  $m$ -by- $n$  matrix  $A$  over  $\mathbb{R}$ .

We can view  $A$  as

- ▶  $n$  columns of  $m$ -vectors:  $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$
- ▶  $m$  rows of  $n$ -vectors:  $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m$

When we have a set of vectors, recall that its span forms a vector space.

We have

- ▶ Column space:  $\text{Span} \{ \mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n \} \subseteq \mathbb{R}^m$
- ▶ Row space:  $\text{Span} \{ \mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m \} \subseteq \mathbb{R}^n$

# Subspaces

## Definition

Let  $\mathcal{V}$  and  $\mathcal{W}$  be vector spaces such that  $\mathcal{V} \subseteq \mathcal{W}$ . We say that  $\mathcal{V}$  is a **subspace** of  $\mathcal{W}$ .

# Subspaces

## Definition

Let  $\mathcal{V}$  and  $\mathcal{W}$  be vector spaces such that  $\mathcal{V} \subseteq \mathcal{W}$ . We say that  $\mathcal{V}$  is a **subspace** of  $\mathcal{W}$ .

## Examples:

- ▶  $\text{Span} \{[1, 1]\}$  is a subspace of  $\mathbb{R}^2$ .
- ▶  $\text{Span} \{[1, 0, 0], [0, 1, 1]\}$  is a subspace of  $\mathbb{R}^3$ .
- ▶  $\text{Span} \{[1, 0, 0], [0, 1, 1], [1, 1, 2]\}$  is a subspace of  $\mathbb{R}^3$ .

## Example 1

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$



## Example 1

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$

► Column space:

$$\mathcal{C}(A) = \{\alpha_1[1, 0] + \alpha_2[2, 1] + \alpha_3[4, 3] \mid \alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}\}$$

## Example 1

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$

► Column space:

$$\mathcal{C}(A) = \{\alpha_1[1, 0] + \alpha_2[2, 1] + \alpha_3[4, 3] \mid \alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}\} = \mathbb{R}^2.$$

► Row space:

$$\mathcal{C}(A^T) = \{\alpha_1[1, 2, 4] + \alpha_2[0, 1, 3] \mid \alpha_1, \alpha_2 \in \mathbb{R}\}$$

## Example 1

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$

► Column space:

$$\mathcal{C}(A) = \{\alpha_1[1, 0] + \alpha_2[2, 1] + \alpha_3[4, 3] \mid \alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}\} = \mathbb{R}^2.$$

► Row space:

$$\mathcal{C}(A^T) = \{\alpha_1[1, 2, 4] + \alpha_2[0, 1, 3] \mid \alpha_1, \alpha_2 \in \mathbb{R}\} \subseteq \mathbb{R}^3.$$

## Example 1 (cont.)

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$

Is there any other way to obtain vector spaces from  $A$ ?

## Example 1 (cont.)

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$

Is there any other way to obtain vector spaces from  $A$ ?

We can think of  $A$  as a coefficient matrix of a system of homogenous linear equations:

$$A\mathbf{x} = \mathbf{0}.$$

In this case, we have

$$\begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

## Example 1 (cont.)

Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix}$$

Is there any other way to obtain vector spaces from  $A$ ?

We can think of  $A$  as a coefficient matrix of a system of homogenous linear equations:

$$A\mathbf{x} = \mathbf{0}.$$

In this case, we have

$$\begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The set of solutions  $\{\mathbf{x} \mid A\mathbf{x} = \mathbf{0}\}$  form a vector space.

## Example 1 (cont.)

Given a matrix  $A$ , we can look at the matrix-vector product  $A\mathbf{x}$ .

Consider

$$\begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}.$$

# Four fundamental subspaces

## Four fundamental subspaces

Given an  $m$ -by- $n$  matrix  $A$ , we have the following subspaces

- ▶ The column space of  $A$  (denoted by  $\mathcal{C}(A)$  )
- ▶ The row space of  $A$  (denoted by  $\mathcal{C}(A^T)$  )
- ▶ The nullspace of  $A$

$$\mathcal{N}(A) = \{\mathbf{x} \mid A\mathbf{x} = \mathbf{0}\}$$

- ▶ The left nullspace of  $A$

$$\mathcal{N}(A^T) = \{\mathbf{y} \mid A^T\mathbf{y} = \mathbf{0}\}$$



# Four fundamental subspaces

## Four fundamental subspaces

Given an  $m$ -by- $n$  matrix  $A$ , we have the following subspaces

- ▶ The column space of  $A$  (denoted by  $\mathcal{C}(A) \subseteq \mathbb{R}^m$  )
- ▶ The row space of  $A$  (denoted by  $\mathcal{C}(A^T)$  )
- ▶ The nullspace of  $A$

$$\mathcal{N}(A) = \{\mathbf{x} \mid A\mathbf{x} = \mathbf{0}\}$$

- ▶ The left nullspace of  $A$

$$\mathcal{N}(A^T) = \{\mathbf{y} \mid A^T\mathbf{y} = \mathbf{0}\}$$

# Four fundamental subspaces

## Four fundamental subspaces

Given an  $m$ -by- $n$  matrix  $A$ , we have the following subspaces

- ▶ The column space of  $A$  (denoted by  $\mathcal{C}(A) \subseteq \mathbb{R}^m$ )
- ▶ The row space of  $A$  (denoted by  $\mathcal{C}(A^T) \subseteq \mathbb{R}^n$ )
- ▶ The nullspace of  $A$

$$\mathcal{N}(A) = \{ \underline{x} \mid A \underline{x} = \mathbf{0} \}$$

- ▶ The left nullspace of  $A$

$$\mathcal{N}(A^T) = \{ \mathbf{y} \mid A^T \mathbf{y} = \mathbf{0} \}$$

$$x \rightarrow \boxed{f} \rightarrow \cdot$$



$$\boxed{A} \begin{bmatrix} x \\ \vdots \end{bmatrix} = 0$$

$$\boxed{Ax = 0}$$
$$\underline{f(x) = Ax}$$

# Four fundamental subspaces

## Four fundamental subspaces

Given an  $m$ -by- $n$  matrix  $A$ , we have the following subspaces


- ▶ The column space of  $A$  (denoted by  $\mathcal{C}(A) \subseteq \mathbb{R}^m$  )
- ▶ The row space of  $A$  (denoted by  $\mathcal{C}(A^T) \subseteq \mathbb{R}^n$  )
- ▶ The nullspace of  $A$

$$\mathcal{N}(A) = \{\mathbf{x} \mid A\mathbf{x} = \mathbf{0}\} \subseteq \mathbb{R}^n$$

- ▶ The left nullspace of  $A$

$$\mathcal{N}(A^T) = \{\mathbf{y} \mid A^T\mathbf{y} = \mathbf{0}\}$$

# Four fundamental subspaces



## Four fundamental subspaces

Given an  $m$ -by- $n$  matrix  $A$ , we have the following subspaces

- ▶ The column space of  $A$  (denoted by  $\mathcal{C}(A) \subseteq \mathbb{R}^m$  )
- ▶ The row space of  $A$  (denoted by  $\mathcal{C}(A^T) \subseteq \mathbb{R}^n$  )
- ▶ The nullspace of  $A$

$$\mathcal{N}(A) = \{\mathbf{x} \mid A\mathbf{x} = \mathbf{0}\} \subseteq \mathbb{R}^n$$

$$\dim \mathcal{C}(A) = \dim \mathcal{C}(A^T) = k$$

$$\dim \mathcal{N}(A) = n - k$$

- ▶ The left nullspace of  $A$

$$\mathcal{N}(A^T) = \{\mathbf{y} \mid A^T \mathbf{y} = \mathbf{0}\} \subseteq \mathbb{R}^m$$

$$\dim \mathcal{N}(A^T) = m - k$$