01204211 Discrete Mathematics Lecture 4c: Mathematical Induction 3

Jittat Fakcharoenphol

July 17, 2021

Review: Mathematical Induction

Suppose that you want to prove that property P(n) is true for every natural number n.

Suppose that we can prove the following two facts:

Base case: P(1)

Inductive step: For any $k \ge 1$, $P(k) \Rightarrow P(k+1)$

The **Principle of Mathematical Induction** states that P(n) is true for every natural number n.

The assumption P(k) in the inductive step is usually referred to as the Induction Hypothesis.

Theorem 1 For any integer $n \geq 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} < 2$.

Theorem 1

For any integer
$$n \ge 1$$
, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} < 2$.

Proof.

The statement P(n) that we want to prove is " $\frac{1}{1^2}+\frac{1}{2^2}+\frac{1}{3^2}+\cdots+\frac{1}{n^2}<2$ ".

Theorem 1

For any integer $n \ge 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} < 2$.

Proof.

The statement P(n) that we want to prove is " $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{n^2} < 2$ ".

Case case: For n = 1, the statement is true because 1 < 2.

Theorem 1

For any integer $n \ge 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} < 2$.

Proof.

The statement P(n) that we want to prove is " $\frac{1}{1^2}+\frac{1}{2^2}+\frac{1}{3^2}+\cdots+\frac{1}{n^2}<2$ ".

Case case: For n = 1, the statement is true because 1 < 2.

Inductive step: For $k \geq 1$, let's assume P(k) and we prove that P(k+1) is true.

Theorem 1

For any integer $n \ge 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} < 2$.

Proof.

The statement P(n) that we want to prove is " $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} < 2$ ".

Case case: For n = 1, the statement is true because 1 < 2.

Inductive step: For $k \geq 1$, let's assume P(k) and we prove that P(k+1) is true.

The induction hypothesis is: $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{k^2} < 2$.

We want to show P(k+1), i.e.,

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} < 2.$$

Then...

▶ Is the assumption

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{k^2} < 2.$$

"strong" enough to prove

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} < 2$$
?

Why?

▶ Is the assumption

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{k^2} < 2.$$

"strong" enough to prove

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} < 2$$
?

Why?

▶ To prove P(k+1), we need a "gap" between the LHS and 2, so that we can add 1/(k+1) without blowing up the RHS.



- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.

- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.
 - 1/1 + 1/4 = 1.25.

- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.
 - ightharpoonup 1/1 + 1/4 = 1.25.
 - $1/1 + 1/4 + 1/9 \approx 1.361.$

- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.
 - 1/1 + 1/4 = 1.25.
 - $1/1 + 1/4 + 1/9 \approx 1.361.$
 - $1/1 + 1/4 + 1/9 + 1/16 \approx 1.4236.$

- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.
 - ightharpoonup 1/1 + 1/4 = 1.25.
 - $1/1 + 1/4 + 1/9 \approx 1.361.$
 - $1/1 + 1/4 + 1/9 + 1/16 \approx 1.4236.$
 - $1/1 + 1/4 + 1/9 + 1/16 + 1/25 \approx 1.4636.$

Yes, there is a gap. But how large?

- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.
 - 1/1 + 1/4 = 1.25.
 - $1/1 + 1/4 + 1/9 \approx 1.361.$
 - $1/1 + 1/4 + 1/9 + 1/16 \approx 1.4236.$
 - $1/1 + 1/4 + 1/9 + 1/16 + 1/25 \approx 1.4636.$

Yes, there is a gap. But how large?

▶ We need the gap to be large enough to insert $1/(k+1)^2$.

- Let's see a few values of the sum:
 - ightharpoonup 1/1 = 1.
 - 1/1 + 1/4 = 1.25.
 - $1/1 + 1/4 + 1/9 \approx 1.361.$
 - $1/1 + 1/4 + 1/9 + 1/16 \approx 1.4236.$
 - $1/1 + 1/4 + 1/9 + 1/16 + 1/25 \approx 1.4636.$

Yes, there is a gap. But how large?

- ▶ We need the gap to be large enough to insert $1/(k+1)^2$.
- After a "mysterious" moment, we observe that

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \le 2 - \frac{1}{n}.$$

Theorem 2 For any integer $n \ge 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \le 2 - \frac{1}{n}$.

Theorem 2

For any integer
$$n \ge 1$$
, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \le 2 - \frac{1}{n}$.

Proof.

(... the beginning is left out ...)

Inductive step: For $k \geq 1$, assume that $\frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{k^2} \leq 2 - \frac{1}{k}$.

Theorem 2

For any integer $n \ge 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \le 2 - \frac{1}{n}$.

Proof.

(... the beginning is left out ...)

Inductive step: For $k \ge 1$, assume that $\frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{k^2} \le 2 - \frac{1}{k}$.

Adding $1/(k+1)^2$ on both sides, we get

$$\frac{1}{1^2} + \frac{1}{2^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} \le 2 - \frac{1}{k} + \frac{1}{(k+1)^2} = 2 - \left(\frac{1}{k} - \frac{1}{(k+1)^2}\right).$$

Theorem 2

For any integer $n \ge 1$, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \le 2 - \frac{1}{n}$.

Proof.

(... the beginning is left out ...)

Inductive step: For $k \ge 1$, assume that $\frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{k^2} \le 2 - \frac{1}{k}$. Adding $1/(k+1)^2$ on both sides, we get

$$\frac{1}{1^2} + \frac{1}{2^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} \le 2 - \frac{1}{k} + \frac{1}{(k+1)^2} = 2 - \left(\frac{1}{k} - \frac{1}{(k+1)^2}\right).$$

Since 1/k - 1/(k+1) = 1/(k(k+1)), we have that

$$1/(k+1) = 1/k - 1/(k(k+1)) < 1/k - 1/(k+1)^{2}.$$

Theorem 2

For any integer
$$n \ge 1$$
, $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \le 2 - \frac{1}{n}$.

Proof.

(... the beginning is left out ...)

Inductive step: For $k \geq 1$, assume that $\frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{k^2} \leq 2 - \frac{1}{k}$. Adding $1/(k+1)^2$ on both sides, we get

$$\frac{1}{1^2} + \frac{1}{2^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} \le 2 - \frac{1}{k} + \frac{1}{(k+1)^2} = 2 - \left(\frac{1}{k} - \frac{1}{(k+1)^2}\right).$$

Since 1/k - 1/(k+1) = 1/(k(k+1)), we have that

$$1/(k+1) = 1/k - 1/(k(k+1)) < 1/k - 1/(k+1)^{2}.$$

Therefore, we conclude that

$$\frac{1}{1^2} + \frac{1}{2^2} + \dots + \frac{1}{k^2} + \frac{1}{(k+1)^2} \le 2 - \left(\frac{1}{k} - \frac{1}{(k+1)^2}\right) \le 2 - \frac{1}{k+1},$$

as required.



A Lesson learned

▶ Is a stronger statement easier to prove?

A Lesson learned

- Is a stronger statement easier to prove?
- In this case, the statement is indeed stronger, but the induction hypothesis gets stronger as well. Sometimes, this works out nicely.