




# 01204211 Discrete Mathematics



## Lecture 12b: Linear functions (I)

Jittat Fakcharoenphol



November 6, 2025

# Linear functions




## Linear functions

Consider vector spaces  $\mathcal{V}$  and  $\mathcal{W}$  over  $\mathbb{R}$ . A function  $f: \mathcal{V} \rightarrow \mathcal{W}$  is **linear** if

1. for all  $\underline{x}, \underline{y} \in \mathcal{V}$ ,  $f(\underline{x} + \underline{y}) = \underline{f(x)} + \underline{f(y)}$  and
2. for all  $\alpha \in \mathbb{R}$  and  $\underline{x} \in \mathcal{V}$ ,  $\underline{f(\alpha x)} = \alpha \underline{f(x)}$ .

## Example 1 - MLP



$$y_1 = \sum_{i=0}^n w_{1i} x_i$$

$$\bar{y} = W \bar{x}$$

$$\alpha \bar{y} = W \alpha \bar{x} = \alpha (W \bar{x})$$

$$\underline{W(\bar{x}_1 + \bar{x}_2)} = (W \bar{x}_1) + (W \bar{x}_2)$$

## Example 2 - Page rank (1)

## Example 2 - Page rank (2)

# Matrix-vector multiplication

Given an  $m \times n$  matrix  $M$  over  $\mathbb{R}$ , consider a product

$$M\mathbf{x}.$$

Note that for the multiplication to work,  $\mathbf{x}$  must be in  $\mathbb{R}^n$  and the result vector is in  $\mathbb{R}^m$ . Therefore, we can define a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  as

$$f(\mathbf{x}) = M\mathbf{x}.$$

Note that  $f$  is linear because:

$$f(\mathbf{x} + \mathbf{y}) = M(\mathbf{x} + \mathbf{y}) = M\mathbf{x} + M\mathbf{y} = f(\mathbf{x}) + f(\mathbf{y}),$$

and

$$f(\alpha\mathbf{x}) = M(\alpha\mathbf{x}) = \alpha M\mathbf{x} = \alpha f(\mathbf{x}).$$

# The converse

## Lemma 1

*For any linear function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , there exists an  $m \times n$  matrix  $M$  such that*

$$f(\mathbf{x}) = M\mathbf{x}.$$

## Proof.

Consider any  $x \in \mathbb{R}^n$ . Let  $\mathbf{x} = [x_1, x_2, \dots, x_n]$ . Note that

$$\mathbf{x} = [x_1, 0, \dots, 0] + [0, x_2, 0, \dots, 0] + \dots + [0, \dots, 0, x_n].$$

Let  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n \in \mathbb{R}^n$  be standard generators, i.e.,  $\mathbf{e}_i$  be a vector with 1 at the  $i$ -th row and 0 at every other positions. (For example  $\mathbf{e}_1 = [1, 0, \dots, 0]$  and  $\mathbf{e}_3 = [0, 0, 1, 0, \dots, 0]$ .)

We thus have

$$\mathbf{x} = x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \dots + x_n\mathbf{e}_n.$$

Since  $f$  is linear, this implies that

$$f(\mathbf{x}) = x_1f(\mathbf{e}_1) + x_2f(\mathbf{e}_2) + \dots + x_nf(\mathbf{e}_n).$$



## Proof (cont.)

Define  $M$  as follows


$$M = \left[ \begin{array}{c|c|c|c} f(\mathbf{e}_1) & f(\mathbf{e}_2) & \cdots & f(\mathbf{e}_n) \end{array} \right].$$

Hence,

$$\begin{aligned} M\mathbf{x} &= \left[ \begin{array}{c|c|c|c} f(\mathbf{e}_1) & f(\mathbf{e}_2) & \cdots & f(\mathbf{e}_n) \end{array} \right] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \\ &= x_1 f(\mathbf{e}_1) + x_2 f(\mathbf{e}_2) + \cdots + x_n f(\mathbf{e}_n) = f(\mathbf{x}), \end{aligned}$$

as required. □


## Structures of linear functions (overview)



$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} M = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

# Structures of linear functions (overview)


$$\{x \mid Mx = 0\}$$



$$M \quad m \times n \text{ matrix} \quad x \in \mathbb{R}^n$$

$$\begin{bmatrix} m & n \end{bmatrix} \quad \begin{bmatrix} x \end{bmatrix} = \begin{bmatrix} \end{bmatrix}_m \in \mathbb{R}^m$$

# Structures of linear functions (overview)



# Structures of linear functions (overview)

