



Course Name: Linear Algebra (MT 104)

Topic: Subspaces, Dimension, Bases (Exercise 2.8 & 2.9)

Instructor: Dr. Sara Aziz

saraazizpk@gmail.com

October 10, 2020

Subspaces of \mathbb{R}^n

One motivation for notion of subspaces of \mathbb{R}^n } algebraic generalization of geometric examples of lines and planes through the origin

$$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \quad \text{point} \\ \text{(no direction vectors)}$$

$$t \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \quad \text{line} \\ \text{(one direction vector)}$$

$$s \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \quad \text{plane} \\ \text{(two direction vectors)} \\ \underbrace{\hspace{1.5cm}}_{\text{linearly independent}}$$

$$r \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \quad \text{name?}$$

$$p \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + r \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \quad \text{name?}$$



Subspace

Definition A *subspace* S of \mathbb{R}^n is a set of vectors in \mathbb{R}^n such that

- (1) $\vec{0} \in S$ [contains zero vector]
- (2) if $\vec{u}, \vec{v} \in S$, then $\vec{u} + \vec{v} \in S$ [closed under addition]
- (3) if $\vec{u} \in S$ and $c \in \mathbb{R}$, then $c\vec{u} \in S$ [closed under scalar mult.]

Subspace

Example

Is $S = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : x \geq 0, y \geq 0 \right\}$
a subspace of \mathbb{R}^2 ?

No!

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \in S \quad \text{but} \quad - \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \notin S$$

$\Rightarrow S$ is not closed under scalar multiplication

Definition A *subspace* S of \mathbb{R}^n is a set of vectors in \mathbb{R}^n such that

- (1) $\vec{0} \in S$
- (2) if $\vec{u}, \vec{v} \in S$, then $\vec{u} + \vec{v} \in S$
- (3) if $\vec{u} \in S$ and $c \in \mathbb{R}$, then $c\vec{u} \in S$

Subspace

Example

Is $S = \left\{ \begin{bmatrix} a \\ b \\ 0 \end{bmatrix} : a, b \in \mathbb{R} \right\}$
a subspace of \mathbb{R}^3 ?

YES!

Definition A *subspace* S of \mathbb{R}^n is a set of vectors in \mathbb{R}^n such that

- (1) $\vec{0} \in S$
- (2) if $\vec{u}, \vec{v} \in S$, then $\vec{u} + \vec{v} \in S$
- (3) if $\vec{u} \in S$ and $c \in \mathbb{R}$, then $c\vec{u} \in S$

(1) $\vec{0} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \in S \Rightarrow$ contains zero vector ✓

(2) Let $\vec{u}, \vec{v} \in S$. Then $\vec{u} = \begin{bmatrix} a_1 \\ b_1 \\ 0 \end{bmatrix}$ and $\vec{v} = \begin{bmatrix} a_2 \\ b_2 \\ 0 \end{bmatrix}$ for some $a_1, a_2, b_1, b_2 \in \mathbb{R}$.
It follows that

$$\vec{u} + \vec{v} = \begin{bmatrix} a_1 + a_2 \\ b_1 + b_2 \\ 0 \end{bmatrix} \in S \Rightarrow \text{closed under addition} \quad \checkmark$$

(3) Let $\vec{u} \in S, c \in \mathbb{R}$. Then $\vec{u} = \begin{bmatrix} a \\ b \\ 0 \end{bmatrix}$ for some $a, b \in \mathbb{R}$. It follows that

$$c\vec{u} = c \begin{bmatrix} a \\ b \\ 0 \end{bmatrix} = \begin{bmatrix} ca \\ cb \\ 0 \end{bmatrix} \in S \Rightarrow \text{closed under scalar mult.} \quad \checkmark$$

Span is a subspace!

Theorem. Let $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k \in \mathbb{R}^n$. Then $S = \text{span}(\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k)$ is a subspace of \mathbb{R}^n .

Proof. We verify the three properties of the subspace definition.

(1) $\vec{0} = 0\vec{v}_1 + 0\vec{v}_2 + \dots + 0\vec{v}_k$

$\Rightarrow \vec{0}$ is a linear comb. of $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k \quad \Rightarrow \vec{0} \in S$

(2) Let $\vec{u}, \vec{w} \in S$. Then $\vec{u} = c_1\vec{v}_1 + \dots + c_k\vec{v}_k$ and

$\vec{w} = d_1\vec{v}_1 + \dots + d_k\vec{v}_k$ for some scalars c_i, d_i . Thus,

$$\begin{aligned}\vec{u} + \vec{w} &= (c_1\vec{v}_1 + \dots + c_k\vec{v}_k) + (d_1\vec{v}_1 + \dots + d_k\vec{v}_k) \\ &= \underbrace{(c_1 + d_1)\vec{v}_1 + \dots + (c_k + d_k)\vec{v}_k}_{\text{linear comb. of } \vec{v}_1, \dots, \vec{v}_k} \Rightarrow \vec{u} + \vec{w} \in S\end{aligned}$$

(3) Let $\vec{u} \in S, c \in \mathbb{R}$. You finish the proof (show that $c\vec{u} \in S$).



Column and row spaces of a matrix

Definition For an $m \times n$ matrix A with column vectors $v_1, v_2, \dots, v_n \in \mathbb{R}^m$, the *column space* of A is $\text{span}(v_1, v_2, \dots, v_n)$.

$\text{col}(A)$ is a subspace of \mathbb{R}^m since it is the span of a set of vectors in \mathbb{R}^m

Definition For an $m \times n$ matrix A with row vectors $r_1, r_2, \dots, r_m \in \mathbb{R}^n$, the *row space* of A is $\text{span}(r_1, r_2, \dots, r_m)$.

$\text{row}(A)$ is a subspace of \mathbb{R}^n since it is the span of a set of vectors in \mathbb{R}^n



Characterizing column and row spaces

Important relationships:

- **Column space**

$$\vec{b} \in \text{col}(A) \iff \vec{b} \text{ is in span of cols of } A \iff \exists \text{ scalars } x_1, x_2, \dots, x_n \text{ such that } x_1 \begin{bmatrix} | \\ \vec{v}_1 \\ | \end{bmatrix} + \dots + x_n \begin{bmatrix} | \\ \vec{v}_n \\ | \end{bmatrix} = \vec{b} \iff A\vec{x} = \vec{b} \text{ is consistent (has a sol'n)}$$

- **Row space**

$$\vec{b} \in \text{row}(A) \iff \vec{b}^T \in \text{col}(A^T) \iff A^T \vec{x} = \vec{b}^T \text{ has a solution}$$

since columns of A^T are the rows of A

OR

$$\vec{b} \in \text{row}(A) \iff \vec{b} \text{ is linear comb. of rows of } A \iff \left[\begin{array}{c} A \\ \vec{b} \end{array} \right] \xrightarrow[\text{for } i > j]{R_i + kR_j} \left[\begin{array}{c} A' \\ \vec{0} \end{array} \right]$$



Null space of a matrix

Definition For an $m \times n$ matrix A , the null space of A is the set of all solutions to $A\vec{x} = \vec{0}$, i.e.,

$$\text{null}(A) = \{\vec{x} : A\vec{x} = \vec{0}\}.$$

$\text{null}(A)$ is a set of vectors in \mathbb{R}^n

Basis

Definition A set of vectors $B = \{\vec{v}_1, \dots, \vec{v}_k\}$ is a *basis* for a subspace S of \mathbb{R}^n if

- $\text{span}(B) = S$,
- and B is a linearly independent set.

Example Standard basis for \mathbb{R}^3 is $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$

but another basis for \mathbb{R}^3 is $\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}.$

choice of
basis is not
unique

More on basis

Definition A set of vectors $B = \{\vec{v}_1, \dots, \vec{v}_k\}$ is a **basis** for a subspace S of \mathbb{R}^n if

- $\text{span}(B) = S$,
- and B is a linearly independent set.

Standard basis for \mathbb{R}^3 is $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$

but another basis for \mathbb{R}^3 is $\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}$.

Do you believe such bases
exist for \mathbb{R}^3 ?

Consider

$$B_1 = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} \right\}$$

$$B_2 = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}, \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}, \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix} \right\}$$

No!

Why not?

- $\text{span}(B_1) \neq \mathbb{R}^3$
- B_2 not linearly indep.

Definition The number of vectors in a basis of a subspace S is called the *dimension* of S .

Example $\dim(\mathbb{R}^n) = n$

since $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\} = \left\{ \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \right\}$ is a basis for \mathbb{R}^n

Side-note The trivial subspace $\{\vec{0}\}$ has no basis
since any set containing the zero vector is linearly dependent, so $\dim(\{\vec{0}\}) = 0$.



Theorem. The vectors $\{\vec{v}_1, \dots, \vec{v}_n\}$ form a basis of \mathbb{R}^n if and only if $\text{rank}(A) = n$, where A is the matrix with columns $\vec{v}_1, \dots, \vec{v}_n$.

Proof sketch (\Rightarrow).

$$\begin{array}{ccccccc} & & \Rightarrow & A\vec{x} = \vec{0} \text{ has only} & \Rightarrow & \text{cols of } A \text{ are} & \\ & & & \text{trivial sol'n } \vec{x} = \vec{0} & & \text{linearly indep.} & \Rightarrow \\ \text{rank}(A) = n \Rightarrow & \text{RREF of } A & & & & & \\ & \text{is } I & \Rightarrow & A\vec{x} = \vec{b} \text{ is consistent} & \Rightarrow & \text{cols of } A & \Rightarrow \\ & & & \text{for any } \vec{b} \in \mathbb{R}^n & & \text{span } \mathbb{R}^n & \\ & & & & & & \text{cols of } A \text{ are} \\ & & & & & & \text{basis for } \mathbb{R}^n \end{array}$$



Fundamental Theorem of Invertible Matrices (extended)

Theorem. Let A be an $n \times n$ matrix. The following statements are equivalent:

- A is invertible.
- $A\vec{x} = \vec{b}$ has a unique solution for all $\vec{b} \in \mathbb{R}^n$.
- $A\vec{x} = \vec{0}$ has only the trivial solution $\vec{x} = \vec{0}$.
- The RREF of A is I .
- A is the product of elementary matrices.
- $\text{rank}(A) = n$.
- Columns of A form a basis for \mathbb{R}^n .



Finding bases for fundamental subspaces of a matrix

Given matrix A , how do we
find bases for subspaces $\begin{cases} \text{row}(A) \\ \text{col}(A) \\ \text{null}(A) \end{cases} ?$

First, get RREF of A .

$$A \xrightarrow{\text{EROs}} R$$

Finding bases for fundamental subspaces of a matrix

EROs do not change
row space of a matrix.

basis for $\text{row}(A) = \text{basis for } \text{row}(R) \Rightarrow \text{nonzero rows of } R$

Columns of A have the
same dependence
relationship
as columns of R .

basis for $\text{col}(A) \Rightarrow \text{columns of } A \text{ that correspond to columns of } R \text{ w/leading 1's}$

- solve $A\vec{x} = \vec{0}$, i.e. solve $R\vec{x} = \vec{0}$
- express sol'n's in terms of free variables, e.g.,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = x_1 \begin{bmatrix} \vdots \\ \vdots \\ \vdots \end{bmatrix} + x_3 \begin{bmatrix} \vdots \\ \vdots \\ \vdots \end{bmatrix} \quad \text{basis vectors for null}(A)$$

Example of matrix subspaces' bases

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 10 \\ 11 & 12 & 13 & 14 & 15 \end{bmatrix} \quad A \xrightarrow{\text{EROs}} R = \begin{bmatrix} 1 & 0 & -1 & -2 & -3 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{basis for row}(A) = \{ [1 \ 0 \ -1 \ -2 \ -3], [0 \ 1 \ 2 \ 3 \ 4] \}$$

$$\text{basis for col}(A) = \left\{ \begin{bmatrix} 1 \\ 6 \\ 11 \end{bmatrix}, \begin{bmatrix} 2 \\ 7 \\ 12 \end{bmatrix} \right\}$$

Example of matrix subspaces' bases

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 10 \\ 11 & 12 & 13 & 14 & 15 \end{bmatrix} \quad A \xrightarrow{\text{EROs}} R = \begin{bmatrix} 1 & 0 & -1 & -2 & -3 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{aligned} x_1 - x_3 - 2x_4 - 3x_5 &= 0 \\ x_2 + 2x_3 + 3x_4 + 4x_5 &= 0 \\ x_3, x_4, x_5 &\text{ free} \end{aligned}$$

$$\text{basis for null}(A) = \left\{ \begin{bmatrix} 1 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ -4 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} x_3 + 2x_4 + 3x_5 \\ -2x_3 - 3x_4 - 4x_5 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_3 \begin{bmatrix} 1 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 2 \\ -3 \\ 0 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} 3 \\ -4 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Example related to column space

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \quad \vec{c} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad \begin{array}{l} \text{Is } b \in \text{col}(A)? \\ \text{Is } c \in \text{col}(A)? \end{array}$$

Determine if $A\vec{x} = \vec{b}$ has a solution.

$$\left[\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 1 & 1 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_2 - R_1} \left[\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

system is consistent
(has infinite # of sol'ns)

Yes, it is in column space of A.

A solution to the system gives scalar coefficients for linear combination.

$$\begin{array}{l} x_1 = 2 - x_3 \\ x_2 = 1 + x_3 \\ x_3 \text{ free} \end{array} \quad \begin{array}{l} \text{one} \\ \text{sol'n} \end{array} \quad \vec{x} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} \quad \vec{b} = 2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$



Example related to column space

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \quad \vec{c} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad \begin{array}{l} \text{Is } \vec{b} \in \text{col}(A)? \\ \text{Is } \vec{c} \in \text{col}(A)? \end{array}$$

Any vector in the column space of A
has 0 in its third component.

Thus, the vector \vec{c} is not in the column space of A .

Example related to row space

$$A = \begin{bmatrix} -6 & 3 \\ 1 & -\frac{1}{2} \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 2 & 1 \end{bmatrix} \quad \text{Is } \vec{b} \in \text{row}(A)?$$

Approach 1:

Is $\vec{b}^T \in \text{col}(A^T)$?

Determine if $A^T \vec{x} = \vec{b}^T$ has a solution.

$$\left[\begin{array}{cc|c} -6 & 1 & 2 \\ 3 & -\frac{1}{2} & 1 \end{array} \right] \xrightarrow{R_2 + \frac{1}{2}R_1} \left[\begin{array}{cc|c} -6 & 1 & 2 \\ 0 & 0 & 2 \end{array} \right]$$

inconsistent system

No, $\vec{b} \notin \text{row}(A)$.

Approach 2:

$\vec{b} \in \text{row}(A) \iff$

$$\left[\begin{array}{c} A \\ \vec{b} \end{array} \right] \xrightarrow[\text{for } i > j]{R_i + kR_j} \left[\begin{array}{c} A' \\ \mathbf{0} \end{array} \right]$$

$$\left[\begin{array}{cc|c} -6 & 3 & 2 \\ 1 & -\frac{1}{2} & 1 \\ 2 & 1 & 1 \end{array} \right] \xrightarrow[R_2 + \frac{1}{6}R_1]{R_3 - 2R_2} \left[\begin{array}{cc|c} -6 & 3 & 2 \\ 0 & 0 & 0 \\ 0 & 2 & -1 \end{array} \right]$$

No, $\vec{b} \notin \text{row}(A)$.

Example related to null space

$$A = \begin{bmatrix} 1 & -1 & 0 & 2 \\ 2 & -2 & 1 & 1 \\ 4 & -4 & 3 & -1 \end{bmatrix}$$

Find $\text{null}(A)$.

We need to solve $A\vec{x} = \vec{0}$.

We have

$$\left[\begin{array}{cccc|c} 1 & -1 & 0 & 2 & 0 \\ 2 & -2 & 1 & 1 & 0 \\ 4 & -4 & 3 & -1 & 0 \end{array} \right] \xrightarrow{\text{EROs}} \left[\begin{array}{cccc|c} 1 & -1 & 0 & 2 & 0 \\ 0 & 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

x_2, x_4 free vars

Convert to equations.

$$x_1 - x_2 + 2x_4 = 0$$

$$x_3 - 3x_4 = 0$$

Solve for x_1 and x_3 .

$$\vec{x} = \begin{bmatrix} x_2 - 2x_4 \\ x_2 \\ 3x_4 \\ x_4 \end{bmatrix} = x_2 \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -2 \\ 0 \\ 3 \\ 1 \end{bmatrix}$$

for $x_2, x_4 \in \mathbb{R}$

$$\text{null}(A) = \text{span} \left(\begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 3 \\ 1 \end{bmatrix} \right)$$