



Course Name: Linear Algebra

Course Code: MT 104

Instructor: Dr. Sara Aziz

saraazizpk@gmail.com

October 27, 2020



3.1 Introduction to Determinants

- Determinant via Cofactor Expansion
 - Examples
 - Theorem
- Triangular Matrices
 - Examples
 - Theorem



Notation: A_{ij} is the matrix obtained from matrix A by deleting the i th row and j th column of A .

Example

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix} \quad A_{23} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$$

Recall that $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$ and we let $\det[a] = a$.

For $n \geq 2$, the **determinant** of an $n \times n$ matrix $A = [a_{ij}]$ is

$$\det A = a_{11} \det A_{11} - a_{12} \det A_{12} + \cdots + (-1)^{1+n} a_{1n} \det A_{1n}$$
$$= \sum_{j=1}^n (-1)^{1+j} a_{1j} \det A_{1j}$$

Example

Compute the determinant of $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$

Solution

$$\det A = 1 \det \begin{bmatrix} -1 & 2 \\ 0 & 1 \end{bmatrix} - 2 \det \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} + 0 \det \begin{bmatrix} 3 & -1 \\ 2 & 0 \end{bmatrix}$$
$$= \text{-----} = \text{-----}$$

Common notation: $\det \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} = \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix}$.

So

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 1 \begin{vmatrix} -1 & 2 \\ 0 & 1 \end{vmatrix} - 2 \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} + 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix}$$



Cofactor

The **(i,j)-cofactor** of A is the number C_{ij} where

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

Example (Cofactor Expansion)

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 1C_{11} + 2C_{12} + 0C_{13}$$

(cofactor expansion across row 1)

Theorem (Cofactor Expansion)

The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column:

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}$$

(expansion across row i)

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}$$

(expansion down column j)

Use a matrix of signs to determine $(-1)^{i+j}$

$$\begin{bmatrix} + & - & + & \cdots \\ - & + & - & \cdots \\ + & - & + & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Example

Compute the determinant of $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$ using cofactor expansion down column 3.

Solution

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & 0 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 3 & -1 \end{vmatrix} = 1.$$

Example

Compute the determinant of $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{bmatrix}$

Solution:

$$\begin{aligned} & \begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \\ &= 1 \begin{vmatrix} 2 & 1 & 5 \\ 0 & 2 & 1 \\ 0 & 3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 2 & 3 & 4 \\ 0 & 2 & 1 \\ 0 & 3 & 5 \end{vmatrix} + 0 \begin{vmatrix} 2 & 3 & 4 \\ 2 & 1 & 5 \\ 0 & 3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 2 & 3 & 4 \\ 2 & 1 & 5 \\ 0 & 2 & 1 \end{vmatrix} \\ &= 1 \cdot 2 \begin{vmatrix} 2 & 1 \\ 3 & 5 \end{vmatrix} = 14 \end{aligned}$$



Triangular Matrices

$$\begin{bmatrix} * & * & \cdots & * & * \\ 0 & * & \cdots & * & * \\ 0 & 0 & \ddots & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & * \end{bmatrix}$$

(upper triangular)

$$\begin{bmatrix} * & 0 & 0 & 0 & 0 \\ * & * & 0 & 0 & 0 \\ * & * & \ddots & 0 & 0 \\ * & * & \cdots & * & 0 \\ * & * & \cdots & * & * \end{bmatrix}$$

(lower triangular)

Theorem

If A is a triangular matrix, then $\det A$ is the product of the main diagonal entries of A .



Example

$$\begin{vmatrix} 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & -3 & 5 \\ 0 & 0 & 0 & 4 \end{vmatrix} = \text{-----} = -24$$



3.2 Properties of Determinants

- Elementary Row Operations
 - Examples
 - Theorem
- Triangular Matrices
 - Examples
 - Theorem
- Determinant of the Transpose
- Multiplicative Property
 - Examples
 - Theorem

Theorem (Elementary Row Operations)

Let A be a square matrix.

- a. If a multiple of one row of A is added to another row of A to produce a matrix B , then $\det A = \det B$.
- b. If two rows of A are interchanged to produce B , then $\det B = -\det A$.
- c. If one row of A is multiplied by k to produce B , then $\det B = k \cdot \det A$.

Theorem still holds if the word *row* is replaced
with _____.

Example

$$\text{Compute } \begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 5 & 0 & 0 \\ 2 & 7 & 6 & 10 \\ 2 & 9 & 7 & 11 \end{vmatrix}.$$

Solution

$$\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 5 & 0 & 0 \\ 2 & 7 & 6 & 10 \\ 2 & 9 & 7 & 11 \end{vmatrix} = 5 \begin{vmatrix} 1 & 3 & 4 \\ 2 & 6 & 10 \\ 2 & 7 & 11 \end{vmatrix} = 5 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 0 & 2 \\ 2 & 7 & 11 \end{vmatrix} \\ = 5 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 0 & 2 \\ 0 & 1 & 3 \end{vmatrix} = -5 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 1 & 3 \\ 0 & 0 & 2 \end{vmatrix} = \text{-----} = \text{-----}.$$

Theorem (c) indicates that $\begin{vmatrix} * & * & * \\ -2k & 5k & 4k \\ * & * & * \end{vmatrix} = k \begin{vmatrix} * & * & * \\ -2 & 5 & 4 \\ * & * & * \end{vmatrix}.$

Example

Compute $\begin{vmatrix} 2 & 4 & 6 \\ 5 & 6 & 7 \\ 7 & 6 & 10 \end{vmatrix}$

Solution

$$\begin{vmatrix} 2 & 4 & 6 \\ 5 & 6 & 7 \\ 7 & 6 & 10 \end{vmatrix} = 2 \begin{vmatrix} 1 & 2 & 3 \\ 5 & 6 & 7 \\ 7 & 6 & 10 \end{vmatrix} = 2 \begin{vmatrix} 1 & 2 & 3 \\ 0 & -4 & -8 \\ 0 & -8 & -11 \end{vmatrix} \\ = 2(-4) \begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & -8 & -11 \end{vmatrix} = 2(-4) \begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 5 \end{vmatrix} = -40$$

Example

Compute $\begin{vmatrix} 2 & 3 & 0 & 1 \\ 4 & 7 & 0 & 3 \\ 7 & 9 & -2 & 4 \\ 1 & 2 & 0 & 4 \end{vmatrix}$ by row reduction and cofac. expansion.

Solution $\begin{vmatrix} 2 & 3 & 0 & 1 \\ 4 & 7 & 0 & 3 \\ 7 & 9 & -2 & 4 \\ 1 & 2 & 0 & 4 \end{vmatrix} = -2 \begin{vmatrix} 2 & 3 & 1 \\ 4 & 7 & 3 \\ 1 & 2 & 4 \end{vmatrix} = -2 \begin{vmatrix} 2 & 3 & 1 \\ 0 & 1 & 1 \\ 1 & 2 & 4 \end{vmatrix}$

$$= 2 \begin{vmatrix} 2 & 3 & 1 \\ 1 & 2 & 4 \\ 0 & 1 & 1 \end{vmatrix} = -2 \begin{vmatrix} 1 & 2 & 4 \\ 2 & 3 & 1 \\ 0 & 1 & 1 \end{vmatrix} = -2 \begin{vmatrix} 1 & 2 & 4 \\ 0 & -1 & -7 \\ 0 & 1 & 1 \end{vmatrix}$$

$$= -2 \begin{vmatrix} 1 & 2 & 4 \\ 0 & -1 & -7 \\ 0 & 0 & -6 \end{vmatrix} = -2(1)(-1)(-6) = -12.$$



Suppose A has been reduced to

$$U = \begin{bmatrix} \blacksquare & * & * & \cdots & * \\ 0 & \blacksquare & * & \cdots & * \\ 0 & 0 & \blacksquare & \cdots & * \\ 0 & 0 & 0 & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \blacksquare \end{bmatrix}$$

by row replacements and row interchanges, then

$$\det A = \begin{cases} (-1)^r \left(\begin{array}{l} \text{product of} \\ \text{pivots in } U \end{array} \right) & \text{when } A \text{ is invertible} \\ 0 & \text{when } A \text{ is not invertible} \end{cases}$$

Theorem

A square matrix is invertible if and only if $\det A \neq 0$.

Theorem

If A is an $n \times n$ matrix, then $\det A^T = \det A$.

Partial proof (2×2 case)

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc \quad \text{and}$$

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix}^T = \det \begin{bmatrix} a & c \\ b & d \end{bmatrix} = ad - bc$$

$$\Rightarrow \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \det \begin{bmatrix} a & c \\ b & d \end{bmatrix}.$$

(3 × 3 case)

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

$$\det \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & i \end{bmatrix} = a \begin{vmatrix} e & h \\ f & i \end{vmatrix} - b \begin{vmatrix} d & g \\ f & i \end{vmatrix} + c \begin{vmatrix} d & g \\ e & h \end{vmatrix}$$

$$\Rightarrow \det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \det \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & i \end{bmatrix}.$$

Theorem (Multiplicative Property)

For $n \times n$ matrices A and B , $\det(AB) = (\det A)(\det B)$.

Example

Compute $\det A^3$ if $\det A = 5$.

Solution: $\det A^3 = \det(AAA) = (\det A)(\det A)(\det A)$
 $= \text{-----} = \text{-----}.$

Example

For $n \times n$ matrices A and B , show that A is singular if $\det B \neq 0$ and $\det AB = 0$.

Solution: Since $(\det A)(\det B) = \det AB = 0$ and $\det B \neq 0$, then $\det A = 0$. Therefore A is singular.