

---

## Fall History

In the last \_\_\_\_\_ , how many falls have you experienced?

In the last 12 months, how many falls have you experienced that resulted in injuries and required medical attention?

---

### Interpretation

None

***Minimally Clinically Important Difference (MCID)***

***Minimal Detectable Change (MDC)***

None

---

### Selected References

1. Miller WC, Speechley M, Deathe B. The prevalence and risk factors of falling and fear of falling among lower extremity amputees. *Arch Phys Med Rehabil.* 2011;82(8):1031-1037.
  2. Hafner BJ, Smith DG. Differences in function and safety between Medicare Functional Classification Level-2 and -3 transfemoral amputees and influence of prosthetic knee joint control. *J Rehabil Res Dev.* 2009;46(3):417-434.
  3. Kahle JT, Highsmith MJ, Hubbard SL. Comparison of nonmicroprocessor knee mechanism versus C-Leg on Prosthesis Evaluation Questionnaire, stumbles, falls, walking tests, stair descent, and knee preference. *J Rehabil Res Dev.* 2008;45(1):1-14.
  4. Lord SR, Sherrington C, Menz HB, et al. Falls in Older People: Risk Factors and Strategies for Prevention (2nd ed.). New York, USA: Cambridge University Press, 2011.
-