

A Mini-Course on Optimization and Dynamics

From Euclidean Gradient Descent to Wasserstein Gradient Flow

Jia-Jie Zhu

Weierstrass Institute for Applied Analysis and Stochastics, Berlin

August 15, 2023

Euclidean gradient descent

Optimization in \mathbb{R}^d

$$\min_{x \in \mathbb{R}^d} f(x).$$

Euclidean gradient descent

Optimization in \mathbb{R}^d

$$\min_{x \in \mathbb{R}^d} f(x).$$

We optimize using the *gradient descent* algorithm

$$x_{k+1} = x_k - \tau_k \cdot \nabla f(x_k)$$

using the “variational principle”

$$x_{k+1} \in \operatorname{argmin}_x f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \frac{1}{2\tau_k} \|x - x_k\|_2^2$$

From GD to Mirror descent

We define the Bregman divergence associated with ϕ as

$$D_{\phi}(x, y) = \phi(x) - \phi(y) - \nabla\phi(y)^{\top}(x - y).$$

From GD to Mirror descent

We define the Bregman divergence associated with ϕ as

$$D_{\phi}(x, y) = \phi(x) - \phi(y) - \nabla\phi(y)^{\top}(x - y).$$

Mirror descent update (with quadratic term replaced by Bregman)

$$x_{k+1} = \operatorname{argmin}_x f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \frac{1}{\tau_k} D_{\phi}(x, x_k)$$

Example of MD: Euclidean norm

Mirror map $\phi(x) = \frac{1}{2}\|x\|_2^2$.

Example of MD: Euclidean norm

Mirror map $\phi(x) = \frac{1}{2}\|x\|_2^2$.

The resulting mirror descent algorithm

$$x_{k+1} = \operatorname{argmin}_x f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \frac{1}{2\tau_k} \|x - x_k\|_2^2.$$

This is equivalent to gradient descent with stepsize τ_k .

Example of MD: negative entropy

Mirror map $\phi(x) = \sum_{i=1}^d x(i) \log x(i)$.

Example of MD: negative entropy

Mirror map $\phi(x) = \sum_{i=1}^d x(i) \log x(i)$. The resulting Bregman divergence is the KL-divergence $D_\phi(x, y) = \sum_{i=1}^d x(i) \log \frac{x(i)}{y(i)}$.

Example of MD: negative entropy

Mirror map $\phi(x) = \sum_{i=1}^d x(i) \log x(i)$. The resulting Bregman divergence is the KL-divergence $D_\phi(x, y) = \sum_{i=1}^d x(i) \log \frac{x(i)}{y(i)}$.

If we restrict x to a (discrete probability) simplex $1^\top x = 1$, then the MD update

$$x_{k+1} = \operatorname{argmin}_{1^\top x=1} f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \frac{1}{\tau_k} D_\phi(x, x_k)$$

has the closed-form as *exponentiated gradient*

$$x(i)_{k+1} = \frac{x(i)_k e^{-\tau_k g_k}}{\sum_{j=1}^n x(j)_k e^{-\tau_k g_k}}, \quad i = 1, 2, \dots, n$$

Euclidean gradient descent as discretization of ODE gradient flow

$$x^{k+1} \in \operatorname{argmin}_x \langle \nabla f(x^k), x \rangle_{\mathbb{E}^d} + \frac{1}{2\tau} \|x - x^k\|^2$$

is the explicit Euler scheme for the ODE (for simplicity, we take constant time step τ)

$$\dot{x}(t) = -\nabla f(x(t)).$$

The solution $x(t)$ is an ODE gradient flow and the ODE is the gradient flow equation (GFE). In GF terms, the solution $x(t)$ is also called a **curve of maximal slope** (steepest descent).

Gradient flow dynamics: (nonlinear) ODE

$$\dot{x}(t) = -\nabla f(x(t))$$

Gradient flow dynamics: (nonlinear) ODE

$$\dot{x}(t) = -\nabla f(x(t))$$

$\dot{x}(t) \in X$ provides the **rate (or velocity)** (we can see)

Gradient flow dynamics: (nonlinear) ODE

$$\dot{x}(t) = -\nabla f(x(t))$$

$\dot{x}(t) \in X$ provides the **rate (or velocity)** (we can see)

$-\nabla f(x(t)) \in X^*$ provides the **(thermodynamic) force** (can't see; shadow price)

Gradient flow dynamics: (nonlinear) ODE

$$\dot{x}(t) = -\nabla f(x(t))$$

$\dot{x}(t) \in X$ provides the **rate (or velocity)** (we can see)

$-\nabla f(x(t)) \in X^*$ provides the **(thermodynamic) force** (can't see; shadow price)

The equation should be written in the **force-balance** form

$$\mathbb{I}_R \dot{x}(t) = -\nabla f(x(t)) \in X^*, \quad \mathbb{I}_R : X \rightarrow X^* \text{ is the Riesz isomorphism.}$$

If, in the non-Euclidean setting, $X \not\cong X^*$, then we have both force space and rate space GFE.

Energy dissipation balance (equality)

Fenchel-Young (proof is trivial; $\frac{a^2+b^2}{2} \geq ab$)

$$\psi(x) + \psi^*(\xi) \geq \langle x, \xi \rangle, \forall (x, \xi) \in X \times X^*.$$

Energy dissipation balance (equality)

Fenchel-Young (proof is trivial; $\frac{a^2+b^2}{2} \geq ab$)

$$\psi(x) + \psi^*(\xi) \geq \langle x, \xi \rangle, \forall (x, \xi) \in X \times X^*.$$

Furthermore, if ψ is proper, lsc, and convex, (x^*, ξ^*) is optimal.

By **Fenchel(-Young) duality and optimality**

$$\frac{d}{dt} f(x(t)) =_{X^*} \langle \nabla f(x(t)), \dot{x} \rangle_X$$

Energy dissipation balance (equality)

Fenchel-Young (proof is trivial; $\frac{a^2+b^2}{2} \geq ab$)

$$\psi(x) + \psi^*(\xi) \geq \langle x, \xi \rangle, \forall (x, \xi) \in X \times X^*.$$

Furthermore, if ψ is proper, lsc, and convex, (x^*, ξ^*) is optimal.

By **Fenchel(-Young) duality and optimality**

$$\frac{d}{dt} f(x(t)) =_{x^*} \langle \nabla f(x(t)), \dot{x} \rangle_X = -\|\nabla f(x(t))\|^2$$

Energy dissipation balance (equality)

Fenchel-Young (proof is trivial; $\frac{a^2+b^2}{2} \geq ab$)

$$\psi(x) + \psi^*(\xi) \geq \langle x, \xi \rangle, \forall (x, \xi) \in X \times X^*.$$

Furthermore, if ψ is proper, lsc, and convex, (x^*, ξ^*) is optimal.

By **Fenchel(-Young) duality and optimality**

$$\frac{d}{dt} f(x(t)) =_{x^*} \langle \nabla f(x(t)), \dot{x} \rangle_X = -\|\nabla f(x(t))\|^2 = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

Energy dissipation balance (equality)

Fenchel-Young (proof is trivial; $\frac{a^2+b^2}{2} \geq ab$)

$$\psi(x) + \psi^*(\xi) \geq \langle x, \xi \rangle, \forall (x, \xi) \in X \times X^*.$$

Furthermore, if ψ is proper, lsc, and convex, (x^*, ξ^*) is optimal.

By **Fenchel(-Young) duality and optimality**

$$\frac{d}{dt} f(x(t)) =_{x^*} \langle \nabla f(x(t)), \dot{x} \rangle_X = -\|\nabla f(x(t))\|^2 = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

Energy does not necessarily decrease along non-solutions, i.e., only inequality

$$\frac{d}{dt} f(z(t)) \geq -\left(\frac{1}{2}\|\dot{z}\|^2 + \frac{1}{2}\|\nabla f(z(t))\|^2\right).$$

Evolutionary variational inequality (EVI) $_{\lambda}$: ODE

Suppose the energy functional f is proper, upper semicontinuous, λ -convex for some $\lambda \in \mathbb{R}$, i.e., either convex or concave, $\forall s \in [0, 1], \forall u_0, u_1 \in \mathbb{R}^d$

$$f((1-s)u_0 + su_1) \leq (1-s)f(u_0) + sf(u_1) - \frac{\lambda}{2}s(1-s)\|u_0 - u_1\|^2.$$

and has compact sublevel sets. Then for any initial condition in the $x(0) \in \mathbb{R}^d$, there exists a unique solution at time t , $x(t) \in \mathbb{R}^d$.

Evolutionary variational inequality (EVI) $_{\lambda}$: ODE

Suppose the energy functional f is proper, upper semicontinuous, λ -convex for some $\lambda \in \mathbb{R}$, i.e., either convex or concave, $\forall s \in [0, 1], \forall u_0, u_1 \in \mathbb{R}^d$

$$f((1-s)u_0 + su_1) \leq (1-s)f(u_0) + sf(u_1) - \frac{\lambda}{2}s(1-s)\|u_0 - u_1\|^2.$$

and has compact sublevel sets. Then for any initial condition in the $x(0) \in \mathbb{R}^d$, there exists a unique solution at time t , $x(t) \in \mathbb{R}^d$.

Furthermore, the ODE solution $x(t)$ satisfies (EVI) $_{\lambda}$, for $t, s \in [0, T]$.

$$\frac{1}{2}\|x(t) - \nu\|^2 \leq \frac{1}{2}e^{-\lambda(t-s)}\|x(s) - \nu\|^2 + M_{\lambda}(t-s)(f(\nu) - f(x(t))),$$
$$M_{\lambda}(\tau) = \int_0^{\tau} e^{-\lambda(\tau-s)} \, ds, \quad \forall \nu \in \text{dom}(F) \subset \mathbb{R}^d.$$

Evolutionary variational inequality (EVI) $_{\lambda}$: ODE

Suppose the energy functional f is proper, upper semicontinuous, λ -convex for some $\lambda \in \mathbb{R}$, i.e., either convex or concave, $\forall s \in [0, 1], \forall u_0, u_1 \in \mathbb{R}^d$

$$f((1-s)u_0 + su_1) \leq (1-s)f(u_0) + sf(u_1) - \frac{\lambda}{2}s(1-s)\|u_0 - u_1\|^2.$$

and has compact sublevel sets. Then for any initial condition in the $x(0) \in \mathbb{R}^d$, there exists a unique solution at time t , $x(t) \in \mathbb{R}^d$.

Furthermore, the ODE solution $x(t)$ satisfies (EVI) $_{\lambda}$, for $t, s \in [0, T]$.

$$\frac{1}{2}\|x(t) - \nu\|^2 \leq \frac{1}{2}e^{-\lambda(t-s)}\|x(s) - \nu\|^2 + M_{\lambda}(t-s)(f(\nu) - f(x(t))),$$
$$M_{\lambda}(\tau) = \int_0^{\tau} e^{-\lambda(\tau-s)} ds, \quad \forall \nu \in \text{dom}(F) \subset \mathbb{R}^d.$$

Using (EVI) $_{\lambda}$, we can effortlessly extract convergence results. Suppose a minimizer of the energy exists $x^* \in \operatorname{arginf}_{x \in \mathbb{R}^d} f(x)$, we set $\nu = x^*, s = 0$ in (EVI) $_{\lambda}$

$$\begin{aligned} \|x(t) - x^*\|^2 &\leq e^{-\lambda t}\|x(0) - x^*\|^2 + 2M_{\lambda}(t-s) \left(\inf_{x \in \mathbb{R}^d} f(x) - f(x(t)) \right) \\ &\leq e^{-\lambda t}\|x(0) - x^*\|^2 \end{aligned}$$

Optimization over probability measures

$$\inf_{\mu \in \mathcal{P}} F(\mu)$$

- ▶ \mathcal{P} : set of probability measures; the probability “simplex”

Optimization over probability measures

$$\inf_{\mu \in \mathcal{P}} F(\mu)$$

- ▶ \mathcal{P} : set of probability measures; the probability “simplex”
- ▶ We will work with two types of (probability) measures

$$d\mu(x) = \rho(x) \, dx, \quad \mu = \sum_{i \in I} \alpha_i \delta_{x_i} \quad \alpha \in \Delta$$

Optimization over probability measures

$$\inf_{\mu \in \mathcal{P}} F(\mu)$$

- ▶ \mathcal{P} : set of probability measures; the probability “simplex”
- ▶ We will work with two types of (probability) measures

$$d\mu(x) = \rho(x) \, dx, \quad \mu = \sum_{i \in I} \alpha_i \delta_{x_i} \quad \alpha \in \Delta$$

- ▶ $M^+ \supseteq \mathcal{P}$: non-negative measures; “cone”

Optimization over probability measures

$$\inf_{\mu \in \mathcal{P}} F(\mu)$$

- ▶ \mathcal{P} : set of probability measures; the probability “simplex”
- ▶ We will work with two types of (probability) measures

$$d\mu(x) = \rho(x) \, dx, \quad \mu = \sum_{i \in I} \alpha_i \delta_{x_i} \quad \alpha \in \Delta$$

- ▶ $M^+ \supseteq \mathcal{P}$: non-negative measures; “cone”
- ▶ F : objective function; “energy”

Optimization over probability measures

What can't we just do gradient descent?

$$\mu^{k+1} = \mu^k - \tau_k \cdot \nabla F(\mu^k)$$

Optimization over probability measures

What can't we just do gradient descent?

$$\mu^{k+1} = \mu^k - \tau_k \cdot \nabla F(\mu^k)$$

- ▶ $\nabla F(\mu^k)$ is undefined

Optimization over probability measures

What can't we just do gradient descent?

$$\mu^{k+1} = \mu^k - \tau_k \cdot \nabla F(\mu^k)$$

- ▶ $\nabla F(\mu^k)$ is undefined
- ▶ μ^{k+1} must be a probability measure; care needs to be taken

Optimization over probability measures

What can't we just do gradient descent?

$$\mu^{k+1} = \mu^k - \tau_k \cdot \nabla F(\mu^k)$$

- ▶ $\nabla F(\mu^k)$ is undefined
- ▶ μ^{k+1} must be a probability measure; care needs to be taken
- ▶ What can we do instead?

A variational approach

Recall the “variational” formulation of gradient descent

$$x^{k+1} \in \operatorname{argmin}_x \langle \nabla f(x^k), x \rangle_{\mathbb{R}^d} + \frac{1}{2\tau} \|x - x^k\|^2 \iff x_{k+1} = x_k - \tau \cdot \nabla f(x_k)$$

for a suitable τ . This is the variational principle.

A variational approach

Recall the “variational” formulation of gradient descent

$$x^{k+1} \in \operatorname{argmin}_x \langle \nabla f(x^k), x \rangle_{\mathbb{R}^d} + \frac{1}{2\tau} \|x - x^k\|^2 \iff x_{k+1} = x_k - \tau \cdot \nabla f(x_k)$$

for a suitable τ . This is the variational principle.

Can we do the same for probability measures?

$$\mu^{k+1} \in \operatorname{arginf}_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{\tau} \mathcal{D}^2(\mu, \mu^k)$$

for some “distance” measure \mathcal{D} . This is sometimes called the *Minimizing Movement Scheme* (MMS).

Variational approach and MMS

$$\mu^{k+1} \in \operatorname{arginf}_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{\tau} \mathcal{D}(\mu, \mu^k)$$

Variational approach and MMS

$$\mu^{k+1} \in \operatorname{arginf}_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{\tau} \mathcal{D}(\mu, \mu^k)$$

We must specify the important ingredients

Energy : F

Geometry : \mathcal{D}

Variational approach and MMS

$$\mu^{k+1} \in \operatorname{arginf}_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{\tau} \mathcal{D}(\mu, \mu^k)$$

We must specify the important ingredients

Energy : F

Geometry : \mathcal{D}

The merit of the right gradient flow formulation of a dissipative evolution equation is that it separates energetics and kinetics: The **energetics** endow the state space with a **functional**, the **kinetics** endow the state space with a (Riemannian) **geometry** via the metric tensor. [Otto 2001]

Geometry: Wasserstein distance

Definition. The p -Wasserstein distance** between probability measures P, Q on \mathbb{R}^d (with p -th finite moments, $p \geq 1$) is defined through the following Kantorovich problem

$$W_p^p(P, Q) := \inf \left\{ \int |x - y|^p d\Pi(x, y) \mid \pi_{\#}^{(1)} \Pi = P, \pi_{\#}^{(2)} \Pi = Q \right\}$$

Geometry: Wasserstein distance

Definition. The p -Wasserstein distance** between probability measures P, Q on \mathbb{R}^d (with p -th finite moments, $p \geq 1$) is defined through the following Kantorovich problem

$$W_p^p(P, Q) := \inf \left\{ \int |x - y|^p d\Pi(x, y) \mid \pi_{\#}^{(1)}\Pi = P, \pi_{\#}^{(2)}\Pi = Q \right\}$$

Dual Kantorovich problem

$$W_p^p(P, Q) = \sup \left\{ \int \psi_1(x) dP(x) + \int \psi_2(y) dQ(y) \mid \psi_1(x) + \psi_2(y) \leq |x - y|^p \right\}$$

Geometry: Wasserstein distance

Definition. The p -Wasserstein distance** between probability measures P, Q on \mathbb{R}^d (with p -th finite moments, $p \geq 1$) is defined through the following Kantorovich problem

$$W_p^p(P, Q) := \inf \left\{ \int |x - y|^p d\Pi(x, y) \mid \pi_{\#}^{(1)}\Pi = P, \pi_{\#}^{(2)}\Pi = Q \right\}$$

Dual Kantorovich problem

$$W_p^p(P, Q) = \sup \left\{ \int \psi_1(x) dP(x) + \int \psi_2(y) dQ(y) \mid \psi_1(x) + \psi_2(y) \leq |x - y|^p \right\}$$

Dynamic formulation: **Benamou–Brenier**

$$W_2^2(P, Q) = \inf \left\{ \int_0^1 \int |v_t|^2 d\mu_t dt \mid \mu_0 = P, \mu_1 = Q, \frac{d}{dt}\mu_t + \operatorname{div}(v_t \mu_t) = 0 \right\}$$

Geometry: Wasserstein distance

Definition. The p -Wasserstein distance** between probability measures P, Q on \mathbb{R}^d (with p -th finite moments, $p \geq 1$) is defined through the following Kantorovich problem

$$W_p^p(P, Q) := \inf \left\{ \int |x - y|^p d\Pi(x, y) \mid \pi_{\#}^{(1)} \Pi = P, \pi_{\#}^{(2)} \Pi = Q \right\}$$

Dual Kantorovich problem

$$W_p^p(P, Q) = \sup \left\{ \int \psi_1(x) dP(x) + \int \psi_2(y) dQ(y) \mid \psi_1(x) + \psi_2(y) \leq |x - y|^p \right\}$$

Dynamic formulation: **Benamou–Brenier**

$$W_2^2(P, Q) = \inf \left\{ \int_0^1 \int |v_t|^2 d\mu_t dt \mid \mu_0 = P, \mu_1 = Q, \frac{d}{dt} \mu_t + \operatorname{div}(v_t \mu_t) = 0 \right\}$$

Entropy regularization (Sinkhorn divergence)

$$\inf_{\Pi} \int c(x, y) d\Pi(x, y) + \lambda D_{\phi}(\Pi \| P \otimes Q)$$

Geometry: (Csizsar) ϕ -divergence

Relative entropy is defined as

$$D_{\phi}(\mu|\nu) = \begin{cases} \int \phi\left(\frac{d\mu}{d\nu}\right) d\nu & \text{if } \mu \ll \nu \\ +\infty & \text{otherwise} \end{cases}$$

Geometry: (Csizsar) ϕ -divergence

Relative entropy is defined as

$$D_{\phi}(\mu|\nu) = \begin{cases} \int \phi\left(\frac{d\mu}{d\nu}\right) d\nu & \text{if } \mu \ll \nu \\ +\infty & \text{otherwise} \end{cases}$$

We can choose the ϕ functions from the following table to obtain: identity (trivial), Kullback, Hellinger, χ^2

Table 1: Entropy functions, their corresponding reverse entropy, and convex conjugates

Entropy f	f^*	Reverse entropy r	r^*
$f_{\text{Id}}(t) = \begin{cases} 0 & \text{if } t = 1 \\ +\infty & \text{otherwise} \end{cases}$	$f_{\text{Id}}^* = \text{Id}$	$r_{\text{Id}}(t) = \begin{cases} 0 & \text{if } t = 1 \\ +\infty & \text{otherwise} \end{cases}$	$r_{\text{Id}}^* = \text{Id}$
$f_{\text{KL}}(t) = t \log t - t + 1$	$f_{\text{KL}}^*(s) = e^s - 1$	$r_{\text{KL}}(t) = t - 1 - \log t$	$r_{\text{KL}}^*(s) = -\log(1 - s)$
$f_{\text{H}}(t) = (\sqrt{t} - 1)^2$	$f_{\text{H}}^*(s) = s/(1 - s)$	$r_{\text{H}}(t) = (\sqrt{t} - 1)^2$	$r_{\text{H}}^*(s) = s/(1 - s)$
$f_{\chi^2}(t) = (t - 1)^2$	$f_{\chi^2}^*(s) = s^2/4 + s$	$r_{\chi^2}(t) = (t - 1)^2/t$	$r_{\chi^2}^*(s) = 2 - \sqrt{1 - s}$

(table: J. Zhu)

Preliminary: first variation over measures and subdifferentials

The **first variation of a functional** F at $\mu \in \mathcal{P}$ is defined as a function $\frac{\delta F}{\delta \mu}[\mu]$

$$\frac{d}{d\epsilon} F(\mu + \epsilon \cdot \nu)|_{\epsilon=0} = \int \frac{\delta F}{\delta \mu}[\mu](x) \, d\nu(x)$$

for any perturbation in measure ν such that $\mu + \epsilon \cdot \nu \in \mathcal{P}$.

The **variational principle**

$$\frac{d}{d\epsilon} F(\mu + \epsilon \cdot \nu)|_{\epsilon=0} = 0$$

for all variation ν , states the “optimality condition”.

We also summon the Fréchet differential on a Banach space X as a set in the dual space

$$DF := \{\xi \in X^* \mid F(\mu) \geq F(\nu) + \langle \xi, \mu - \nu \rangle_X + o(\|\mu - \nu\|_X) \text{ for } \mu \rightarrow \nu\}$$

Three types of energy functionals

Suppose $\mu(x) = \rho(x) \, dx$ WLOG,

$$\mathcal{F}(\varrho) = \int f(\varrho(x)) dx, \quad \mathcal{V}(\varrho) = \int V(x) d\varrho, \quad \mathcal{W}(\varrho) = \frac{1}{2} \iint W(x-y) d\varrho(x) d\varrho(y)$$

Three types of energy functionals

Suppose $\mu(x) = \rho(x) \, dx$ WLOG,

$$\mathcal{F}(\varrho) = \int f(\varrho(x)) dx, \quad \mathcal{V}(\varrho) = \int V(x) d\varrho, \quad \mathcal{W}(\varrho) = \frac{1}{2} \iint W(x-y) d\varrho(x) d\varrho(y)$$

We calculate the first variations (by following the definition)

$$\frac{\delta \mathcal{F}}{\delta \varrho}(\varrho) = f'(\varrho), \quad \frac{\delta \mathcal{V}}{\delta \varrho}(\varrho) = V, \quad \frac{\delta \mathcal{W}}{\delta \varrho}(\varrho) = W * \varrho$$

Back to (discrete-time) gradient flow

Optimization

$$\inf_{\mu \in \mathcal{P}} F(\mu)$$

Back to (discrete-time) gradient flow

Optimization

$$\inf_{\mu \in \mathcal{P}} F(\mu)$$

Recall MMS

$$\mu^{k+1} \in \operatorname{arginf}_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{\tau} \mathcal{D}(\mu, \mu^k)$$

We have specified the important ingredients

$$\text{Energy} : F, \quad \text{Geometry} : \mathcal{D}$$

We can construct a concrete instance of MMS for gradient flow by “mix-and-match”.

Wasserstein-MMS: Jordan-Kinderlehrer-Otto (JKO) scheme

a.k.a. Minimizing Movement Scheme (MMS):

$$\mu^{k+1} \in \inf_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{2\tau} W_2^2(\mu, \mu^k)$$

This formulation is very general in the sense that it includes **nonlinear-in-measure** F .
We should think of this as the *gradient descent algorithm for prob. measures*.

Otto's Gradient flow equation in the Wasserstein space

$$\mu^{k+1} \in \inf_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{2\tau} W_2^2(\mu, \mu^k)$$

Continuous-time limit $\tau \rightarrow 0$, we have (non-trivially) the **gradient flow equation** (GFE)

$$\partial_t \mu - \nabla \cdot \left(\mu \nabla \frac{\delta F}{\delta \mu}[\mu] \right) = 0$$

which describes the dissipation of energy F in $(\text{Prob}(\bar{X}), W_2)$. [Otto et al 90s-2000s, Ambrosio 2005]

In a different flavor, we can write it just like ODE $\dot{x} = -\nabla f(x)$ (in the **rate** form; primal vs. dual force-balance)

$$\partial_t \mu = -\mathbb{K}_{\text{Otto}}(\mu) DF$$

Otto's Gradient flow equation in the Wasserstein space

$$\mu^{k+1} \in \inf_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{2\tau} W_2^2(\mu, \mu^k)$$

Continuous-time limit $\tau \rightarrow 0$, we have (non-trivially) the **gradient flow equation** (GFE)

$$\partial_t \mu - \nabla \cdot (\mu \nabla \frac{\delta F}{\delta \mu}[\mu]) = 0$$

which describes the dissipation of energy F in $(\text{Prob}(\bar{X}), W_2)$. [Otto et al 90s-2000s, Ambrosio 2005]

In a different flavor, we can write it just like ODE $\dot{x} = -\nabla f(x)$ (in the **rate** form; primal vs. dual force-balance)

$$\partial_t \mu = -\mathbb{K}_{\text{Otto}}(\mu) DF = \nabla \cdot (\mu \nabla DF).$$

Otto's Gradient flow equation in the Wasserstein space

$$\mu^{k+1} \in \inf_{\mu \in \mathcal{P}} F(\mu) + \frac{1}{2\tau} W_2^2(\mu, \mu^k)$$

Continuous-time limit $\tau \rightarrow 0$, we have (non-trivially) the **gradient flow equation** (GFE)

$$\partial_t \mu - \nabla \cdot (\mu \nabla \frac{\delta F}{\delta \mu}[\mu]) = 0$$

which describes the dissipation of energy F in $(\text{Prob}(\bar{X}), W_2)$. [Otto et al 90s-2000s, Ambrosio 2005]

In a different flavor, we can write it just like ODE $\dot{x} = -\nabla f(x)$ (in the **rate** form; primal vs. dual force-balance)

$$\partial_t \mu = -\mathbb{K}_{\text{Otto}}(\mu) DF = \nabla \cdot (\mu \nabla DF).$$

Example: WGF of (Boltzmann/KL/relative) Entropy

nonlinear (in measure) energy (e.g., in variational inference)

$$F(\mu) = D_{\text{KL}}(\mu \parallel \pi) = \int \log\left(\frac{\delta\mu}{\delta\pi}(x)\right) \rho(x) \, dx$$

Example: WGF of (Boltzmann/KL/relative) Entropy

nonlinear (in measure) energy (e.g., in variational inference)

$$F(\mu) = D_{\text{KL}}(\mu \parallel \pi) = \int \log\left(\frac{\delta\mu}{\delta\pi}(x)\right) \rho(x) \, dx$$

$$\frac{\delta F}{\delta\mu} [\mu] = \log \rho - \log \pi,$$

density $\rho := \frac{d\mu}{d\mathcal{L}}$ The Fokker-Planck equation as the **Wasserstein gradient flow** [Otto et al. 90s-2000s]

$$\begin{aligned} \partial_t \mu &= \nabla \cdot \left(\mu \nabla \frac{\delta F}{\delta\mu} [\mu] \right) \\ &= \nabla \cdot (\mu (\nabla \log \rho - \nabla \log \pi)) \\ &= \Delta \rho + \nabla \cdot (\rho \nabla \log \pi) \end{aligned}$$

Example: WGF of (Boltzmann/KL/relative) Entropy

nonlinear (in measure) energy (e.g., in variational inference)

$$F(\mu) = D_{\text{KL}}(\mu \| \pi) = \int \log\left(\frac{\delta\mu}{\delta\pi}(x)\right) \rho(x) \, dx$$

$$\frac{\delta F}{\delta\mu} [\mu] = \log \rho - \log \pi,$$

density $\rho := \frac{d\mu}{d\mathcal{L}}$ The Fokker-Planck equation as the **Wasserstein gradient flow** [Otto et al. 90s-2000s]

$$\begin{aligned} \partial_t \mu &= \nabla \cdot \left(\mu \nabla \frac{\delta F}{\delta\mu} [\mu] \right) \\ &= \nabla \cdot (\mu (\nabla \log \rho - \nabla \log \pi)) \\ &= \Delta \rho + \nabla \cdot (\rho \nabla \log \pi) \end{aligned}$$

► If π is the Lebesgue measure, we obtain the heat equation $\partial_t \mu = \Delta \mu$

Example: WGF of (Boltzmann/KL/relative) Entropy

nonlinear (in measure) energy (e.g., in variational inference)

$$F(\mu) = D_{\text{KL}}(\mu \parallel \pi) = \int \log\left(\frac{\delta\mu}{\delta\pi}(x)\right) \rho(x) \, dx$$

$$\frac{\delta F}{\delta\mu} [\mu] = \log \rho - \log \pi,$$

density $\rho := \frac{d\mu}{d\mathcal{L}}$ The Fokker-Planck equation as the **Wasserstein gradient flow** [Otto et al. 90s-2000s]

$$\begin{aligned}\partial_t \mu &= \nabla \cdot \left(\mu \nabla \frac{\delta F}{\delta\mu} [\mu] \right) \\ &= \nabla \cdot (\mu (\nabla \log \rho - \nabla \log \pi)) \\ &= \Delta \rho + \nabla \cdot (\rho \nabla \log \pi)\end{aligned}$$

- ▶ If π is the Lebesgue measure, we obtain the heat equation $\partial_t \mu = \Delta \mu$
- ▶ Note: The force field $\frac{\delta F}{\delta\mu} [\mu]$ and the “score” $\nabla \frac{\delta F}{\delta\mu} [\mu]$ are **not accessible** if μ is atomic. \implies “score-matching”...

Application: sampling and variational inference

Suppose $\pi \propto e^{-V(x)}$, but with unknown normalizing constant, we want

$$\inf_{\mu \in \mathcal{P}} D_{KL}(\mu \| \pi).$$

Using the WGF, we have the Fokker-Planck equation

$$\partial_t \mu = \nabla \cdot (\mu (\nabla \log \rho(x) - \nabla V(x)))$$

Suppose there is a single atom whose state is X_t (R.V.), it is pushed towards the velocity field

$$\nabla \log \rho(X_t) - \nabla V(X_t)$$

We can construct gradient descent

$$X_{t+1} = X_t + \tau \cdot (\nabla \log \rho(X_t) - \nabla V(X_t))$$

Langevin Monte-Carlo forward-Euler discretization

$$X_{t+1} = X_t - \tau \cdot \nabla V(X_t) + \sqrt{2\tau} Z, Z \sim N(0, \text{Id})$$

Application: (distributionally) robust learning with Otto's WGF

We can use our WGF theory (invented 20yr ago; nothing new) to solve Wasserstein DRO for robust learning (also adversarial robustness in [Sinha et al. 2017])

$$\min_{\theta} \sup_{\mu} \mathbb{E}_{\mu} l(\theta, x) - \gamma \cdot W_2^2(\mu, \hat{\mu}_N)$$

Application: (distributionally) robust learning with Otto's WGF

We can use our WGF theory (invented 20yr ago; nothing new) to solve Wasserstein DRO for robust learning (also adversarial robustness in [Sinha et al. 2017])

$$\min_{\theta} \sup_{\mu} \mathbb{E}_{\mu} l(\theta, x) - \gamma \cdot W_2^2(\mu, \hat{\mu}_N)$$

The inner measure-update step is gradient ascent

$$X_{t+1} = X_t + \tau \nabla l(\theta_t, X_t)$$

where $\tau = \frac{1}{2\gamma}$.

Application: (distributionally) robust learning with Otto's WGF

We can use our WGF theory (invented 20yr ago; nothing new) to solve Wasserstein DRO for robust learning (also adversarial robustness in [Sinha et al. 2017])

$$\min_{\theta} \sup_{\mu} \mathbb{E}_{\mu} l(\theta, x) - \gamma \cdot W_2^2(\mu, \hat{\mu}_N)$$

The inner measure-update step is gradient ascent

$$X_{t+1} = X_t + \tau \nabla l(\theta_t, X_t)$$

where $\tau = \frac{1}{2\gamma}$. Then the whole Wasserstein robust learning is simply gradient descent-ascent (GDA).

Energy dissipation balance of WGF

Recall the ODE case

$$\frac{d}{dt}f(x(t)) = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

Energy dissipation balance of WGF

Recall the ODE case

$$\frac{d}{dt}f(x(t)) = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

In $(\text{Prob}(\bar{X}), F, W_2)$, **Fenchel(-Young)** yields the **Energy dissipation balance** (equality) [Ambrosio et al. 2007]

$$\frac{d}{dt}F(\mu(t)) = -\frac{1}{2}|\mu'|_{W_2}(t)^2 - \frac{1}{2}|\nabla^- F|_{W_2}(\mu(t))^2$$

$$F(\mu(t)) - F(\mu(s)) = -\frac{1}{2} \int_s^t |\mu'|_{W_2}(r)^2 + |\nabla^- F|_{W_2}(\mu(r))^2 \, dr$$

Energy dissipation balance of WGF

Recall the ODE case

$$\frac{d}{dt}f(x(t)) = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

In $(\text{Prob}(\bar{X}), F, W_2)$, **Fenchel(-Young)** yields the **Energy dissipation balance** (equality) [Ambrosio et al. 2007]

$$\frac{d}{dt}F(\mu(t)) = -\frac{1}{2}|\mu'|_{W_2}(t)^2 - \frac{1}{2}|\nabla^- F|_{W_2}(\mu(t))^2$$

$$F(\mu(t)) - F(\mu(s)) = -\frac{1}{2} \int_s^t |\mu'|_{W_2}(r)^2 + |\nabla^- F|_{W_2}(\mu(r))^2 \, dr$$

- ▶ metric speed with velocity v_t : $|\mu'|_{W_2}(t) = \sqrt{\int |v_t|^2 \, d\mu}$
- ▶ metric slope: $|\nabla^- F|_{W_2}(\mu(t)) = \sqrt{\int |\nabla \frac{\delta F}{\delta \mu} [\mu](x)|^2 \, d\mu}$

Energy dissipation balance of WGF

Recall the ODE case

$$\frac{d}{dt}f(x(t)) = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

In $(\text{Prob}(\bar{X}), F, W_2)$, **Fenchel(-Young)** yields the **Energy dissipation balance** (equality) [Ambrosio et al. 2007]

$$\frac{d}{dt}F(\mu(t)) = -\frac{1}{2}|\mu'|_{W_2}(t)^2 - \frac{1}{2}|\nabla^- F|_{W_2}(\mu(t))^2$$

$$F(\mu(t)) - F(\mu(s)) = -\frac{1}{2} \int_s^t |\mu'|_{W_2}(r)^2 + |\nabla^- F|_{W_2}(\mu(r))^2 \, dr$$

- ▶ metric speed with velocity v_t : $|\mu'|_{W_2}(t) = \sqrt{\int |v_t|^2 \, d\mu}$
- ▶ metric slope: $|\nabla^- F|_{W_2}(\mu(t)) = \sqrt{\int |\nabla \frac{\delta F}{\delta \mu} [\mu](x)|^2 \, d\mu}$

The velocity field can be identified as $v_t = -\nabla \frac{\delta F}{\delta \mu} [\mu]$. EDB can then be used as the definition of gradient flows (curves of maximal slopes), even without GFE.

Energy dissipation balance of WGF

Recall the ODE case

$$\frac{d}{dt}f(x(t)) = -\left(\frac{1}{2}\|\dot{x}\|^2 + \frac{1}{2}\|\nabla f(x)\|^2\right)$$

In $(\text{Prob}(\bar{X}), F, W_2)$, **Fenchel(-Young)** yields the **Energy dissipation balance** (equality) [Ambrosio et al. 2007]

$$\frac{d}{dt}F(\mu(t)) = -\frac{1}{2}|\mu'|_{W_2}(t)^2 - \frac{1}{2}|\nabla^- F|_{W_2}(\mu(t))^2$$

$$F(\mu(t)) - F(\mu(s)) = -\frac{1}{2} \int_s^t |\mu'|_{W_2}(r)^2 + |\nabla^- F|_{W_2}(\mu(r))^2 \, dr$$

- ▶ metric speed with velocity v_t : $|\mu'|_{W_2}(t) = \sqrt{\int |v_t|^2 \, d\mu}$
- ▶ metric slope: $|\nabla^- F|_{W_2}(\mu(t)) = \sqrt{\int |\nabla \frac{\delta F}{\delta \mu} [\mu](x)|^2 \, d\mu}$

The velocity field can be identified as $v_t = -\nabla \frac{\delta F}{\delta \mu} [\mu]$. EDB can then be used as the definition of gradient flows (curves of maximal slopes), even without GFE.

For (Boltzmann) entropy $F(u) = \rho \log \rho$, EDB gives $\frac{d}{dt}F(\mu(t)) = -\int |\nabla \log \rho|^2 \rho \, dx$

Evolutionary variational inequality (EVI) $_{\lambda}$: Wasserstein GF

Under a few technical assumptions and the so-called λ -geodesic-convexity of the energy F , if along a geodesic curve γ ,

$$F(\gamma(s)) \leq (1-s)F(\gamma(0)) + sF(\gamma(1)) - \frac{\lambda}{2}s(1-s)W_2^2(\gamma(0), \gamma(1)), \quad \forall s \in [0, 1].$$

Evolutionary variational inequality (EVI) $_{\lambda}$: Wasserstein GF

Under a few technical assumptions and the so-called λ -geodesic-convexity of the energy F , if along a geodesic curve γ ,

$$F(\gamma(s)) \leq (1-s)F(\gamma(0)) + sF(\gamma(1)) - \frac{\lambda}{2}s(1-s)W_2^2(\gamma(0), \gamma(1)), \quad \forall s \in [0, 1].$$

Then, there exists unique gradient flow solution satisfies (EVI) $_{\lambda}$, for .

$$\frac{1}{2}W_2^2(\mu(t), \nu) \leq \frac{1}{2}e^{-\lambda(t-s)}W_2^2(\mu(s), \nu) + M_{\lambda}(t-s)(F(\nu) - F(\mu(t))),$$
$$\forall \nu \in \text{dom}(\mathcal{F}), M_{\lambda}(\tau) = \int_0^{\tau} e^{-\lambda(\tau-s)} \, ds.$$

Set $\nu \in \operatorname{arginf}_{\mu} F(\mu)$, we have exponential convergence in-time and uniqueness of gradient flow.

Thank you!

There are many other active research topics in GF for ML

Thank you!

There are many other active research topics in GF for ML

- ▶ Gradient flow structure with *kernel geometry* [also some of my past / current works]

Thank you!

There are many other active research topics in GF for ML

- ▶ Gradient flow structure with *kernel geometry* [also some of my past / current works]
- ▶ Unbalanced transport and its gradient flow

Thank you!

There are many other active research topics in GF for ML

- ▶ Gradient flow structure with *kernel geometry* [also some of my past / current works]
- ▶ Unbalanced transport and its gradient flow
- ▶ Applications: causal inference, mean-field NN, Nash equilibrium, offline RL, policy optimization. . .

Thank you!

There are many other active research topics in GF for ML

- ▶ Gradient flow structure with *kernel geometry* [also some of my past / current works]
- ▶ Unbalanced transport and its gradient flow
- ▶ Applications: causal inference, mean-field NN, Nash equilibrium, offline RL, policy optimization. . .

Reference

1. Mielke, A. An introduction to the analysis of gradients systems. Preprint at <http://arxiv.org/abs/2306.05026> (2023).
2. Otto, F. The geometry of dissipative evolution equations: the porous medium equation. (2001).
3. Ambrosio, L. & Savaré, G. Gradient Flows of Probability Measures. in Handbook of Differential Equations: Evolutionary Equations vol. 3 1–136 (Elsevier, 2007).
4. Santambrogio, F. Optimal transport for applied mathematicians. Birkhäuser, NY 55, 94 (2015).
5. Peletier, M. A. Variational Modelling: Energies, gradient flows, and large deviations. arXiv:1402.1990 [math-ph] (2014).