

We start with the given equation:

$$\text{ck}(k, t_0, T) = \frac{2T^2 \sin\left(\frac{\pi kt_0}{2T}\right)}{k\pi(4T^2 - k^2 t_0^2)}$$

**Step 1: Factor the denominator.**

We factor the denominator as follows:

$$\text{ck}(k, t_0, T) = \frac{2T^2 \sin\left(\frac{\pi kt_0}{2T}\right)}{k\pi \left(2T - \frac{kt_0}{2}\right) \left(2T + \frac{kt_0}{2}\right)}$$

**Step 2: Use the definition of sinc.**

The sinc function is defined as:

$$\text{sinc}(x) = \frac{\sin(\pi x)}{\pi x}$$

Using this definition, we express the sine term in the numerator in terms of a sinc function:

$$\sin\left(\frac{\pi kt_0}{2T}\right) = \frac{kt_0}{2T} \cdot \pi \cdot \text{sinc}\left(\frac{kt_0}{2T}\right)$$

**Step 3: Substituting into the equation.**

Now, substitute the sinc form into the original equation:

$$\text{ck}(k, t_0, T) = \frac{2T^2 \cdot \frac{kt_0}{2T} \cdot \pi \cdot \text{sinc}\left(\frac{kt_0}{2T}\right)}{k\pi(4T^2 - k^2 t_0^2)}$$

**Step 4: Simplify the equation.**

First, cancel out the common factors of  $\pi$  and  $k$ :

$$\text{ck}(k, t_0, T) = \frac{2T^2 \cdot \frac{t_0}{2T} \cdot \text{sinc}\left(\frac{kt_0}{2T}\right)}{4T^2 - k^2 t_0^2}$$

Now, simplify the terms:

$$\text{ck}(k, t_0, T) = \frac{t_0}{4T} \cdot \frac{2T^2 \cdot \text{sinc}\left(\frac{kt_0}{2T}\right)}{4T^2 - k^2 t_0^2}$$

**Step 5: Expand the final expression.**

Now we split the term inside the bracket:

$$\text{ck}(k, t_0, T) = \frac{t_0}{4T} \cdot \text{sinc}\left(\frac{kt_0}{2T}\right) \left(1 - \frac{k^2 t_0^2}{4T^2}\right)$$

**Step 6: Final expression:**

$$\text{ck}(k, t_0, T) = \frac{t_0}{4T} \cdot \text{sinc}\left(\frac{kt_0}{2T}\right) - \frac{t_0}{4T} \cdot \frac{k^2 t_0^2 \cdot \text{sinc}\left(\frac{kt_0}{2T}\right)}{4T^2}$$