




GENERAL:

- The brain accounts for just 2% of our body weight but receives 15% of resting cardiac output and consumes 20% of the oxygen produced by the cardiovascular system
- The brain depends primarily on oxidative metabolism for generation of ATP
- The brain uses glucose and has a minimal capacity to use glycolysis or other energy substrates
- A blockage or leak in an artery or arteriole will cause an interruption of blood flow and damage the area of the brain supplied by that vessel
- 2 million nerve cells are lost each minute in which reperfusion has not been achieved
- Minor occlusions may occur in the intracranial or extracranial vessels without causing neurological signs and symptoms due to extensive collateral connections exist between the arterial systems of the brain
- Size and distribution of an infarct depends on the speed of thrombosis and the presence of collateral circulation
- Drop in systemic blood pressure may lead to an infarct even without complete vascular occlusion in severe atherosclerosis

DEFINITIONS

Tissue based definitions according to consensus statements from the American Heart Association and American Stroke Association (AHA/ASA)

TRANSIENT ISCHEMIC ATTACK (TIA):

Transient episode of neurologic dysfunction caused by focal brain, spinal cord, or retinal ischemia, **without** acute infarction.

CENTRAL NERVOUS SYSTEM (CNS) INFARCTION:

Brain, spinal cord, or retinal cell death attributable to ischemia, based on neuropathologic, neuroimaging, and/or clinical evidence of permanent injury.

ISCHEMIC STROKE:


An episode of neurologic dysfunction caused by focal cerebral, spinal, or retinal infarction.

SILENT CNS INFARCTION:


Imaging or neuropathological evidence of CNS infarction without a history of acute neurological dysfunction

TIA

- **Self-limited** episodes of vascular obstruction by atheromatous emboli and/or platelet and fibrin aggregates
- Lasts from a few seconds to minutes and usually disappear within an hour
- If symptoms last longer (up to 24 hours), higher probability of infarction
- **Prelude to a stroke:** One third of patients with TIAs develop clinically significant infarcts within 5 years
- **Most frequent symptoms:**
 - Temporary loss of vision (amaurosis fugax),
 - Difficulty speaking (aphasia)
 - Numbness or tingling (paresthesia), usually on one side of the body
 - Weakness on one side of the body (hemiparesis)
- **Less common symptoms:**
 - Impairment of consciousness
 - Dizziness, lack of coordination or poor balance
 - Fatigue
 - Feeling of “something is wrong with me”




Geographic Patterns: Stroke Death Rates for 2018 through 2020 for Adults Aged 35+ years by County. The map shows that concentrations of counties with the highest stroke death rates – meaning the top quintile – are located primarily in the Southeast, with heavy concentrations of high-rate counties in Arkansas, Alabama, Louisiana, Mississippi, Tennessee, Texas, Kentucky, Michigan, Ohio, North Carolina, South Carolina, Guam, and the Northern Mariana Islands. Pockets of high-rate counties are also found in Delaware, Maryland, Oklahoma, Nevada, Oregon, Kansas, California, and Alaska.




Geographic Patterns: Stroke Death Rates for 2018 through 2020 for Adults Aged 65 Years and Older by County. The map shows that concentrations of counties with the highest stroke death rates – meaning the top quintile – are located primarily in the Southeast, with heavy concentrations of high-rate counties in Arkansas, Alabama, Louisiana, Mississippi, Tennessee, Texas, Kentucky, Michigan, Ohio, North Carolina, South Carolina, Guam, and the Northern Mariana Islands. Pockets of high-rate counties are also found in Delaware, Maryland, Oklahoma, Nevada, Oregon, Kansas, California, and Alaska.


AHA: HEART DISEASE AND STROKE STATISTICS-2022 UPDATE



AHA: HEART DISEASE AND STROKE STATISTICS-2023 UPDATE



Estimated 10-year stroke risk in adults 55 years of age according to levels of various risk factors (Framingham Heart Study).




	A	B	C	D	E	F
Blood Pressure*	95-105	138-148	138-148	138-148	138-148	138-148
Diabetes	No	No	Yes	Yes	Yes	Yes
Cigarette Smoking	No	No	No	Yes	Yes	Yes
Prior AF	No	No	No	No	Yes	Yes
Prior CVD	No	No	No	No	No	Yes

* - Closest ranges for women are : 95-104 and 115-124.




Mozaffarian D et al. *Circulation*. 2015;131:e29-e322
Copyright © American Heart Association, Inc. All rights reserved.



INFARCT

ETIOLOGY AND PATHOGENESIS:

- Caused by local interruption of blood flow.
- Most common form of cerebrovascular disease
- Constitutes 70-80% of cerebrovascular accidents.
- Most common in patients >60 years of age.
- More common in males than females (until the age of 65).
- Most common cause: atherosclerosis and plaque rupture
- Most common predisposing factors:
 - Hypertension
 - diabetes mellitus
 - smoking



Genetic and inflammatory mechanisms in stroke. Gray background = subendothelium; white background = blood vessel lumen; SMC = intraluminal smooth muscle cells.

CAUSES OF ISCHEMIC INFARCTION

- Atherosclerosis-atherothrombosis
- Small vessel disease
- Cardioembolism, including paradoxical embolism through a patent foramen ovale
- Vasculitis
- Arterial dissection
- Polycythemia, thrombocythemia, thrombotic thrombocytopenic purpura (TTP)
- Systemic lupus erythematosus (SLE)
- Sickle cell disease
- Vascular spasm
- Meningitis and fungal vasculitis
- Hypercoagulability (Factor V Leiden, Prothrombin 20210A mutation, antiphospholipid antibody syndrome)
- Inherited metabolic disorders (Fabry disease, homocystinuria, mitochondrial disorders)
- Fibromuscular dysplasia and other angiopathies

This is for future reference. Lysosomal storage diseases pop up in the boards from time to time.

Fabry disease:

- Lysosomal storage disease (sphingolipidosis)
- Caused by the lack of alpha-Galactosidase A (alpha-GAL), which hydrolyses the terminal α -galactosyl moieties of glycolipids and glycoproteins
- Lack of alpha-GAL leads to the accumulation of a glycolipid, globotriaosylceramide (GL3), and its metabolites within several types of cells, including endothelial, cardiac and nerve cells
- Dysfunction of multiple systems is characteristic: progressive heart and kidney damage, stroke, acroparesthesias, angiokeratomas, hypohidrosis, corneal opacities, GI-problems (GI-dysmotility due to neuropathic and myopathic changes)
- X-linked, but unlike other x-linked diseases, females carrying the faulty gene may also develop cardiac and kidney problems.
- Frequency: 1 in 3400-4000 newborns; the second most common lysosomal storage disease (Gaucher disease with a defective beta-Glucosidase is the most common)

INFARCT

THROMBOTIC OCCLUSIONS:


Most common sites:

- Internal carotid arteries at the carotid bifurcation
- Middle cerebral arteries at the origin
- Basilar artery at both ends


EMBOLIC OCCLUSIONS:

- Most common cause of an intracranial arterial occlusion
- Thrombo-emboli and emboli of atheromatous material most commonly originate in:
 - Heart
 - Carotids
 - More proximal arterial segment
- Less common emboli:
 - Vegetations from endocarditis
 - Fat
 - Bone marrow
 - Air

CIRCLE OF WILLIS




STENOSIS OF INTERNAL CAROTID ARTERY




HIGH RESOLUTION CONTRAST
ENHANCED MAGNETIC RESONANCE
ANGIOGRAPHY (MRA)


- A STENOSIS OF INTERNAL CAROTID ARTERY
- B ALTERED BLOOD FLOW DUE TO STENOSIS
- C NORMAL BLOOD FLOW



Brain. Severe atherosclerosis in internal carotid arteries, basilar and vertebral arteries and in the circle of Willis.




Left: Severe atherosclerosis in cerebral arteries. **Right:** Severe atherosclerosis and thrombosis of the basilar artery.



Diagnosis: Stent in the internal carotid artery. **Description:** Atherosclerosis of carotid artery with a stent in the internal carotid artery. **Clinical findings:** Hypertension, nicotine abuse, generalized atherosclerosis. Asymptomatic stenosis of the left internal carotid artery. Percutaneous transluminal angioplasty and stenting. Two hours after the operation patient became confused, disoriented, and dysphasic and died. An acute cerebral infarct due to collapse of the stent was found.

INFARCT

DISSECTING ANEURYSM IN CAROTID ARTERY




- Dissection of the carotid and vertebral arteries causes arterial occlusion and embolism
- Accounts for up to 20% of strokes in younger patients


19

Dissecting aneurysm in carotid artery of a young man. Blood has dissected between the internal elastic lamina and the media and is obstructing blood flow to the brain. Elastic van Gieson stain


SOURCES OF CARDIOEMBOLIC STROKE



Fibrillation




Left: Complication of MI: mural thrombus attached to damaged ventricular wall. Old infarct appears white. **Right:** Here is a thromboembolus that originated from mural thrombus in the left atrium. The heart is a common source for such emboli.



Diagnosis: infectious endocarditis with septic emboli to the brain.

FAT EMBOLISM SYNDROME



EXTENSIVE WHITE MATTER PETECHIAL HEMORRHAGES

Complication of long bone fractures and traumatic lesions of viscera and soft tissues.

40-60% of patients with total joint arthroplasty have fat embolism, but only 1-5% develop the syndrome.


Neurologic signs and symptoms typically present 24-72 hrs. after the initial trauma but may start up to a week later.

Fat globules can be demonstrated in lung post-capillary venules indicating that ASD or VSD (allowing paradoxical embolization) are not necessary.

Non-traumatic settings:

- osteomyelitis
- pancreatitis
- diabetes

The extensive white matter petechial hemorrhages seen here are typical for fat embolism syndrome. The patient manifests generalized cerebral dysfunction with disturbances of higher cortical function and consciousness, often without localizing signs, usually about a week after the initiating event, such as long bone fractures in a vehicular accident.



Brain. Diagnosis: Air embolism. **Description:** Numerous intraluminal air bubbles are visible. **Comment:** Bubbles in the arterial circulation can arise from basically three sources: venous gas embolism with breach of the pulmonary vascular filter (paradoxical gas embolism), patent foramen ovale (paradoxical gas embolism) and tear of the pulmonary parenchyma with entry of gas into pulmonary venous outflow. Studies show that systemic venous bubbles are trapped in the pulmonary arterial tree and are usually completely eliminated from that site. The other main mechanism for arterial gas embolism is by way of the pulmonary overpressure syndrome or 'burst lung'. This occurs from baropressure increases as the diver on compressed air ascends with a closed glottis or a free diver takes a breath of compressed air at depth and ascends. Because of Boyle's law, maximal changes in volume occur in the 4 feet (1.22 m.) closest to the surface and the diver sustains a tear in the pulmonary parenchyma with the escape of air into the pulmonary venous outflow.

LESS COMMON CAUSES OF INFARCTS

INFECTIOUS VASCULITIS

- Syphilis and TB used to be common causes
- Now more common in the setting of immunosuppression and opportunistic infection
 - Aspergillosis and mucor most common
 - CMV encephalitis

COLLAGEN VASCULAR DISEASE

- SLE, RA, Bechet


NON-INFECTIOUS SYSTEMIC VASCULITIS WITH CNS INVOLVEMENT

- **Giant cell (temporal) arteritis most common**
- Polyarteritis nodosa, granulomatosis with polyangiitis, Takayasu arteritis, Kawasaki disease – all are rare in the CNS

PRIMARY GRANULOMATOUS ANGIITIS OF THE CNS


- Involves multiple small- to medium-sized parenchymal and subarachnoid vessels and causes destruction of the vessel walls
- Clinically diffuse encephalopathy or multifocal manifestations and cognitive dysfunction
- Patients improve with steroids and immunosuppressive therapy

THROMBOSIS CAUSED BY ANGIOINVASIVE FUNGI



SILVER STAIN

ANGIOINVASIVE ASPERGILLOSIS




INFARCT
IN PONS

BASILAR ARTERY THROMBOSIS CAUSED
BY MUCORMYCOSIS IN A LEUKEMIA
PATIENT

26


Left: Angioinvasive aspergillosis. **Right:** Basilar artery thrombosis caused by mucormycosis in a leukemia patient. The pons is infarcted and has green discoloration, because the patient was in liver failure and had high serum bilirubin.

GIANT-CELL (TEMPORAL) ARTERITIS



Kumar: Robbins and Cotran Pathologic Basis of Disease, Professional Edition , 8th ed.

FIGURE 11-23 Giant-cell (temporal) arteritis. **A**, H&E stain of section of temporal artery showing giant cells at the degenerated internal elastic lamina in active arteritis (*arrow*). **B**, Elastic tissue stain demonstrating focal destruction of internal elastic lamina (*arrow*) and intimal thickening (IT) characteristic of long-standing or healed arteritis. **C**, Examination of the temporal artery of a patient with giant-cell arteritis shows a thickened, nodular, and tender segment of a vessel on the surface of head (*arrow*). (**C** from Salvarani C et al.: *Polymyalgia rheumatica and giant-cell arteritis*. *N Engl J Med* 347:261, 2002.)




Left: Giant cell arteritis. **Right:** Swollen optic disc caused by ischemia due to occlusion of the ciliary feeder vessels of the optic nerve. The patient experiences sudden, painless, usually devastating, permanent loss of vision in one eye. Most individuals are aged over 70 years and have a combination of limb girdle myalgias, fever and lassitude ("polymyalgia rheumatica"), caused by an idiopathic autoimmune disorder of the elderly. Some patients have scalp tenderness and pain with chewing, which may be manifestations of external carotid artery ischemia.


Laboratory findings: elevated sedimentation rate and C-reactive protein.

Biopsy of a temporal artery shows granulomatous inflammation.

The diagnosis must be made immediately because there is a strong chance that the optic nerve of the contralateral eye will become involved within days unless high-dose corticosteroid treatment is begun. The treatment is nearly 100% effective in preventing second eye involvement.

THE CONSEQUENCES OF BLOOD CLOTS DEPEND ON THE LOCATION





ANTERIOR CIRCULATION ISCHEMIA

- **The most common artery involved in stroke is the middle cerebral artery (MCA)** which supplies blood to the lateral surface of the frontal, temporal and parietal lobes
- Deep penetrating branches of the MCA perfuse the subcortical regions of the brain including the basal ganglia, internal capsule and the corona radiata
- Damage to the MCA causes:
 - Contralateral weakness and sensory loss in the face, neck and arm and to a lesser degree in the leg
 - Aphasia, if the damage is in the left (dominant) hemisphere

ANTERIOR CIRCULATION ISCHEMIA

- **The second most common artery damaged in stroke is the anterior cerebral artery (ACA)** which perfuses the anterior surface of the frontal lobe and the medial surfaces of the frontal and parietal lobes.
- **Damage to the ACA causes:**
 - Contralateral weakness
 - Sensory loss primarily in the leg
- **Damage to the opthalmic artery causes:**
 - Ipsilateral monocular loss of vision
 - Homonymous hemianopsia
 - Amaurosis fugax (fleeting blindness)

POSTERIOR CIRCULATION ISCHEMIA

Interruption of blood supply to the brainstem leads to :

- loss of brainstem function
- cranial nerve abnormalities with or without hemiparesis
- hemisensory deficits

Brainstem damage may lead to:

- dysphagia (difficulty swallowing due to paralysis of tongue and larynx)
- dysarthria (inability to articulate clearly due to facial paralysis)

Loss of blood flow to the cerebellum causes:

- ataxia (gross motor incoordination due to cerebellar damage)
- hypotonia
- dysmetria
- loss of equilibrium
- vertigo

Loss of blood flow in the posterior cerebral artery causes :

- diplopia
- bilateral hemianopsia

INFARCT

MORPHOLOGY:

First 4-12 hrs:

No gross or microscopic change

By 12-18 hrs:

Pink neurons

Neuronal changes similar to global ischemia, but accompanied by PMNs

By 36-48 hrs:

Swelling and softening of parenchyma and blurring of the demarcation between gray and white matter

Hemorrhages

By 72 hrs:

Phagocytosis of the dead cells results in sharper demarcation of the infarct, peak macrophages at 7-14 days

By 1 month:


Extensive phagocytosis results in further softening and liquefaction and development of irregular cavities

By 6 months:

Complete liquefaction

A thin rim of subpial parenchyma supplied by leptomeningeal arteries is preserved over a cortical infarct (helps to differentiate an old infarct from an old contusion).


MASSIVE ACUTE INFARCT IN THE MCA TERRITORY




- CEREBRAL EDEMA CAUSED A MIDLINE SHIFT
- BLURRING OF THE GRAY-WHITE MATTER DEMARCATION
- SECONDARY BLEEDING AT THE EDGES OF THE INFARCT

Brain, gross, coronal section. Massive acute infarct, middle cerebral artery territory.


FRESH ANEMIC BRAIN INFARCT



- THE NORMAL SHARP DEMARCATION BETWEEN THE GRAY AND WHITE MATTER IS LOST
- FOCAL HEMORRHAGES



NORMAL




ACUTE INFARCT


Left: 87 y o woman. Brain, parietal lobe. **Diagnosis:** Fresh anemic brain infarct.

Right top: Normal. **Right bottom:**

**INFARCT IN THE ACA-MCA WATERSHED AREA
WITH SECONDARY HEMORRHAGE**




Brain, Gross, coronal section.



- A. Fresh anemic brain infarction in the parietal lobe (MCA territory) of an 87-yo woman. Note hyper-eosinophilic dead neurons and edematous neuropil.
- B. Different patient: Successful clot retrieved from the basilar artery using the Merci retriever.
- C. Partially lysed thromboembolus after administration of a thrombolytic agent.
 - In the treatment of acute cerebral ischemia by thrombolytic agents, time lost is brain lost.
 - In a typical middle cerebral artery ischemic stroke, 2 million nerve cells are lost each minute in which reperfusion has not been achieved.
 - In several large clinical trials treatment within the first 90 minutes of onset increased the odds of an excellent outcome by 2.6-fold, in the 91- to 180-minute window by 1.6-fold, and in the 181- to 270-minute window by 1.3-fold, while treatment in the 271- to 360-minute window did not improve outcome in a statistically significant manner.
- D. Complication of thrombolytic therapy: secondary hemorrhage.


SUBACUTE BRAIN INFARCT



87 y o woman. Brain, parietal lobe. **Diagnosis:** Subacute brain infarct.

Left: Description: Edema and neovascularisation.

Right: Description: Organized thrombus with recanalization.



78 y o woman. Parietal lobe. **Diagnosis:** Old anemic infarct. **Description:** Brown discoloration in the frontoparietal area of the brain.


OLD ANEMIC INFARCT




CYSTIC LESION IN THE MCA TERRITORY
18 MONTHS AFTER THE PATIENT SUFFERED A STROKE

Brain, gross, whole brain.


History: A 68r-old right-handed woman awoke one morning aphasic and unable to move her right arm and leg. The patient died 18 months after the onset of the CNS symptoms.




Acc305-77 439.svs, Brain, HE, 0.3X




Acc305-77 439.svs, Brain, HE, 5X



80 y o man. Basal ganglia. **Diagnosis:** Remote infarct.





78 yom. Diagnosis: Atrophy of the motor tracts after medial infarction.

Description: Cross section through the mesencephalon shows a marked atrophy of the left motor tract.

Additional findings: Old medial infarction of the left hemisphere 10 years ago. **Clinical findings:** Cerebrovascular insult with right hemiparesis.


Comment: Wallerian degeneration is defined as anterograde degeneration of a nerve fiber following damage to nerve cell body or disruption of the proximal axon. It occurs both in peripheral and central nervous system. The breakdown of myelin in CNS is slower, because macrophage recruitment takes longer. **The main cause of Wallerian degeneration in the CNS is an infarct, less often tumor, hemorrhage, or vascular malformation. The most common location is the pyramidal tract.** In the atrophic stage months or years after the injury there is a conspicuous decrease of the volume of the pyramidal tract after a medial infarction.

LACUNAR INFARCTS

- **Small, slit-shaped, <1.5 cm in diameter**
- **Usually multiple**
- Located at sites supplied by perforating branches:
 - Basal ganglia
 - Internal capsule
 - Thalamus
 - Pons
- **Arteriolosclerosis**
 - Small vessel disease associated with **chronic hypertension** is the most common cause
 - Endothelial injury and leakage of plasma proteins from vessels cause development of typical hyaline arteriolosclerosis
- Lacunar infarcts may be asymptomatic or cause focal neurologic deficits
- Multiple lacunar infarcts are associated with ischemic white matter degeneration and s.c. **vascular dementia** (Binswanger disease)
- Multiple lacunar infarcts in basal ganglia cause a syndrome of rigidity and abnormal gait resembling Parkinson disease



MULTIPLE LACUNAR INFARCTS WITH WHITE MATTER ATROPHY; AN OLD INFARCT IN THE MCA TERRITORY



Brain– Gross, coronal section. Multiple lacunar infarcts are associated with ischemic white matter degeneration and dementia: **Binswanger disease.**


SMALL VESSEL DISEASE - OTHER EXAMPLES

- **Diabetic microangiopathy**
 - Probably caused by glycation of proteins, which leads to diffuse basement membrane thickening and endothelial cell damage
- **Cerebral autosomal dominant arteriopathy with subcortical infarcts and ischemic leukoencephalopathy (CADASIL)**
 - Caused by mutations of the *notch3* gene in chromosome 19
 - The problem is degeneration of vascular smooth muscle cells
 - Migraine headache and TIAs beginning at ca 30 years of age with detectable tissue changes in white matter by imaging studies
 - Recurrent strokes and dementia
- **Genetic angiopathy due to mutations of collagen 4A1**
 - Collagen 4A1 is a constituent of the vascular adventitia
 - Causes porencephaly (CNS cysts and cavities) in infants – rare
 - Causes lacunar infarcts and leukoencephalopathy in adults

SMALL VESSEL DISEASE IN DIABETES



BILATERAL LACUNAR INFARCTS IN BASAL GANGLIA
IN A HYPERTENSIVE DM PATIENT



HYALINE ARTERIOLOSCLEROSIS
HEMOSIDERIN-FILLED MACROPHAGES
INDICATE PREVIOUS BLEEDING

Left: Bilateral lacunar infarcts in the basal ganglia of a diabetic patient. Lacunar infarcts are caused are associated with small vessel disease. **Right:** Hyaline arteriosclerosis in a diabetic patient.

VENOUS INFARCTS

The least common pattern of cerebral infarction

Caused by occlusion of the venous sinuses and cerebral cortical veins by local thrombosis

Predisposing factors: disorders that cause hypercoagulability

- Oral contraceptives
 - 8 per 100,000 users
- Pregnancy
 - 11.6 per 100,000 deliveries in developed countries
 - 450 per 100,000 deliveries in developing countries
 - Highest risk during the third trimester and in the postpartum period; highest within the 6 weeks after delivery
- Polycythemia
- Systemic lupus
- Spread of infection from adjacent foci
 - nasal sinuses, middle ear, sometimes skin
- Dehydration in children

INTRACRANIAL HEMORRHAGE

CLASSIFICATION:

Based on the partition of the intracranial space into four anatomic compartments:

- Brain parenchyma
- Subarachnoid space
- Subdural space
- Epidural space

ETIOLOGY AND PATHOGENESIS:

- **Hypertension (Charcot-Bouchard microaneurysms)**
- **Berry aneurysm**
- **Trauma**
- Vascular malformations (AVM)
- Primary and metastatic neoplasms
- Bleeding associated with hematologic malignancies
- Coagulation disorders (anticoagulation therapy)
- Cerebral amyloid angiopathy (CAA)
- Vasculitis (polyarteritis nodosa, SLE, bacterial and fungal arteritis)


INTRACRANIAL HEMORRHAGE

INTRAPARENCHYMAL HEMORRHAGE:

- Lesions are predominantly in three locations:
 - 65% in the basal ganglia and thalamus
 - 15% in the pons
 - 10% in the cerebellum
- Minute hemorrhages (petechiae) are less harmful than the obstruction of the associated small vessels
- Large hematomas act as space occupying lesions and have the potential to produce focal neurologic deficits and to cause lethal transtentorial herniation

HEMORRHAGE IN THALAMUS

RISK FACTOR: HYPERTENSION



- Hypertensive hemorrhage in the right thalamus extends into the ventricles.
- Hemorrhages in this location are not amenable to surgical intervention.



62 yom with a 3-week-old rust-colored hemorrhage in thalamus.


53

Left: A CT scan demonstrating a hypertensive hemorrhage in the right thalamus that has extended into the ventricular system. Hemorrhages in this location are not amenable to surgical intervention with removal of the blood.

Right: 62 y o man. Thalamus. **Diagnosis:** Old intracerebral hemorrhage

FRESH MASSIVE HEMORRHAGE IN BASAL GANGLIA


RISK FACTOR: HYPERTENSION



Diagnosis: Fresh massive hemorrhage in basal ganglia (typical location) in a hypertensive patient. Hemorrhages involving the basal ganglia area (the putamen in particular) tend to be non-traumatic and caused by hypertension, which damages and weakens the small penetrating arteries. A mass effect with midline shift, often with secondary edema, may lead to herniation.


MASSIVE INTRACEREBRAL HEMORRHAGE

RISK FACTOR:
ETOH CIRRHOSIS WITH COAGULOPATHY



A

RISK FACTOR:
CHRONIC MYELOID LEUKEMIA; BLAST CRISIS



B



- A. 65 y o man. **Diagnosis:** Intracerebral hemorrhage penetrating the ventricular system. **Description:** Hemorrhage originating from the left basal ganglia. **Clinical findings:** Alcoholic liver disease/cirrhosis.
- B. 59 y o man with CML in blast crisis. Cerebellum. **Diagnosis:** Fresh hemorrhage in cerebellum.

SMALL VESSEL DISEASE ASSOCIATED WITH INTRA-CEREBRAL HEMORRHAGE

CEREBRAL AMYLOID ANGIOPATHY (CAA)

- Dense uniform deposits of various types of amyloid in the vessel walls (mostly A β)
- Small vessels become rigid tubes; no fibrosis
- Affects primarily leptomeningeal and cerebral cortical arterioles and capillaries
- Molecular layer of cerebellum may be involved
- More often associated with brain hemorrhages than lacunar infarcts
- May cause both microbleeds and large hemorrhages
- Lobar distribution in contrast with HTN-associated bleeds, which occur more often in thalamus and basal ganglia
- Most cases are sporadic, but a few are familial
 - The majority of Alzheimer patients have also CAA
 - Autosomal dominant Alzheimer disease caused by trisomy 21
 - Mutations in genes coding for the Amyloid Precursor Protein and Presenilin

CEREBRAL AMYLOID ANGIOPATHY (CAA)




- CAA-related hemorrhages can occur anywhere but are most commonly located in the frontal and parietal lobes and involve the cortex and the subcortical white matter.
- A spontaneous cerebral hemorrhage in an elderly person without an apparent cause should raise a suspicion for CAA.
- Amyloid deposition may incite granulomatous angiitis.

57

Left: Massive cerebral hemorrhage in a patient with cerebral amyloid angiopathy (CAA). **Right:** Immunostain for beta-amyloid.


RECAP: ISCHEMIC VS. HEMORRHAGIC STROKE

Ischemic stroke



A clot blocks blood flow to an area of the brain

Hemorrhagic stroke



Bleeding occurs inside or around brain tissue

SUBARACHNOID HEMORRHAGE

Most common causes:

- Rupture of a berry aneurysm
- Trauma

Less common cause:

- Arteriovenous malformation

Clinical signs:

- Headache (increasing intracranial pressure)
- Nuchal rigidity (meningeal irritation)
- Alteration in mental status

Complications:

- Hydrocephalus resulting from organization of the blood in the CSF
 - **Communicating:** CSF flows through the ventricles, but is not adequately absorbed into the bloodstream.
 - **Non-communicating:** The aqueduct and/or the foramina are closed and the flow of CSF is impaired.

SUBARACHNOID HEMORRHAGE



Organization of the blood may result in a blockage of the foramina or cerebral aqueduct and disruption of CSF circulation or meningeal fibrosis and disruption of absorption of the CSF.

Brain. Gross, right lateral view. The convexity of the brain is covered by blood, which may organize resulting in meningeal fibrosis, if the patient survives.

ANEURYSMS

BERRY (SACCULAR) ANEURYSM

- Originates in a structurally weak area at the branching point of a large cerebral artery in or close to the circle of Willis
- 90% occur within the carotid supply
- 20-30% are multiple
- Rupture of a berry aneurysm is the most common cause of symptomatic subarachnoid hemorrhage in adults


ATHEROSCLEROTIC ANEURYSM

- Fusiform
- Typical locations:
 - internal carotid artery
 - vertebral artery
 - basilar artery

MYCOTIC (SEPTIC) ANEURYSM

- Most commonly associated with subacute bacterial endocarditis
- Due to weakening of the walls of usually small arteries by infected emboli
- May cause hemorrhages anywhere in the brain
- Infection may spread into the leptomeninges (leptomeningitis) or parenchyma resulting in an abscess

COMMON SITES OF BERRY ANEURYSMS



62


Robbins 7th Ed. Figure 28-18 Common sites of saccular (berry) aneurysms in the circle of Willis.

RUPTURED BERRY ANEURYSMS CAUSE SUBARACHNOID HEMORRHAGE

MULTIPLE BERRY ANEURYSMS




CIRCLE OF WILLIS



63

Left: Circle of Willis with three berry aneurysms. Multiple aneurysms are seen in about 20-30% of cases of berry aneurysm. Such aneurysms are "congenital" in the sense that the defect in the arterial wall is present from birth, but the actual aneurysm takes years to develop, so that rupture is most likely to occur in young to middle-aged adults.

Right: The subarachnoid hemorrhage from a ruptured aneurysm is more of an irritant producing vasospasm than a mass lesion.




Left: 66 yom. Base of brain. **Diagnosis:** Non-ruptured berry (=saccular = spherical) aneurysm with thrombosis.

Right: Clipped berry aneurysm.


LARGE SACCULAR ANEURYSM

- A SPACE-OCCUPYING LESION
- MAY MIMIC TUMOR CLINICALLY



A large saccular aneurysm at the cerebello-pontine angle may clinically present with the same symptoms as a tumor.

WHAT IS THE MOST LIKELY ETIOLOGY FOR THIS ANEURYSM?



A fusiform aneurysm in the basilar artery is typically caused by atherosclerosis.

ARTERIO-VENOUS MALFORMATION (AVM)

ETIOLOGY:

- Congenital

DISTRIBUTION:


- Cerebral hemispheres within the distribution of middle cerebral artery
most common location

PATHOPHYSIOLOGY:

- Low-resistance AV shunt
- AVM siphons blood from the adjacent parenchyma
- Rupture may cause an intraparenchymal and/or subarachnoid hemorrhage

CLINICAL FEATURES:


- Seizures
- Neurologic deficits
- Chronic mass effects



A cause for hemorrhage, particularly in persons aged 10 to 30, is a vascular malformation. **Left:** This angiogram demonstrates a tortuous collection of irregular small vessels in the temporal region of the brain. **Right:** Seen here is a mass of irregular, tortuous vessels over the left posterior parietal region.

ARTERIOVENOUS MALFORMATION

- A. AVM in the MCA territory:
- Most common location: convexity of the brain
 - Surgical treatment may be possible




- B. An AVM deep in the brain:
- Difficult to treat surgically
 - Rupture results in an intracerebral hemorrhage, which may extend into the ventricles and subarachnoid space

69

A. Brain, gross, coronal section: Diagnosis: A large Arteriovenous malformation in the MCA territory. AVMs are the most dangerous of the various forms of intracranial vascular malformation because of their tendency to undergo spontaneous, often catastrophic, hemorrhage. AVMs may also cause neurological abnormalities (e.g., seizures) even in the absence of hemorrhage.


B. Brain, gross, coronal section: Diagnosis: Ruptured AVM with intraventricular and intracerebral hemorrhage. The hemorrhage can be intracerebral or extend into ventricles or subarachnoid space.

ARTERIOVENOUS MALFORMATION



TORTUOUS WORM-LIKE VASCULAR CHANNELS OF VARYING CALIBER AND WALL THICKNESSS
INTERMINGLE WITH BRAIN TISSUE

Brain. HE, 5X. Arteriovenous malformation is a low-pressure arteriovenous shunt composed of tortuous vessels of varying caliber and varying wall thickness surrounded by brain tissue. AVMs are present at birth and can increase in size. They are most common in the convexities of the hemispheres, but can occur anywhere. Small leaks may cause a seizure disorder or other



REFERENCES:


1. TamerYahya, et al
State-of-the-Art Review:
Stroke in young adults: Current trends, opportunities for prevention and pathways forward,
Am J of Preventive Cardiology, Volume 3, September 2020, 100085
<https://doi.org/10.1016/j.ajpc.2020.100085>
2. Jeffrey L Saver, MD, et al:
Thrombolytic Therapy in Stroke
<https://emedicine.medscape.com/article/1160840-overview#a2>



II CENTRAL NERVOUS SYSTEM TRAUMA

TRAUMATIC BRAIN INJURY

- Concussions
- Contusions (coup and contre-coup injury)
- Skull fractures
 - Lacerations
- Vascular injuries associated with trauma
 - Epidural, subdural, subarachnoid, intraparenchymal hemorrhages



Robbins and Kumar Basic Pathology, 11th Ed, Chapter 21, Fig 21-14, modified.

A. Epidural hematoma is caused by rupture of a meningeal artery and usually due to a skull fracture; leads to accumulation of blood between the skull and dura.

B. Subdural hematoma is caused by damage to bridging veins between the brain and the superior sagittal sinus; leads to accumulation of blood between the two layers of dura.

HEAD TRAUMA

- Penetrating or blunt; open or closed
- The magnitude and distribution of brain lesions depend on
 - the shape of the object causing the trauma,
 - the force of impact,
 - whether the head is in motion at the time of injury
- Severe brain damage can occur in the absence of external signs of head injury
- Severe lacerations and even skull fractures may not be always associated with injury to the brain (?)

CEREBRAL CONCUSSION

- Mild form of diffuse axonal injury (DAI) without permanent pathology (but: repeated concussion may cause CTE)
- Loss of consciousness is probably due to a functional disturbance of the reticular activating substance of the upper brainstem (the part of the CNS that is subjected to the highest twisting force during sagittal rotation of the hemispheres)


CTE = chronic traumatic encephalopathy

CEREBRAL CONTUSION

- Caused by rapid tissue displacement, disruption of vascular channels, and subsequent hemorrhage, tissue injury, and edema.
- The crests of the gyri are most susceptible to traumatic injury, because they are closest to the skull
- Contusions are common in regions, where the inside of the skull is irregular, e.g., the orbitofrontal regions and the temporal lobe tips.
- Contusions are wedge-shaped, with the widest aspect closest to the point of impact
- Blood seeps into the involved tissue across the cerebral cortex and into the white matter and subarachnoid spaces within hours.
- Function is impaired early, but morphologic changes (nuclear pyknosis, cytoplasmic eosinophilia, cellular disintegration) take about 24 hours to appear.
- Inflammatory response by PMNs is followed by macrophages.
- The superficial cortex is most severely affected in contrast with ischemic lesions, in which the superficial layer of cortex may be preserved.
- Coup injury occurs under the site of impact with an object.
- Contre coup injury occurs on the side opposite the area of the impact

CEREBRAL CONTUSION - CONTRE COUP INJURY

ACUTE CONTRE COUP CONTUSION




Contusions are common in areas, where the inside of the skull is irregular


- the orbitofrontal region
- tips of temporal lobes

76

Brain. Gross, ventral view. The crests of the gyri are most susceptible to traumatic forces producing this injury pattern



43 y o man. Base of brain. **Diagnosis:** Fresh traumatic subarachnoid hemorrhage. Coup contrecoup injury




A. Brain – gross; ventral view. 78 y o man. Remote contusions are present, mainly involving the right inferior frontal lobe. Note that the right olfactory bulb is absent, which would manifest as unilateral anosmia (the least tested cranial nerve on examination).

B. Brain – gross; ventral view. 61 y o woman. Base of frontal lobe. Remote frontal cerebral contusion. **Description:** Cystic frontobasal lesion of the cerebral cortex in a classic location. **Additional Findings:** Grand-Mal epilepsy.

LACERATION - PENETRATING HEAD TRAUMA

- Laceration and uncal herniation in the left hemisphere
- Brain edema with flattening of gyri and narrowing of sulci



Brain – gross. Ventral aspect of brain and a view of the convexity. Acute brain swelling in the closed cranial cavity is a serious condition. There is a laceration in the right hemisphere. The brain is edematous with flattening and widening of gyri and narrowing of sulci. Swelling of the right cerebral hemisphere has produced a shift with herniation of the uncus of the hippocampus through the tentorium, leading to the groove seen at the white arrow.

DIFFUSE AXONAL INJURY (DAI)

ETIOLOGY:

- Caused by blows to unsupported head
- Occurs most frequently in motor vehicle accidents

MECHANISM:

- Cerebrum moves back and forth, pivoting around the upper brainstem, which together with cerebellum is held firmly fixed by the tentorium
- Falx cerebri prevents side-to-side motion

CLINICAL FEATURES:


- Severe DAI causes immediate loss of consciousness after the injury
- Patients remain comatose or go into a persistent vegetative state
- 50% of patients who develop coma shortly after trauma are thought to have white matter damage and DAI

DIFFUSE AXONAL INJURY (DAI) continued

PATHOPHYSIOLOGY:

- Axons are stretched but not severed
- Sudden deformation causes compaction of neurofilaments, loss of microtubules and an arrest of the fast axoplasmic flow
- Mitochondria and other organelles accumulate proximal to the lesion and cause axonal swelling, which may lead to axonal rupture several hours later
- Influx of calcium through the stretched axolemma
- The swellings are located at nodes of Ranvier where the axolemma is more liable to deform because there is no myelin*
- Brain damage is most severe along midline structures (corpus callosum, brainstem), where the shear forces are greatest, and at the cortex-white matter junction because of the change in the consistency of brain tissue
- In severe traumatic brain injury DAI is made worse by widespread vascular injury and other traumatic lesions which cause cerebral edema and hypoxic ischemic encephalopathy (HIE)

*An HE stained section of DAI would look similar to the findings on slide 34 in the previous lecture on "Injury and Repair in the CNS and PNS"



Left: Axonal swellings can be detected with H&E and silver stains 15 hours after injury. Immunostains with antibodies to **Beta Amyloid Precursor Protein (BAPP)** can detect axonal lesions 2-3 hours after injury. BAPP is produced by neurons as a reaction to injury. **Right:** Brownish discoloration of white matter is due to old micro hemorrhages.

Glial Fibrillary acid protein (GFAP) can be detected in blood 1-2 hours after injury and reaches the peak in 20 hours.

DIFFUSE AXONAL INJURY

DAI: hemorrhages in corpus callosum



DAI: Duret hemorrhages in brain stem



- CAUSED BY BLOWS TO AN UNSUPPORTED HEAD
- OCCURS MOST FREQUENTLY IN MOTOR VEHICLE ACCIDENTS

83

ACUTE TRAUMATIC ENCEPHALOPATHY MARKERS

Serum levels of biomarkers correlate with the degree of traumatic brain injury (TBI)

- Used in the clinical settings to predict the severity of TBI.
- Used in Forensic pathology: there is a survival-time-dependent increase in astroglial GFAP and neuronal IL-6 expression in fatally injured human brain.
- **GFAP (Glial fibrillary acid protein)***
 - **Detectable in serum in <15 minutes after acute traumatic brain injury (ATBI)****
- **Ubiquitin C-terminal hydroxylase (UCH-L1)***
 - **Detectable in serum within an hour of injury**; the level correlates with the severity of injury determined by imaging findings and the Glasgow Coma Score.

* The FDA authorized the use of a blood test for glial fibrillary acidic protein (GFAP) and ubiquitin carboxy-terminal hydrolase L1 (UCH-L1) in mild traumatic brain injury (mTBI) in 2018.

A cartridge that can measure GFAP and UCH-L1 in whole blood within minutes after a TBI used in a hand-held iSTAT device (which can also measure several other analytes in whole blood) was tested by the US Military and was cleared by FDA for diagnosing acute TBI on April 1, 2024.


The cartridge uses a sandwich enzyme-linked immunosorbent assay (ELISA) to measure the amount of GFAP in the sample.

ACUTE TRAUMATIC ENCEPHALOPATHY MARKERS, cont.

- **IL-6:**
 - Elevated levels immediately after the trauma are associated with a worse outcome.
 - Produced by reactive astrocytes, microglial cells and neurons
 - May contribute to disruptions of the BBB and the progression of cerebral edema.
 - Promotes neurogenesis and regeneration in animal models of TBI.
- **Neurofilament light chain (NFL)**
 - Released from damaged neurons (analogous to troponin release from myocardium).
- **Tau:**
 - Significantly increased within hours of injury.

TRAUMATIC CNS HEMORRHAGE IN AN IMMATURE BABY


IMMATURE BRAIN; 28 WEEKS' GESTATION



- Immature brain is more susceptible to birth trauma
- Intraventricular hemorrhage typically arises in the germinal matrix, which is sensitive to hypoxia

86


This is intraventricular hemorrhage (IVH) which arose in the germinal matrix (subependymal region) of a 28 week gestational age newborn. The germinal matrix region is very susceptible to such lesions at this period of development, leading to IVH as a complication of prematurity.



EPIDURAL HEMATOMA


- Results from a skull fracture, which shears the **middle meningeal artery** resulting in bleeding and dissection of the dura from the inner table of the skull
- The expanding mass increases intracranial pressure and may produce herniation
- Patient may be initially unconscious, then have a **lucid period of minutes or hours** before developing increasing neurologic symptoms and losing consciousness
- Prompt evacuation of hematoma is necessary to save the patient's life

EPIDURAL HEMATOMA




The top of the skull is removed to reveal the middle meningeal artery which has emerged from the foramen spinosum to branch over the surface of the dura


Two views of a sagittal head CT- scan without contrast demonstrating a large epidural hematoma with right to left shift and ventricular narrowing.



EPIDURAL HEMATOMA



Left: Brain, gross, superior view. Right: Skull and dura. Horizontal section.



The dura has been reflected above to reveal the bridging veins that extend across to the superior aspect of the cerebral hemispheres. These can be torn with trauma, particularly if there is significant cerebral atrophy that exposes these veins even more.

ACUTE SUBDURAL HEMORRHAGE

- Results from trauma that shears the bridging veins and/or small arteries that traverse the dura
- Most vulnerable: superior cortical veins that lead to the superior sagittal sinus



CHRONIC SUBDURAL HEMATOMA

- May occur after a trivial head trauma in the young and the old
- Slowly accumulating hematoma becomes “localized” = surrounded by a membrane of fibrovascular tissue
- Degenerating blood clot increases in size by drawing fluid from the CSF and by new hemorrhages from the fragile vessels of the organizing granulation tissue
- Clinical symptoms and signs appear as a result of increased Intracranial pressure

ACUTE SUBDURAL HEMATOMA *

NO-CONTRAST CT:

A large acute subdural hematoma acts as a space-occupying lesion and causes a midline shift




Skull side




Bilateral subdural hematomas, larger in the left hemisphere

CHRONIC SUBDURAL HEMATOMA WITH MEMBRANE FORMATION *




These are two views of the same sagittal head CT scan without contrast demonstrating a large subdural hematoma with left to right shift and ventricular narrowing.

CHRONIC SUBDURAL HEMATOMA




80 y o man. **Diagnosis:** Chronic subdural hematoma. **Description:** Brown discoloration of the right hemisphere and the inside of the dura.



Brain – gross. Coronal section.

CHRONIC SUBDURAL HEMATOMA*



- A blood clot is located between two fibrotic membranes.
 - The inner membrane below the clot results entirely from organization.
 - The age of the hematoma, i.e., the time of the injury, can be estimated on the basis of the degree of the organization.
 - The hematoma may increase in size by drawing CSF, or because of rebleeding from the fragile vessels of the granulation tissue
- 96

Meninges, subgross, trichrome stain. Organizing subdural hematoma.

INCREASED INTRACRANIAL PRESSURE

DEFINITION:

- Raised intracranial pressure is an increase in mean CSF pressure **above 200 mm H₂O** with the patient recumbent

PATHOPHYSIOLOGY:

- Intracranial pressure increases when the volume of brain tissue increases beyond the limit permitted by compression of veins and displacement of CSF

ETIOLOGY:

- Most cases are associated with a mass effect, either diffuse, as in generalized brain edema, or focal, as with tumors, abscesses, or hemorrhages

CNS HERNIATIONS

SUBFALCINE (CINGULATE) HERNIATION:


- Unilateral or asymmetric expansion of a cerebral hemisphere displaces the cingulate gyrus under the falx cerebri.
- May be associated with compression of branches of the anterior cerebral artery.

TRANSTENTORIAL (UNCAL, MESIAL TEMPORAL) HERNIATION:

- The medial aspect of the temporal lobe is compressed against the free margin of the tentorium cerebelli.
- Increasing displacement of the temporal lobe compresses the third cranial nerve resulting in pupillary dilation and impairment of ocular movements on the side of the lesion.
- Compression of the posterior cerebral artery results in ischemic injury to the primary visual cortex.
- A large herniation compresses the contralateral cerebral peduncle resulting in **Kernohan's notch** and in ipsilateral hemiparesis.
- Progression of transtentorial herniation is often accompanied by flame shaped Duret hemorrhages in the midbrain and pons in the midline and paramedian regions due to tearing of penetrating veins and arteries that supply the upper brainstem.

James Watson Kernohan (1896-1981), an Irish-American pathologist, described a notched cerebral peduncle caused by contralateral transtentorial herniation.

MAJOR CNS HERNIATIONS



PATHOPHYSIOLOGY:

The cranial vault is subdivided by rigid dural folds (the falx and tentorium) into compartments.


If the expansion of brain in any of the compartments is sufficiently severe, the ICP rises, and **herniation** will occur.

Kernohan's notch:

In transtentorial herniation imaging studies show a midline shift and an indentation in the contralateral cerebral peduncle, which is compressed against the tentorium.

99

Modified from Robbins 10th ed, Figure 23-4. Major herniations of the brain: subfalcine, transtentorial, and tonsillar.




Brain – gross, mid-sagittal section and a cross section through midbrain and pons.

- There is kinking of the brainstem due to herniation and **Duret hemorrhages** are present in the midbrain and pons.
- Duret hemorrhages are small linear areas of bleeding in the midbrain and upper pons of the brainstem caused by **downward displacement of the brainstem**, which leads to stretching and rupture of small perforating branches of the basilar artery.
- The displacement is a result of uncal (temporal medial lobe) herniation due to increased intracranial pressure following acute hemorrhage, trauma (diffuse axonal "shearing" injury), tumor or inflammation/abscess.
- Infarction and venous thrombosis may cause global edema and elevated ICP leading to Duret hemorrhages.
- The diagnosis is made by CT or MRI.
- The prognosis is poor.

TONSILLAR HERNIATION *

- Displacement of the cerebellar tonsils through the foramen magnum
- Life-threatening because it causes brainstem compression and compromises vital respiratory and cardiac centers in the medulla oblongata



HERNIATED CEREBELLAR TONSILS

101

Cerebellar tonsillar herniation with displacement of the cerebellar tonsils into the foramen magnum and compression of medulla oblongata.

THE “SHAKEN BABY” SYNDROME*

PREVALENCE:

- Child abuse is the second most common cause of death in children under one year of age, after the sudden infant death syndrome
- Estimated 50,000 cases of shaken baby syndrome in the United States each year
- Nearly 25% of infants with shaken baby syndrome die from the sustained brain injuries
- Victims range in age from just a few days to five years, with an average age of six to eight months
- Young men (early 20s) are the usual perpetrators

CLINICAL PICTURE:

- Infants with these injuries are usually brought to the ER with a nonspecific clinical picture of hypotonia, listlessness, vomiting, irritability, and lethargy.
- Combination of subdural hematoma, subarachnoid hemorrhage, contusions, DAI, and retinal hemorrhage are often present
- Diagnosis may be missed unless a skeletal survey and funduscopic examination are done.

*Also known as abusive head trauma, shaken impact syndrome, inflicted head injury or whiplash shaken infant syndrome.

THE “SHAKEN BABY” SYNDROME continued

MECHANISM AND PATHOGENESIS:

- Shaking sets the head into a dangling motion and can cause **whiplash spinal cord injury**
- Movement of the brain inside the skull causes **subdural and subarachnoid hemorrhage, contusions, white matter hemorrhages (gliding hemorrhages), tears of the corpus callosum, and DAI**
- **Injuries are a combination of shaking and blunt impact**, thus, the term "shaking-impact syndrome" is more accurate
- **Retinal hemorrhages and hemorrhages within the optic nerve sheaths** are caused by shaking of the globe and tearing of delicate retinal vessels


THE “SHAKEN BABY” SYNDROME continued

SIGNIFICANCE OF RETINAL AND OPTIC NERVE INJURY:

- Retinal and optic nerve injury are an important component of the “shaken baby” syndrome and correlate with the severity of brain damage
- Retinal hemorrhages are extremely uncommon in accidental brain injury and thus documentation of retinal hemorrhage by funduscopic examination is important in the clinical and medico-legal evaluation of such cases

PROGNOSIS:

- Usually poor; 20% of cases result in death within the first few days
Survivors will most often be left with intellectual and developmental disabilities such as mental retardation or cerebral palsy
- Damage to the eyes can cause partial or total loss of vision
- Specialized care is usually required for the rest of the child’s life




Brain T2 MRI, coronal view and a funduscopy view on an infant. Suspicion of a shaken baby syndrome (SBS).

- Severe brain atrophy and large bilateral subdural hematomas.
- The subdural hematomas may have evolved from smaller ones that were present on admission.
- Subdural hematomas can develop following severe cerebral atrophy.

Clinical findings:

- **Retinal hemorrhages** are often large and confluent, an important component of the SBS, and correlate with the severity of brain damage.
- Retinal hemorrhages are probably caused by shaking of the vitreous body which pulls the retina and tears its delicate vessels.
- Retinal hemorrhages are **uncommon in accidental brain injury and are infrequently seen in patients with bleeding disorders, angiopathies, sepsis, or other non-traumatic settings.**
- **Documentation** by funduscopy examination is important in the clinical evaluation of the SBS.
- **Subdural hematomas and retinal hemorrhages are markers of the SBS.**
- Encephalopathy and death are caused mainly by **global HIE-the key pathology of the SBS-and increased intracranial pressure.**

- The back-and-forth motion of the head during shaking applies great stress to the cranio-cervical junction and the lower brainstem.
- In extreme cases, the spinal cord may separate from the brainstem.
- Lesser forces impair the function of vital centers in the brainstem causing **central apnea**.
- **Cerebral edema**, which follows HIE, raises intracranial pressure and makes HIE worse.
- **Vascular tears** from shaking cause microscopic or large parenchymal hemorrhages.
- **Infarcts** in an irregular geographic pattern are common.
- **DAI** may affect the cortex-white matter junction, the corpus callosum, internal capsule, or the brainstem and contributes to the encephalopathy.
- **Infants with the SBS usually also have rib, skull, and other fractures (often of varying ages), skin and soft tissue bruises, internal injuries, and other evidence of trauma.**



THE “SHAKEN BABY” SYNDROME

- Partial detachment of retina
- Retinal and optic nerve hemorrhages
- Severe brain atrophy secondary to hypoxic- ischemic encephalopathy (HIE) and bilateral subdural hemorrhages

See the T2 MRI image of this brain on the previous slide.

Note: Subdural hematomas may develop after minimal trauma in patients with severe cerebral atrophy.



THE “SHAKEN BABY” SYNDROME

OPTIC NERVE AND RETINAL HEMORRHAGES

