

7.0 SEED TREATMENT DIRECTIONS

7.1 Domestic Use

Brassica (Cole) Leafy Vegetables Crop Group 5

Crops		
Broccoli (<i>Brassica oleracea</i> var. <i>botrytis</i>)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)	
Broccoli, Chinese (gai lon) (<i>Brassica alboglabra</i>)	Cavalo Broccolo (<i>Brassica oleracea</i> var. <i>botrytis</i>)	
Broccoli Raab (rapini) (<i>Brassica campestris</i>)	Collards (<i>Brassica oleracea</i> var. <i>acephala</i>)	
Brussels sprouts (<i>Brassica oleracea</i> var. <i>gemmifera</i>)	Kale (<i>Brassica oleracea</i> var. <i>acephala</i>)	
Cabbage (<i>Brassica oleracea</i>)	Kohlrabi (<i>Brassica oleracea</i> var. <i>gongylodes</i>)	
Cabbage, Chinese (bok choy) (<i>Brassica chinensis</i>)	Mizuna (<i>Brassica rapa</i> var. <i>japonica</i>)	
Cabbage, Chinese (napa) (<i>Brassica pekinensis</i>)	Mustard greens (<i>Brassica juncea</i>)	
Cabbage, Chinese mustard (gai choy) (<i>Brassica campestris</i>)	Mustard spinach (<i>Brassica rapa</i> var. <i>perviridis</i>)	
	Rape greens (<i>Brassica napus</i>)	
Diseases	Use Rate (fl oz/100 lb)	Mefenoxam (g ai/100 kg)
Pythium Damping-off	0.09 – 0.68	2 – 15.5
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.1. 		
USE RESTRICTIONS		
<ul style="list-style-type: none"> Do not apply a pre-plant or an at-plant application of mefenoxam- or metalaxyl-containing products to fields planted with seeds treated with Apron XL 350ES. 		
Additional Use Directions		
Ridomil Gold® (foliar-applied mefenoxam) products are allowed to be applied beginning 6 weeks after transplanting.		