

565 IR STARK SHIFT THINGY

THE AUTHOR

ABSTRACT. Our chosen system of study is the 2-[(7-4-[N,N-Bis(4-methylphenyl)amino]phenyl-2,1,3- benzothiadiazol-4-yl)methylene]propanedinitrile (DTDCPB) chromophore prevalent in organic photovoltaic materials research. DTDCPB has nitrile groups that make it ideal for IR studies due to its strong signature stretching frequency. For the first component of our project, we hope to simulate some ground state IR spectra of DTDCPB in the presence of varying external fields. This can then be used to calculate the Stark shift and build a calibration curve for the effect of external field on the Stark shift of this molecule. Furthermore, we plan to probe the excited state of DTDCPB as well and see if we can look at Stark effects in the excited state as well. The third step we hope to investigate is a pump-probe transient IR study of this molecule where the excited state dynamics of DTDCPB can be observed. Hopefully our understanding of Stark effects in the ground and excited states will elucidate some of the dynamics of the transient IR study.

1. INTRODUCTION

Weeeeeee Typing thangs

2. THEORY

Some theoretical hand-wavey bullshit and lots of "beautiful mathematics" shall go here. Someday.

Here is an inline expression: $N = \frac{1}{2}\hbar$. Whereas here we have a single line expression:

$$\Psi = \sum_i A_{ij} c^\dagger.$$

$$A + (B + C) = (A + B) + C$$

$$A = \sum k_{ij}^2$$

3. METHODOLOGY

4. COMPUTATIONAL RESULTS

5. DISCUSSION

6. CONCLUSION

Date: May 17, 2016.