
Kinematic coverage of the closure-test datasets

- fixed-target e/μ DIS (SLAC, BCDMS, NMC, HERMES)
- HERA DIS (NC, CC, bottom)
- ν DIS (CHORUS, Pb)
- fixed-target Drell–Yan (E866 pp/pd, E605 Cu)
- collider Z/γ* (CDF, ATLAS, LHCb)
- synthetic Z row (plumbing only)
- in the union, not in this selection

JAM24

13 DIS + 3 DY datasets · 2771 points · + 13 lattice ensembles


`closure_JAM_truth_JAM_data, closure_NNPDF_truth_JAM_data`



NNPDF4.0

15 DIS + 17 DY datasets · 4199 points · + 13 lattice ensembles


`closure_JAM_truth_NNPDF_data, closure_NNPDF_truth_NNPDF_data`



Union (JAM24 ∪ NNPDF4.0)

16 DIS + 17 DY datasets · 4233 points · + 13 lattice ensembles


`closure_JAM_truth, closure_NNPDF_truth`



Mock / smoke

9 DIS + 3 DY datasets · 157 points · + 1 lattice ensemble

`closure_JAM_truth_small, closure_NNPDF_truth_small`



Rows after the closure cuts, as generation folds them; dashed lines are the DIS cuts $Q^2 \geq m_c^2$ and $W^2 > 10$ GeV^2. Each JAM/NNPDF truth pair fits identical kinematics.
Drell–Yan at LO: $x_{1,2} = (M/\sqrt{s})\, e^{\pm y}$ at $Q^2 = M^2$, bin centres, both x_1 and x_2 drawn; 1 bin centre with LO $x \geq 1$ (ATLAS 8 TeV high-mass) not drawn.