

# Spectral rigidity of some arithmetic surfaces

Justin Katz

Purdue University

June 22, 2022

# Spectra: rigidity and flexibility

In general, you cannot hear the shape of a drum:

# Spectra: rigidity and flexibility

In general, you cannot hear the shape of a drum:




Figure: Some isospectral drums

# Spectra: rigidity and flexibility

In general, you cannot hear the shape of a drum:




Figure: Some isospectral drums

Soft speculation: sufficiently special drums should sound unique.

# Spectra: rigidity and flexibility

In general, you cannot hear the shape of a drum:




Figure: Some isospectral drums

Soft speculation: sufficiently special drums should sound unique.



Figure: A very special (expensive) drum

# Hurwitz surfaces: the most symmetric of drums

A Hurwitz surface is a compact hyperbolic Riemann surface  $X$  of genus  $g$  such that  $|\operatorname{Aut}(X)| = 84(g - 1)$ .

# Hurwitz surfaces: the most symmetric of drums

A Hurwitz surface is a compact hyperbolic Riemann surface  $X$  of genus  $g$  such that  $|\text{Aut}(X)| = 84(g - 1)$ .




Figure: A genus 7 Hurwitz (Macbeath) surface [van Wijk, 2009]

# Specific speculation

In [Reid, 2014], Alan asked whether Hurwitz surfaces are determined up to isometry by their spectrum.



# Specific speculation

In [Reid, 2014], Alan asked whether Hurwitz surfaces are determined up to isometry by their spectrum.



Figure: not a genus 7 Hurwitz (Macbeath) surface

The following theorem answers in the affirmative infinitely many times (though, not all!)

The standard setup for arithmetic hyperbolic surfaces:

The standard setup for arithmetic hyperbolic surfaces:

- $k$  is a totally real number field, with ring of integers  $R_k$

The standard setup for arithmetic hyperbolic surfaces:

- $k$  is a totally real number field, with ring of integers  $R_k$
- $B$  is an indefinite quaternion algebra over  $k$ , split over a unique real place.

The standard setup for arithmetic hyperbolic surfaces:

- $k$  is a totally real number field, with ring of integers  $R_k$
- $B$  is an indefinite quaternion algebra over  $k$ , split over a unique real place.
- $\mathcal{O}$  is a maximal order in  $B$ , and  $\mathcal{O}^1$  its group of norm 1 units.

The standard setup for arithmetic hyperbolic surfaces:

- $k$  is a totally real number field, with ring of integers  $R_k$
- $B$  is an indefinite quaternion algebra over  $k$ , split over a unique real place.
- $\mathcal{O}$  is a maximal order in  $B$ , and  $\mathcal{O}^1$  its group of norm 1 units.
- $\Gamma(1)$  is the image of  $\mathcal{O}^1$  in  $\mathrm{PSL}(2, \mathbb{R})$  (an arithmetic lattice)

The standard setup for arithmetic hyperbolic surfaces:

- $k$  is a totally real number field, with ring of integers  $R_k$
- $B$  is an indefinite quaternion algebra over  $k$ , split over a unique real place.
- $\mathcal{O}$  is a maximal order in  $B$ , and  $\mathcal{O}^1$  its group of norm 1 units.
- $\Gamma(1)$  is the image of  $\mathcal{O}^1$  in  $\mathrm{PSL}(2, \mathbb{R})$  (an arithmetic lattice)
- For an ideal  $I$  of  $R_k$ , let  $\Gamma(I)$  be the kernel of the reduction map  $\Gamma(1) \rightarrow \mathrm{PSL}(2, R_k/I)$ .

## Theorem (McReynolds, Katz *in progress*)

Suppose that  $B$  has **type number 1**. Then for all but finitely many primes  $\mathcal{P}$  of  $R_k$ , and for all  $n \geq 0$ , the hyperbolic surface  $\Gamma(\mathcal{P}^n) \backslash \mathbb{H}$  is spectrally rigid.

The standard setup for arithmetic hyperbolic surfaces:

- $k$  is a totally real number field, with ring of integers  $R_k$
- $B$  is an indefinite quaternion algebra over  $k$ , split over a unique real place.
- $\mathcal{O}$  is a maximal order in  $B$ , and  $\mathcal{O}^1$  its group of norm 1 units.
- $\Gamma(1)$  is the image of  $\mathcal{O}^1$  in  $\mathrm{PSL}(2, \mathbb{R})$  (an arithmetic lattice)
- For an ideal  $I$  of  $R_k$ , let  $\Gamma(I)$  be the kernel of the reduction map  $\Gamma(1) \rightarrow \mathrm{PSL}(2, R_k/I)$ .



# Examples:

- If the field  $k$  has class number 1 (or more generally, the class group has no 2-torsion), then the type number hypothesis is satisfied for every indefinite quaternion algebra  $B$  over  $k$  (provided  $B$  is split over a unique real place)

# Examples:

- If the field  $k$  has class number 1 (or more generally, the class group has no 2-torsion), then the type number hypothesis is satisfied for every indefinite quaternion algebra  $B$  over  $k$  (provided  $B$  is split over a unique real place)
- In particular, this theorem applies to all *rational* indefinite quaternion algebras:

# Examples:

- If the field  $k$  has class number 1 (or more generally, the class group has no 2-torsion), then the type number hypothesis is satisfied for every indefinite quaternion algebra  $B$  over  $k$  (provided  $B$  is split over a unique real place)
- In particular, this theorem applies to all *rational* indefinite quaternion algebras: **it applies to  $\mathrm{PSL}(2, \mathbb{Z})$  and its principal congruence subgroups!**

# Examples:

- If the field  $k$  has class number 1 (or more generally, the class group has no 2-torsion), then the type number hypothesis is satisfied for every indefinite quaternion algebra  $B$  over  $k$  (provided  $B$  is split over a unique real place)
- In particular, this theorem applies to all *rational* indefinite quaternion algebras: **it applies to  $\mathrm{PSL}(2, \mathbb{Z})$  and its principal congruence subgroups!**
- Hurwitz surfaces arise from normal subgroups of the 2, 3, 7 triangle group, which arises as the norm 1 units of a quaternion algebra over the field  $\mathbb{Q}(2 \cos 2\pi/7)$ , which has class number 1. Hence the theorem applies: **principal congruence Hurwitz surfaces are spectrally rigid.**



Reid, A. W. (2014).

Traces, lengths, axes and commensurability.

*Annales de la Faculte des Sciences de Toulouse.*

*Mathematiques. Serie 6*, 23(5):1103–1118.



van Wijk, J. J. (2009).

Symmetric tiling of closed surfaces: Visualization of regular maps.

*ACM Transactions on Graphics*, 28(3):1–12.