
- Summarizing Distributions
- Consider a coin toss with P(H) = 0.6
- We can summarize this using a function:

$$P(n) = \begin{cases} 0.4 & \text{if } n=0 \\ 0.6 & \text{if } n=1 \end{cases}$$



$$\frac{P(n)}{P} = \begin{cases} \frac{1-p}{p} & \text{if } n=0 \\ \frac{1-p}{p} & \text{if } n=1 \end{cases}$$

 $(P(A) + P(\overline{A}) = 1$ Law of total probability)

Or, we can write this as:

$$P(n; p) = P \cdot (1-p)$$

f! called the model parameter Probability density function (PDF)

Set the parameter:
$$p = 0.6$$

The function becomes: $P(n) = 0.6^n \cdot 0.4^{-n}$

Now:

$$P(0) = 0.6^{\circ} \cdot 0.4^{\circ}$$

$$= 0.4$$

$$P(1) = 0.6^{\circ} \cdot 0.4^{\circ}$$

$$= 0.6$$

$$P(1) = 0.6' \cdot 0.4^{\circ}$$

= 0.6

- A PDF calculates probabilities for any value of a RV.
- It is (often) parameterized
- Several common PDFs define well known "distributions"
- The one above is the "Bernoulli Distribution"— single experiment
- "n is bernoulli distributed".

If we do (n) experiments. Success chance = ρ RV: X — number of successes in n experiments If I flip it 10 times, it can have P(10) successes P(10) needed! Let's set n=10, p=0.6 $P(X=1) = -7(0.6 \times 0.4^{9})$ $+ 0.4^{2} \times 0.6 \times 0.4^{9}$ $+ 0.4^{2} \times 0.6 \times 0.4^{9}$ $+ 0.4^{2} \times 0.6 \times 0.4^{9}$ How about P(2) ? But we know that this is a common pattern. X is a RV which belongs to the "Binomial Distribution". $P(X = k ; \underline{n}, \underline{p}) = (\underline{n}) \underline{p}^{k} (1-\underline{p})^{-k}$ Setting parameters: n= 10, p=0.6 $P(X=k) = \binom{10}{k} 0.6^{k} 0.4^{10-k}$ $P(X=2) = {\binom{10}{2}} \cdot 0.6^2 \cdot 0.4^8$ 0.25 Beil shape notice the shape!

0.2

0.15

0.16

0.01

0.02

0.04

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.00

0.0 b(x)

X

Create variations to the model by varying the parameters. Jupyter Lab!

- Practical applications.
 - Model the number of expected virus warnings
 - Don't want to annoy the user
 - Don't want to let viruses through
- Other distributions for discrete random variables
 - Poisson
 - Geometric