


1 | Solve Equations

Operation timed out. Arithmetic errors. #todo



2 | Read 1.B and 1.C

2.1 | General Notes

- The distributive property is extremely useful ### 1.35 Example
 - 1. If $b = 0$ then we can divide all x_3 by 5 and combine the last two terms to get F^3 , which is a vector space, without loss of generality. If not, then when you try to multiply by a scalar then you will find that the above reasoning breaks (i think).
 - 1. $f(x) = 0$ is continuous, so the additive identity exists. All sums of continuous functions result in continuous functions, so it is closed under addition. And all scalar multiples also work out.
 - 1. slightly awkward: i don't actually know what a differentiable real valued function is. #todo-exr0n
 - 1. (see previous)
 - 1. what does it mean for a sequence of complex numbers to have a limit 0? but I think you can use the same argument that the missing elements are just "collapsed" into one invisible one. ### 1.40 Definition direct sum
 - Something about uniqueness?

- If there is only one way to write zero then it works (1.44 Condition for a direct sum)


2.2 | Exercise to present

I would be interested in 7, 8, 10, 12, 14-19

3 | **2x2 Matrices that are Commutative**

(under multiplication, with all other 2×2 matrices)

Starting with $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} w & x \\ y & z \end{bmatrix} = \begin{bmatrix} w & x \\ y & z \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, I got $(x+y)(a-d) = (b+c)(w-z)$ and $by = cx$, but wasn't sure how to further develop it.



4 | Epilogue

Linear algebra homework always takes so long. Even though I skip like half of the problems. This is kind of an issue.