

# Joint stochastic theory of bedload transport and bed elevations: derivation of heavy-tailed resting times

James K. Pierce<sup>1\*</sup>, and Marwan A. Hassan<sup>1</sup>

<sup>1</sup>Department of Geography, University of British Columbia

## Key Points:

- We model fluvial bedload activity and local bed elevation as a two-species stochastic birth-death process.
- Computations show heavy-tailed power-law distributions of resting times for sediment undergoing burial, with universal tail parameters  $\alpha = 1.15$ .
- We discuss implications for bedload diffusion and suggest a new theoretical framework to approach the problem.

---

\*Vancouver, British Columbia, Canada

Corresponding author: James K. Pierce, [kpierce@alumni.ubc.ca](mailto:kpierce@alumni.ubc.ca)

## Abstract

A consensus has formed that fluvial bedload resting times lie on heavy-tailed statistical distributions which may result from sediment burial. However, due to observational difficulties, only a handful of experiments have resolved these distributions, and there have been few theoretical attempts to build understanding, leaving their generating mechanism and specific characteristics uncertain. In this work, we present a new theory describing bedload transport and bed elevation changes as a joint stochastic process and derive resting time distributions for sediment undergoing burial from the joint dynamics. Our theory implies heavy-tailed power-law distributions of resting times with universal tail behavior completely characterized by the mean erosion rate and its scaling with bed elevation changes. This finding provides a new perspective on heavy-tailed resting times with implications for the bedload diffusion problem.

## 1 Introduction

The majority of classic studies into fluvial sediment transport have attempted to relate the bulk downstream flux of bedload to characteristics of the hydraulic forcing (e.g. Yalin, 1972), yet the relevance of this approach to environmental problems is limited, as many contemporary issues require knowledge of the differences between motions of individual grains, and not just their average motion characteristics. For example, the export of contaminants from channels (e.g. Malmon, Reneau, Dunne, Katzman, & Drakos, 2005) and the morphological response of channels to ecological restoration efforts (e.g. Gaeuman, Stewart, Schmandt, & Pryor, 2017) or to changes in hydrology or sediment supply (e.g. Hassan & Bradley, 2017) is not determined by bulk bedload fluxes, highlighting individual motions of bedload as an important topic for geophysics research.

A significant complication is that individual grains transport within a noisy environment, with noise sources ranging across spatial and temporal scales from smaller scale fluid turbulence (Celik, Diplas, & Dancey, 2014) and variability in the arrangement of bed surface grains (Gordon, Carmichael, & Isackson, 1972), to larger scale channel morphology changes (Hassan & Bradley, 2017) and unsteady watershed hydrology (Phillips, Martin, & Jerolmack, 2013). As a result, the transport characteristics of individual grains are not deterministic (e.g. Einstein, 1937), even in the most controlled laboratory experiments (e.g. Böhm, Ancey, Frey, Reboud, & Ducottet, 2004; Charru, Mouilleron, & Eiff, 2004; Fathel, Furbish, & Schmeeckle, 2015; Heyman, Bohorquez, & Ancey, 2016).

In response to this, researchers have long considered probabilistic theories of individual motions based on random walk concepts, whereby bedload motions are approximated as alternating sequences of steps and rests, with step lengths and resting times treated as random variables drawn from statistical distributions (N. D. Bradley & Tucker, 2012; Einstein, 1937; Hassan, Church, & Schick, 1991; Nakagawa & Tsujimoto, 1976; Yano, 1969). In these theories, differences between the random motions of one grain and the next imply bedload diffusion, or a spreading apart of grains through time. Over long timescales, the diffusion characteristics predicted by these models critically differ depending on whether the step length and resting time distributions have light or heavy tails (e.g. N. D. Bradley, 2017).

Heavy-tailed distributions have exceedance functions  $P(X > x) \sim x^{-\alpha}$  with tail parameters  $\alpha < 2$ , meaning large values of  $x$  are relatively common, while light-tailed distributions have  $\alpha \geq 2$ , meaning large values of  $x$  are relatively rare. If both resting time and step distance distributions have light tails, the diffusion is said to be normal or Fickian, with a variance of particle positions  $\sigma_x^2$  scaling with time  $t$  as  $\sigma_x^2 \propto t$ . However, if either distribution has a heavy-tail, the diffusion is called anomalous, with a variance of particle position scaling as  $\sigma_x^2 \propto t^\gamma$ , where  $\gamma \neq 1$ . In this expression,  $\gamma < 1$  is called sub-diffusion and  $\gamma > 1$  is super-diffusion. In strongly asymmetric random walks such as bedload transport, heavy-tailed step lengths imply super-diffusion, while heavy-tailed resting times imply ei-

ther super or sub-diffusion, depending on  $\alpha$  (Weeks & Swinney, 1998; Weeks, Urbach, & Swinney, 1996).

Tracer experiments in gravel bed rivers show anomalous bedload diffusion (N. D. Bradley, 2017; Phillips et al., 2013), light-tailed step lengths (N. D. Bradley & Tucker, 2012; Hassan, Voepel, Schumer, Parker, & Fraccarollo, 2013), and heavy-tailed resting times (N. D. Bradley, 2017; Olinde & Johnson, 2015; Pretzlav, 2016; Voepel, Schumer, & Hassan, 2013), forming a coherent experimental picture of super-diffusive bedload transport, at least at long observation timescales (e.g. Martin, Jerolmack, & Schumer, 2012; Nikora, 2002). However, field studies have not resolved the mechanism generating observed heavy-tailed resting times (e.g. N. D. Bradley, 2017), and empirical distributions display clear differences in their form and characteristics, with different tail parameters (e.g. Olinde & Johnson, 2015) and sometimes truncation (e.g. N. D. Bradley, 2017) or tempering to light tails at large resting times (e.g. Voepel et al., 2013). These differences and the mechanism generating heavy-tailed resting times deserve further research attention.


A predominant hypothesis is that heavy-tailed resting times and anomalous diffusion originate from sediment burial (Martin, Purohit, & Jerolmack, 2014; Voepel et al., 2013; Wu et al., 2019). Conceptually, when grains rest on the bed surface, material transported from upstream can deposit on top of them, preventing entrainment until it’s removed, driving up resting times and imparting a heavy tail to the distribution. Martin et al. (2014) have provided the only direct support for this hypothesis. They traced grains in a narrow flume with clear sidewalls, directly resolving burial as the generator of heavy-tailed resting times, and they described their results with a theoretical model which is formally similar to an earlier effort by Voepel et al. (2013).

The models of Voepel et al. (2013) and Martin et al. (2014) consider bed elevations as a random walk and interpret resting times as return periods from above in the bed elevation time-series (e.g. Redner, 2007). Both models are successful in describing different experimental resting time distributions. However, the assumptions and results of these models are inconsistent with one another, and their treatment of bed elevations as a process independent of sediment transport is questionable, since erosion and deposition are the source of bed elevation changes (e.g. Wong, Parker, DeVries, Brown, & Burges, 2007).

In this work, we approach the problem from a different angle, making an extension of the stochastic bedload transport theory of Ancey, Davison, Böhm, Jodeau, and Frey (2008) to link bed elevation changes to the erosion and deposition events of individual grains, and we derive resting times as a consequence of this theory. The key assumptions of our model are: (1) bedload erosion and deposition can be characterized by probabilities per unit time, or rates (e.g. Ancey et al., 2008; Einstein, 1950); and (2) these rates are contingent on the local bed elevation, encoding the property that erosion of sediment is emphasized from regions of exposure, while deposition is emphasized in regions of shelter (e.g. Sawai, 1987; Wong et al., 2007). Our theory generates heavy-tailed distributions with no tempering and a universal tail parameter  $\alpha \approx 1.18$  for a particular non-dimensionalization of the resting time, showing close correspondence to the findings of Martin et al. (2014) and suggesting a correction to some imperfections in their results. We conclude the paper by framing our work in relation to earlier ideas and discussing the implications of this work on open problems in individual bedload motions and anomalous diffusion.

## 2 Stochastic theory

We define a volume of downstream length  $L$  which contains some number  $n$  of moving particles in the water flow and some number  $m$  of stationary particles composing the bed at time  $t$ . For simplicity, we consider all particles as approximately spherical with the same diameter  $2a$ , so their mobility and packing characteristics are similar. Following Ancey et al. (2008), we prescribe four events which can occur at any instant to modify the populations



**Figure 1.** Definition sketch of a control volume containing  $n$  moving grains and  $m$  resting grains. Migration, entrainment, and deposition processes are represented by curved arrows, and the bed elevation at some instant is depicted by dotted line. The bed is presented in a degraded state, where  $m < m_0$ . The distributions of  $n$  and  $m$  are indicated in the upper right panel, while the bottom panel is a time-series of bed elevations (1).

$n$  and  $m$ , and we characterize these events using probabilities per unit time, or rates. These are: (1) migration of a moving particle into the volume from upstream ( $n \rightarrow n+1$ ); (2) the entrainment (i.e., erosion) of a stationary particle into motion within the volume ( $m \rightarrow m-1$  and  $n \rightarrow n+1$ ); (3) the deposition of a moving particle to rest within the volume ( $m \rightarrow m+1$  and  $n \rightarrow n-1$ ); and (4) the migration of a moving particle out of the volume to downstream ( $n \rightarrow n-1$ ). As the events occur at random intervals, they set up a joint stochastic evolution of the populations  $n$  and  $m$  characterized by a joint probability distribution  $P(n, m, t)$  having marginals  $P(n, t) = \sum_m P(n, m, t)$  and  $P(m, t) = \sum_n P(n, m, t)$  for the number of particles in motion and at rest in the volume at  $t$ . These concepts are depicted in figure 1.

The populations  $n$  and  $m$  provide the bulk bedload flux  $q_s$  and the local bed elevation  $z$ . The mean bedload transport rate is given by  $q_s \propto u_s \langle n \rangle$ , where  $u_s$  is the characteristic velocity of moving bedload and  $\langle n \rangle = \sum_{n,m} n P(n, m)$  is the mean number of grains in motion (e.g. Ancey et al., 2008; Charru et al., 2004; Furbish, Haff, Roseberry, & Schmeeckle, 2012). The bed elevation is related to  $m$  though the packing geometry of the bed. To derive this, we prescribe a mean number of grains at rest  $m_0$  and introduce a packing fraction  $\phi$  of grains in the bed (Torquato, 2018). Considering a two-dimensional bed (e.g. Einstein, 1950; Paintal, 1971), the deviation from the mean bed elevation is

$$z(m) = \frac{\pi a^2}{\phi L} (m - m_0) = z_1 (m - m_0). \quad (1)$$

The constant  $z_1 = \pi a^2 / (\phi L)$  is an important scale of the problem.  $z_1$  is the magnitude of bed elevation change (in an average sense across the control volume) associated with the addition or removal of a single grain. We write the rates of the four possible transitions as

(e.g. Ancey et al., 2008):

$$R_{MI}(n+1, m|n, m) = \nu \quad \text{migration in,} \quad (2)$$

$$R_E(n+1, m-1|n, m) = \lambda(m) + \mu(m)n \quad \text{entrainment,} \quad (3)$$

$$R_D(n-1, m+1|n, m) = \sigma(m)n \quad \text{deposition,} \quad (4)$$

$$R_{MO}(n-1, m|n, m) = \gamma n \quad \text{migration out.} \quad (5)$$

These rates are independent of the past history of the populations and depend only on the current populations  $(n, m)$ . As a result, the system is Markovian (e.g. Cox & Miller, 1965; van Kampen, 1992), meaning time intervals between subsequent transitions are exponentially distributed (e.g. Gillespie, 2007).

In (2-5),  $\nu$  and  $\gamma$  are constants characterizing migration rates of individual grains into and out of the volume. They lack any dependence on the populations  $n$  and  $m$ . In contrast,  $\lambda(m)$ ,  $\mu(m)$ , and  $\sigma(m)$ , characterizing the entrainment, collective entrainment (e.g. Ancey et al., 2008; Heyman, Ma, Mettra, & Ancey, 2014; Heyman, Mettra, Ma, & Ancey, 2013), and deposition rates of individual grains are considered to depend on  $m$ . As is well-known, bed elevation changes modify the likelihood of entrainment and deposition in a negative feedback (Sawai, 1987; Wong et al., 2007); that is, aggradation increases the likelihood of entrainment, while degradation increases the likelihood of deposition. Wong et al. (2007) concluded that bed elevation changes induce an exponential variation in entrainment and deposition probabilities, while Sawai (1987) concluded that the variation is linear. For simplicity, we incorporate the scaling of Sawai (1987) and note its equivalence to the Wong et al. (2007) scaling when bed elevation changes are small. Because experimental distributions of bed elevations are usually symmetrical, (Martin et al., 2014; Singh, Fienberg, Jerolmack, Marr, & Foufoula-Georgiou, 2009; Wong et al., 2007), we expect the erosion and deposition feedbacks to be anti-symmetrical. That is, as bed elevation changes drive up (down) erosion rates, so they drive down (up) deposition rates to the same degree.

Summarizing these ideas, the entrainment and deposition rates can be written  $\chi(m) = \chi_0(1 \pm z_1 z(m)/(2l)^2)$ , where  $\chi = \lambda, \mu, \sigma$ , and the entrainment parameters take the plus sign, while deposition takes the minus, and we have introduced a length scale  $l$ . As we'll see, the variance of bed elevation turns out to be given by  $\text{var}(z) = (l/z_1)^2$ . Accordingly,  $l$  characterizes the range of bed elevation variations, which could be interpreted as the active layer depth (e.g. Church, 2017). Another perspective is that  $l$  is the distance of bed elevation change at which the entrainment and deposition rates are significantly affected. With these substitutions, the local bed elevation-dependent entrainment and deposition rates (3-4) can be written:

$$R_E(n+1, m-1|n, m) = [\lambda_0 + \mu_0 n] \left[ 1 + \frac{z_1 z(m)}{(2l)^2} \right], \quad \text{entrainment,} \quad (6)$$

$$R_D(n-1, m+1|n, m) = \sigma_0 \left[ 1 - \frac{z_1 z(m)}{(2l)^2} \right] n, \quad \text{deposition.} \quad (7)$$

At  $z(m) = 0$ , the rates reduce to those of the Ancey et al. (2008) theory. Away from this elevation, entrainment and deposition are alternatively suppressed and accentuated depending on the sign of  $z(m)$ .

In terms of the transition rates (2-7), we can obtain the Master equation for the probability flow using the forward Kolmogorov equation  $\partial P(n, m; t)/\partial t = \sum_{n', m'} R(n, m)P(n', m'; t)$  (e.g. Ancey et al., 2008; Cox & Miller, 1965; Gillespie, 1992) as

$$\begin{aligned} \frac{\partial P}{\partial t}(n, m; t) = & \nu P(n-1, m; t) + \{\lambda(m+1) + [n-1]\mu(m+1)\}P(n-1, m+1; t) \\ & + [n+1]\sigma(m-1)P(n+1, m-1; t) + [n+1]\gamma P(n+1, m; t) \\ & - \{\nu + \lambda(m) + n\mu(m) + n\sigma(m) + n\gamma\}P(n, m; t). \end{aligned} \quad (8)$$

The joint probability distribution  $P(n, m; t)$  solving this equation will fully characterize the statistics of  $n$  and  $m$ . We anticipate that solutions will adjust from the initial conditions to a steady-state distribution  $P_s(n, m)$ , independent of time, if the constant factors in the transition rates are representative of steady bedload transport conditions. This Master equation describes a two-species stochastic birth-death model (e.g. Cox & Miller, 1965) of a type well-known in the population ecology literature (e.g. Pielou, 1977; Swift, 2002). In our context, the two species are the moving and stationary grains in the volume.

### 3 Numerical simulations

Unfortunately, (8) does not appear to admit an analytical solution (but see Swift (2002) for a standard method which fails in this case). The difficulty stems from the product terms between  $n$  and  $m$ . In response, we numerically simulate (8) using the Gillespie algorithm (Gillespie, 1977, 1992, 2007). The Gillespie algorithm leverages the defining property of a Markov process: when transition rates do not depend on the past, time intervals between transitions are exponentially distributed (e.g. Cox & Miller, 1965).

**Table 1.** Parameters from Ancey et al. (2008) experiments describing the rates of migration in, entrainment, deposition, and migration out when  $z(m) = 0$ . All units are  $s^{-1}$  (probability/time). In our model, bed elevation changes modulate these rates in accord with (2-7).

Flow	$\nu$	$\lambda_0$	$\mu_0$	$\sigma_0$	$\gamma$
(a)	5.45	6.59	3.74	4.67	0.77
(g)	7.74	8.42	4.34	4.95	0.56
(i)	15.56	22.07	3.56	4.52	0.68
(l)	15.52	14.64	4.32	4.77	0.48
(n)	15.45	24.49	3.64	4.21	0.36


Therefore, to step the Markov process through a single transition, it's enough to draw a random value from the exponential distribution of transition intervals to determine the time of the next transition. Then, drawing another random value to choose the type of transition which occurs, using the relative probabilities (2-7), the transition can be enacted by shifting  $t$ ,  $n$  and  $m$  by the appropriate values (i.e., entrainment is  $m \rightarrow m - 1$  and  $n \rightarrow n + 1$ , and so on). This procedure can be iterated to form an exact realization of the stochastic process (e.g. Gillespie, 2007).

In this way, we simulated 5 flow conditions with 10 different values of  $l$  taken across a range from  $l = a$  (1 grain radius) to  $l = 10a$  (10 grain radii). These values lie in the range exhibited in the majority of available experimental data (Martin et al., 2014; Singh et al., 2009; Wong et al., 2007). For the migration, entrainment, and depo-

sition parameters at each flow condition ( $\nu, \lambda_0, \mu_0, \sigma_0, \gamma$ ), we used values measured by Ancey et al. (2008) in a series of flume experiments. These are summarized in table 1. Flow conditions are labeled (a), (g), and so on, roughly in order of increasing bedload flux (see Ancey et al. (2008) for more details). However, our conclusions are not dependent on these parameters. In all simulations, we take the packing fraction  $\phi = 0.6$ , a typical value for a pile of spheres (e.g. Bennett, 1972), and set  $L = 22.5\text{cm}$  and  $a = 0.3\text{cm}$ , in accord with the Ancey et al. (2008) experiments. Each simulation was run for 1500hrs of virtual time, a period selected to ensure convergence of the resting time statistics.

### 4 Results

Our simulations show noisy time-series of bedload activities and bed elevations (as seen in the bottom panel of figure 1). From our chosen initial conditions, all simulations show a rapid attainment of steady state conditions followed by a steady-state stochastic dynamics of  $n$  and  $m$ , supporting a time-independent joint distribution  $P_s(n, m)$ . We compute this joint distribution by counting occurrences of the states  $(n, m)$  in the simulated time series.




**Figure 3.** Marginal distributions of  $n$  and  $m$  for a subset of simulations. Some points have been omitted for clarity.

From this joint distribution we compute marginals  $P(n)$  and  $P(m)$  as explained in section 2.

Some of these marginal distributions are displayed in figure 3. Neglecting changes in bed elevation, Ancey et al. (2008) analytically derived negative binomial (NegBin) distributions for the bedload activity  $n$ , and this functional form seems preserved after including bed elevation changes, because all our computed distributions admit clean NegBin fits (figure 3a). For  $m$ , our computations show Gaussian distributions (figure 3b), consistent with our assumptions about the scaling of erosion and deposition rates with bed elevation changes (e.g. Wong et al., 2007).


From the marginal distributions, we calculate means and variances of bedload activity ( $n$ ) and elevation ( $m$ ). The mean bed elevation is  $m_0$ , the parameter in (1).  $m$  fluctuates around this value because it sets the equilibrium position of the elevation-related feedbacks within (1). The variance of  $m$  appears given by  $z_1^2 \text{var}(m) = l^2$ , as indicated in figure 2, consistent with our interpretation of  $l$  as a measure of bed elevation fluctuations. The moments of  $n$  are more difficult to understand. Without dwelling on the issue, the moments of  $n$  shift with the ratio  $l/z_1$  as a result of the feedback between bed elevation changes and bedload transport.

Now we describe the analysis of bedload resting times from time-series of  $m$ . Following Voepel et al. (2013) and Martin et al. (2014), we concentrate on a particular bed elevation  $m'$ , and find all time intervals separating deposition events at  $m = m'$  from erosion events at  $m = m' + 1$ . These are the return times from above of the bed conditional to the elevation  $m'$ . Binning these conditional return times (with logarithmic bins) and counting



**Figure 2.** Data from all simulations is plotted to show that  $l$  controls deviations of bed elevations:  $\text{var}(m) = (l/z_1)^2$ .





**Figure 4.** Resting time statistics differently with flow conditions and the bed elevation variance (i.e.,  $l$ ). Panel (a) shows differing flow conditions at a fixed  $l$  value, while panel (c) shows fixed flow conditions at differing  $l$ . When scaled by  $T_0$  (10), both types of difference collapse in the tails of the distributions, as shown in panels (b) and (d). Finally, in panel (d), the black dotted line indicates a power law decay of the collapsed tails parameter  $\alpha \approx 1.18$ .

the occurrences in each bin, we obtain a non-exceedance distribution of return times  $t_r$  held conditional to the elevation  $m'$ :  $P(T > t_r | m')$ . Using the marginal probability distribution of bed elevations, we derive the unconditional non-exceedance distribution of resting times as a sum over all elevations (Martin et al., 2014; Nakagawa & Tsujimoto, 1980; Voepel et al., 2013; Yang & Sayre, 1971):

$$P(T > t_r) = \sum_{m'} P(m') P(T > t_r | m'). \quad (9)$$

Some of these results are displayed in figure 4. In contrast to earlier works our analysis does not require binning over the elevation, since our elevation series is discrete (multiples of  $z_1$ ). Comparing panels (a) and (c) shows the resting time distributions scale with the intensity of bedload transport and the standard deviation of bed elevations ( $l$ ) differently. However, as shown in panels (b) and (d), a characteristic timescale  $T_0$  can be found to collapse away both of these differences.

We can obtain  $T_0$  heuristically by finding a characteristic speed of bed elevation change. Formally, the mean erosion rate is  $E = \sum_{n,m} R_E(n, m) P(n, m)$ . This is the number of grains leaving the bed per unit time. Since the removal of a single grain changes the bed elevation by  $z_1$ , bed elevations change with a characteristic speed  $z_1 E$ . Since the characteristic deviation



of elevation is  $l$ , the time required for the bed to shift through a characteristic deviation is

$$T_0 = \frac{l}{z_1 E}. \quad (10)$$

When scaling the resting time by this  $T_0$ , we obtain the collapse shown in figure 4. Using the log-likelihood estimation described by Newman (2005), we estimate for all return times satisfying  $T/T_0 > 10^3$ , the resting time non-exceedance distributions decay as a heavy-tailed power law with parameter  $\alpha = 1.18 \pm 0.32$ .

## 5 Discussion

Einstein (1937) created the first theory of individual bedload motions and diffusion, and his ideas can be viewed as the historical nexus of an entire paradigm of research which extends into the present day (e.g. Ancey et al., 2008; N. D. Bradley, 2017; Hassan et al., 1991; Heyman et al., 2016; Hubbell & Sayre, 1964; Martin et al., 2014; Nakagawa & Tsujimoto, 1976; Phillips et al., 2013; Yano, 1969). Works in this paradigm attempt to understand properties of bedload transport such as bulk fluxes or bedload diffusion from stochastic concepts of individual motions. With a few exceptions (e.g. Nakagawa & Tsujimoto, 1980; Wu et al., 2019; Yang & Sayre, 1971), theories within this paradigm are spatially one-dimensional, concentrating on the motion of grains in the downstream direction but neglecting the vertical dimension, wherein bed elevation changes can imply sediment burial (e.g. Martin et al., 2014; Voepel et al., 2013).

In the presented work, we have built upon these earlier works to create a theory of bedload transport and bed elevation changes based on individual bedload motions. In particular, we have generalized the Ancey et al. (2008) bedload theory to include bed elevation changes, incorporating feedbacks between the local bed elevation and the instantaneous erosion and deposition rates (e.g. Wong et al., 2007) to form a joint description of bulk bedload fluxes and bed elevation changes. Our simulations reveal negative binomial distributions for bedload activities and normal distributions for bed elevations, reproducing a wide set of experimental results (Ancey et al., 2008; Heyman et al., 2016; Martin et al., 2014; Singh et al., 2009; Wong et al., 2007). We believe two key contributions of our theory are that it (1) predicts resting time distributions for sediment undergoing burial, which are otherwise difficult to measure and poorly understood (e.g. N. D. Bradley, 2017; Martin et al., 2014; Voepel et al., 2013); and (2) provides a base for further developing of bedload transport theory including a vertical dimension of bed elevation dynamics. In this section, we will link our work to earlier research and discuss these two contributions in detail.

The link of our theory to earlier descriptions of bedload activities by Ancey et al. (2008) and bed elevations by Martin et al. (2014) follow as particular limit cases of (8). The Ancey et al. (2008) bedload theory is obtained in the limit of vanishing bed elevation fluctuations  $l \ll z_1$  (figure 2). In this case, we can neglect  $z_1 z(m)/(2l)^2$  in comparison to 1 in (6-7). Taking account of this change in (8), summing over  $m$ , and noting that  $n$  and  $m$  are independent in the limit derives the master equation of Ancey et al. (2008) for the bedload activity distribution. Likewise, Martin et al. (2014) bed elevation theory is obtained in the limit of vanishing bedload activity fluctuations. Writing

$$n = \langle n \rangle + \delta n, \quad (11)$$

where  $\delta n$  are the fluctuations of  $n$ , substituting this into (8), taking  $m_0 = 0$  (without any loss of generality), and summing the resulting equations over  $n$  gives

$$\frac{\partial}{\partial t} P(m, t) = E \left\{ \left[ 1 + \left( \frac{z_1}{2l} \right)^2 m \right] P(m+1, t) + \left[ 1 - \left( \frac{z_1}{2l} \right)^2 m \right] P(m-1, t) - 2P(m, t) \right\}, \quad (12)$$

where we have neglected all terms involving  $\delta n$  and used  $E = D$  for steady-state transport (e.g. Einstein, 1950). This is a discrete analog of the mean-reverting random walk used (in continuum form) by Martin et al. (2014) to model bed elevation changes.

Differences between our bed elevation statistics and those predicted Martin et al. (2014) given by (12) are induced by  $\delta n$  – the bedload activity fluctuations. When bedload fluctuations are relatively small  $|\delta n/\langle n \rangle| < 1$ , we expect the Martin et al. (2014) theory to be an excellent description of bed elevation changes, and it follows naturally from our joint treatment. However, in the Ancey et al. (2008) experiments, for example,  $|\delta n/\langle n \rangle|$  has instantaneous values often up to 3 or 5. Accordingly, we expect relatively large correction terms to (12) involving the bedload fluctuations  $\delta n$ . Although we do not expect these to modify bed elevation distributions away from Gaussian (figure 3), we do see a clear signature of these correction terms in our simulations. When we compute temporal autocorrelation functions of bed elevation time-series, we see clear differences from the exponential correlations expected in the Ornstein-Uhlenbeck process (e.g. Horsthemke & Lefever, 1984; Risken, 1989); in our case, the correlations are obviously non-exponential with a much slower decay. We conclude that sufficiently strong bedload fluctuations can induce much longer temporal correlations in the bed elevation time series than those predicted by the Martin et al. (2014) theory. Since the return times of a stochastic process are intimately related to its correlations (e.g. Redner, 2007), we expect bedload fluctuations are capable of inducing departures in the resting times of sediment undergoing burial from those derived by the Martin et al. (2014) theory, but if bedload fluctuations are relatively small, the Ornstein-Uhlenbeck theory seems aptly descriptive of resting times for sediment undergoing burial.

Now we discuss our computed resting time distributions (figure 4) and their properties with special emphasis on the implications of these results for bedload diffusion. To summarize, we have found upon scaling by the  $T_0$  – a timescale relating to the characteristic speed of bed elevation changes – that resting times for sediment undergoing burial should decay as a heavy-tailed power law with universal tail parameter  $\alpha = 1.18 \pm 0.32$ . For a general asymptotic power law, if  $\alpha > 1$ , neither the mean or variance of the resting time will converge, while if  $1 < \alpha < 2$ , the mean will converge while the variance will not (e.g. N. D. Bradley, 2017). Within our computational error, which stems from the finite duration of our simulations and the log-likelihood estimation (e.g. Newman, 2005), we cannot conclude whether the mean resting time will diverge, but we can conclude the variance will diverge. We can also expect anomalous super-diffusion at long time-scales. According to Weeks and Swinney (1998), if the step length distribution has a light tail (e.g. Hassan et al., 2013), our computed power-law resting times are sufficiently heavy-tailed to imply diffusion scaling as either  $\sigma_x^2 \propto t^{3-\alpha} \approx t^{\{1.82 \pm 0.32\}}$  or  $\sigma_x^2 \propto t^{2\alpha} \approx t^{\{3.64 \pm 1.28\}}$  at asymptotically large times (e.g. N. D. Bradley, 2017). In either case, the process of sediment burial induces an extreme super-diffusion of bedload: at long timescales, some grains will continue to transport downstream in alternate motion-rest sequences while others will become trapped under the bed for relatively very long periods of time, driving a rapid spreading of the population. However, we emphasize this is long-time diffusive scaling only. In fact, the conclusions of Weeks and Swinney (1998) are derived entirely from asymptotic arguments. Nikora (2002); Nikora, Goring, McEwan, Griffiths, and Nikora (2001) set an expectation of at least three ranges of bedload diffusion called local, intermediate, and global as a result of the scale dependence of the various processes at play in bedload motions, such as collisions, turbulent fluctuations, and burial. These three ranges have been captured by several theoretical models (Bialik, Nikora, & Rowiński, 2012; Fan, Singh, Guala, Foufoula-Georgiou, & Wu, 2016; Zhang, Meerschaert, & Packman, 2012, e.g.). While the transition point from local to intermediate scales is relatively well-understood, and there is some theoretical support for the notion of super-diffusion in the local range and normal diffusion in the intermediate range, the transition point between the intermediate and global ranges has so far, to our knowledge, defied theoretical description. Given the mounting theoretical arguments here and elsewhere (e.g. Martin et al., 2014; ?) and experimental support for global range super-diffusion (e.g. N. D. Bradley, 2017; Olinde & Johnson, 2015; Phillips et al., 2013; Pretzlav, 2016; Voepel et al., 2013), and the relative lack of understanding of the transition point between the intermediate and global ranges, we believe understanding of diffusion for many timescales

We predict heavy-tailed power law resting times with parameter  $\alpha \approx 1.18$  as a result of sediment burial. The tails of these distributions show near-perfect collapse upon scaling resting times by the factor  $T_0 = l/(z_1 E)$ , a characteristic timescale of bed elevation change, and we find the asymptotic behaviors occur for  $T/T_0 > 10^3$ . We note

In contrast, Martin et al. (2014)

In addition, since  $1 \leq \alpha \leq 2$ , tracer positions will have a convergent mean (e.g. N. D. Bradley, 2017; Weeks & Swinney, 1998). For comparison, Martin et al. (2012) found  $\alpha = 0.85$  from a three-dimensional flume experiment, Voepel et al. (2013) found  $\alpha = 0.88$  in data from the field study of Habersack (2001), Phillips et al. (2013) found  $\alpha = 1.12$  from their field study, Martin et al. (2014) found  $\alpha \approx 1.0$  from their simulations and two-dimensional flume experiments, and N. D. Bradley (2017) found  $\alpha = 0.67$  from field data. All of these values imply anomalous diffusion, like our simulation results, while only Phillips et al. (2013) and perhaps Martin et al. (2014) predict convergent means consistent with our results. These discrepancies may stem from our assumption that burial is the source of heavy tails in these field data. We have analyzed return periods of the bed elevation, but as noted by N. D. Bradley (2017), other return periods impede tracer motions in gravel bed rivers, such as the deposition of sediment on top of floodplains (e.g. D. N. Bradley & Tucker, 2013), or on top of high bars in the bends of a stream (e.g. N. D. Bradley, 2017).

Finally, we discuss the scope for extension of our model. The limiting relationship we have shown to Ancey et al. (2008), and knowledge of several follow-up works to Ancey et al. (2008) (e.g. Ancey & Heyman, 2014; Heyman, Bohórquez, & Ancey, 2015; Heyman et al., 2014), suggest an interesting and potentially useful generalization of our model. These follow-up works are based on chaining many Ancey et al. (2008) single-cell models together along a line, with migration out of one cell being migration into another, and they provide a framework to study spatial correlations in bedload transport (e.g. Heyman et al., 2015, 2014). One can imagine using our model (8) in the same way, chaining an array of volumes together along a line, generating a stochastic 1D morphodynamics theory ultimately rooted in the foundational ideas of Einstein (1950), and providing spatial correlations in bed elevation changes and bedload transport. An obvious issue with this idea is how to parameterize the entrainment and deposition rates within the model in terms of local and non-local bed elevations. Of course, this has been a longstanding issue in deterministic 1D morphodynamics as well (e.g. Tucker & Hancock, 2010). We speculate that the stochastic analogue of this issue could be easier to understand.

## 6 Conclusion

We predict heavy-tailed power law resting times with parameter  $\alpha \approx 1.18$  as a result of sediment burial. The tails of these distributions show near-perfect collapse upon scaling resting times by the factor  $T_0 = l/(z_1 E)$ , a characteristic timescale of bed elevation change, and we find the asymptotic behaviors occur for  $T/T_0 > 10^3$ .

Many open questions remain around the problem of anomalous bedload diffusion. In this paper, we have made incremental steps forward, by framing sediment burial directly in terms of the sediment transport process. When burial is a dominant process affecting the motions of individual grains, our results should serve as a prototype of this process, as they're derived under the assumptions of uniform grain size and equilibrium sediment transport in the spirit of Einstein (1950). We predict anomalous bedload diffusion  $\sigma_x^2 \propto t^{1.85}$  and a convergent mean, corroborating a subset of earlier studies. Most directly, we have derived the Martin et al. (2014) model as an approximate limit of our more general theory, while we found no connection to the Voepel et al. (2013) model. In closing, we highlight that other return processes in nature could impart heavy tails to fluvial bed material resting times (e.g. N. D. Bradley, 2017). To truly understand anomalous diffusion, we need to pin down these processes, quantify their relative importance, and learn to mix their effects.

## Acknowledgments

The simulation code is available upon request from the first author. We would like to thank Shawn Chartrand and Conor McDowell for helpful discussions.

## References

- Ancey, C., Davison, A. C., Böhm, T., Jodeau, M., & Frey, P. (2008). Entrainment and motion of coarse particles in a shallow water stream down a steep slope. *Journal of Fluid Mechanics*, 595(2008), 83–114. doi: 10.1017/S0022112007008774
- Ancey, C., & Heyman, J. (2014). A microstructural approach to bed load transport: mean behaviour and fluctuations of particle transport rates. *Journal of Fluid Mechanics*, 744(2014), 129–168. doi: 10.1017/jfm.2014.74
- Bennett, C. H. (1972). Serially deposited amorphous aggregates of hard spheres. *Journal of Applied Physics*, 43(6), 2727–2734. doi: 10.1063/1.1661585
- Bialik, R. J., Nikora, V. I., & Rowiński, P. (2012). 3D Lagrangian modelling of saltating particles diffusion in turbulent water flow. *Acta Geophysica*, 60(6), 1639–1660. doi: 10.2478/s11600-012-0003-2
- Böhm, T., Ancey, C., Frey, P., Reboud, J. L., & Ducottet, C. (2004). Fluctuations of the solid discharge of gravity-driven particle flows in a turbulent stream. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, 69(6 1), 1–13. doi: 10.1103/PhysRevE.69.061307
- Bradley, D. N., & Tucker, G. E. (2013). The storage time, age, and erosion hazard of laterally accreted sediment on the floodplain of a simulated meandering river. *Journal of Geophysical Research: Earth Surface*, 118(3), 1308–1319. doi: 10.1002/jgrf.20083
- Bradley, N. D. (2017). Direct Observation of Heavy-Tailed Storage Times of Bed Load Tracer Particles Causing Anomalous Superdiffusion. *Geophysical Research Letters*, 44(24), 12,227–12,235. doi: 10.1002/2017GL075045
- Bradley, N. D., & Tucker, G. E. (2012). Measuring gravel transport and dispersion in a mountain river using passive radio tracers. *Earth Surface Processes and Landforms*, 37, 1034–1045. doi: 10.1002/2017GL075045
- Celik, A. O., Diplas, P., & Dancey, C. L. (2014). Instantaneous pressure measurements on a spherical grain under threshold flow conditions. *Journal of Fluid Mechanics*, 741, 60–97. doi: 10.1017/jfm.2013.632
- Charru, F., Mouilleron, H., & Eiff, O. (2004). Erosion and deposition of particles on a bed sheared by a viscous flow. *Journal of Fluid Mechanics*, 519(2004), 55–80. doi: 10.1017/S0022112004001028
- Church, M. (2017). What is the "active layer"? *Water Resources Research*, 53, 5–10. doi: 10.1002/2016WR019675.Received
- Cox, D. R., & Miller, H. (1965). *The Theory of Stochastic Processes*. London: Chapman and Hall.
- Einstein, H. A. (1937). *Bed-load transport as a probability problem* (Unpublished doctoral dissertation). ETH Zurich.
- Einstein, H. A. (1950). *The Bed-Load Function for Sediment Transportation in Open Channel Flows* (Tech. Rep. No. 1026). Washington, DC: United States Department of Agriculture.
- Fan, N., Singh, A., Guala, M., Foufoula-Georgiou, E., & Wu, B. (2016). Exploring a semimechanistic episodic Langevin model for bed load transport: Emergence of normal and anomalous advection and diffusion regimes. *Water Resources Research*, 52(4), 3787–3814. doi: 10.1002/2016WR018704.Received
- Fathel, S. L., Furbish, D. J., & Schmeeckle, M. W. (2015). Experimental evidence of statistical ensemble behavior in bed load sediment transport. *Journal of Geophysical Research F: Earth Surface*, 120(11), 2298–2317. doi: 10.1002/2015JF003552
- Furbish, D. J., Haff, P. K., Roseberry, J. C., & Schmeeckle, M. W. (2012). A probabilistic description of the bed load sediment flux: 1. Theory. *Journal of Geophysical Research: Earth Surface*, 117(3). doi: 10.1029/2012JF002352

- Gaeuman, D., Stewart, R., Schmandt, B., & Pryor, C. (2017). Geomorphic response to gravel augmentation and high-flow dam release in the Trinity River, California. *Earth Surface Processes and Landforms*, 42(15), 2523–2540. doi: 10.1002/esp.4191
- Gillespie, D. T. (1977). Exact stochastic simulation of coupled chemical reactions. *Journal of Physical Chemistry*, 81(25), 2340–2361. doi: 10.1021/j100540a008
- Gillespie, D. T. (1992). *Markov Processes: An Introduction For Physical Sciences*. Academic Press, Inc.
- Gillespie, D. T. (2007). Stochastic Simulation of Chemical Kinetics. *Annual Review of Physical Chemistry*, 58(1), 35–55. doi: 10.1146/annurev.physchem.58.032806.104637
- Gordon, R., Carmichael, J. B., & Isackson, F. J. (1972). Saltation of Plastic Balls in a 'One-Dimensional' Flume. *Water Resources Research*, 8(2), 444–458.
- Habersack, H. M. (2001). Radio-tracking gravel particles in a large braided river in New Zealand: A field test of the stochastic theory of bed load transport proposed by Einstein. *Hydrological Processes*, 15(3), 377–391. doi: 10.1002/hyp.147
- Hassan, M. A., & Bradley, D. N. (2017). Geomorphic controls on tracer particle dispersion in gravel bed rivers. In *Gravel-bed rivers: Processes and disasters* (pp. 159–184). New York, NY: John Wiley & Sons Ltd. doi: 10.16719/j.cnki.1671-6981.2015.03.007
- Hassan, M. A., Church, M., & Schick, A. P. (1991). Distance of movement of coarse particles in gravel bed streams. *Water Resources Research*, 27(4), 503–511. doi: 10.1029/90WR02762
- Hassan, M. A., Voepel, H., Schumer, R., Parker, G., & Fraccarollo, L. (2013). Displacement characteristics of coarse fluvial bed sediment. *Journal of Geophysical Research: Earth Surface*, 118(1), 155–165. doi: 10.1029/2012JF002374
- Heyman, J., Bohórquez, P., & Ancey, C. (2015). Exploring the physics of sediment transport in non-uniform super-critical flows through a large dataset of particle trajectories. *J. Geophys. Res.:Earth Surf.*, submitted.
- Heyman, J., Bohorquez, P., & Ancey, C. (2016). Entrainment, motion, and deposition of coarse particles transported by water over a sloping mobile bed. *Journal of Geophysical Research: Earth Surface*, 121(10), 1931–1952. doi: 10.1002/2015JF003672
- Heyman, J., Ma, H. B., Mettra, F., & Ancey, C. (2014). Spatial correlations in bed load transport: Evidence, importance, and modeling. *Journal of Geophysical Research: Earth Surface*, 119(8), 1751–1767. doi: 10.1002/2013JF003003. Received
- Heyman, J., Mettra, F., Ma, H. B., & Ancey, C. (2013). Statistics of bedload transport over steep slopes: Separation of time scales and collective motion. *Geophysical Research Letters*, 40(1), 128–133. doi: 10.1029/2012GL054280
- Horsthemke, W., & Lefever, R. (1984). *Noise-Induced Transitions: Theory and Applications in Physics, Chemistry, and Biology*. Springer-Verlag.
- Hubbell, D. W., & Sayre, W. W. (1964). Sand Transport Studies with Radioactive Tracers. *J. Hydr. Div.*, 90(HY3), 39–68.
- Malmon, D. V., Reneau, S. L., Dunne, T., Katzman, D., & Drakos, P. G. (2005). Influence of sediment storage on downstream delivery of contaminated sediment. *Water Resources Research*, 41(5), 1–17. doi: 10.1029/2004WR003288
- Martin, R. L., Jerolmack, D. J., & Schumer, R. (2012). The physical basis for anomalous diffusion in bed load transport. *Journal of Geophysical Research: Earth Surface*, 117(1), 1–18. doi: 10.1029/2011JF002075
- Martin, R. L., Purohit, P. K., & Jerolmack, D. J. (2014). Sedimentary bed evolution as a mean-reverting random walk: Implications for tracer statistics. *Geophysical Research Letters*, 41(17), 6152–6159. doi: 10.1002/2014GL060525
- Nakagawa, H., & Tsujimoto, T. (1976). On Probabilistic Characteristics of Motion of Individual Sediment Particles on Stream Beds. In *Hydraulic problems solved by stochastic methods: Second international iahr symposium on stochastic hydraulics* (pp. 293–320). Lund, Sweden.
- Nakagawa, H., & Tsujimoto, T. (1980). Sand bed instability due to bed load motion. *Journal of the Hydraulics Division-ASCE*.
- Newman, M. E. (2005). Power laws, Pareto distributions and Zipf's law. *Contemporary*



- Physics*, 46(5), 323–351. doi: 10.1080/00107510500052444
- Nikora, V. (2002). On bed particle diffusion in gravel bed flows under weak bed load transport. *Water Resources Research*, 38(6), 1–9. Retrieved from <http://doi.wiley.com/10.1029/2001WR000513> doi: 10.1029/2001WR000513
- Nikora, V., Goring, D., McEwan, I., Griffiths, G., & Nikora. (2001). Spatially averaged open-channel flow over rough bed. *Journal of Hydraulic Engineering*, 127(February), 123–133.
- Olinde, L., & Johnson, J. P. L. (2015). Using RFID and accelerometer-embedded tracers to measure probabilities of bed load transport, step lengths, and rest times in a mountain stream. *Water Resources Research*, 51, 7572–7589. doi: 10.1002/2014WR016259
- Paintal, A. S. (1971). A Stochastic Model Of Bed Load Transport. *Journal of Hydraulic Research*, 9(4), 527–554. doi: 10.1080/00221687109500371
- Phillips, C. B., Martin, R. L., & Jerolmack, D. J. (2013). Impulse framework for unsteady flows reveals superdiffusive bed load transport. *Geophysical Research Letters*, 40(7), 1328–1333. doi: 10.1002/grl.50323
- Pielou, E. (1977). *Mathematical Ecology* (1st ed.). New York, NY: John Wiley & Sons Ltd.
- Pretzlav, K. L. (2016). *Armor development and bedload transport processes during snowmelt and flash floods using laboratory experiments, numerical modeling, and field-based motion-sensor tracers* (Unpublished doctoral dissertation). University of Texas.
- Redner, S. (2007). *A guide to first-passage processes*. Cambridge, UK: Cambridge University Press.
- Risken, H. (1989). *The Fokker-Planck Equation: Methods of Solution and Applications* (2nd ed.). Ulm: Springer-Verlag.
- Sawai, K. (1987). Dispersion of bed load particles. *Bull. Disas. Prev. Res. Inst., Kyoto Univ.*, 37(Part 1, No. 323).
- Singh, A., Fienberg, K., Jerolmack, D. J., Marr, J., & Foufoula-Georgiou, E. (2009). Experimental evidence for statistical scaling and intermittency in sediment transport rates. *Journal of Geophysical Research: Earth Surface*, 114(1), 1–16. doi: 10.1029/2007JF000963
- Swift, R. J. (2002). A Stochastic Predator-Prey Model. *Bulletin of the Irish Mathematical Society*, 48, 57–63.
- Torquato, S. (2018). Perspective: Basic understanding of condensed phases of matter via packing models. *Journal of Chemical Physics*, 149(2). doi: 10.1063/1.5036657
- Tucker, G. E., & Hancock, G. R. (2010). Modelling landscape evolution. *Earth Surface Processes and Landforms*, 35, 28–50. doi: 10.1002/esp
- van Kampen, N. G. (1992). *Stochastic Processes in Physics and Chemistry*. Amsterdam: Elsevier Science Publishers B.B.
- Voepel, H., Schumer, R., & Hassan, M. A. (2013). Sediment residence time distributions: Theory and application from bed elevation measurements. *Journal of Geophysical Research: Earth Surface*, 118(4), 2557–2567. doi: 10.1002/jgrf.20151
- Weeks, E. R., & Swinney, H. L. (1998). Anomalous diffusion resulting from strongly asymmetric random walks. *Physical Review E - Statistical Physics, Plasmas, Fluids, and Related Interdisciplinary Topics*, 57(5), 4915–4920. doi: 10.1103/PhysRevE.57.4915
- Weeks, E. R., Urbach, J. S., & Swinney, H. L. (1996). Anomalous diffusion in asymmetric random walks with a quasi-geostrophic flow example. *Physica D: Nonlinear Phenomena*, 97(1-3), 291–310. doi: 10.1016/0167-2789(96)00082-6
- Weiss, G. H. (1994). *Aspects and applications of the random walk*. Amsterdam: North Holland.
- Wong, M., Parker, G., DeVries, P., Brown, T. M., & Burges, S. J. (2007). Experiments on dispersion of tracer stones under lower-regime plane-bed equilibrium bed load transport. *Water Resources Research*, 43(3), 1–23. doi: 10.1029/2006WR005172
- Wu, Z., Foufoula-Georgiou, E., Parker, G., Singh, A., Fu, X., & Wang, G. (2019). Analytical Solution for Anomalous Diffusion of Bedload Tracers Gradually Undergoing Burial. *Journal of Geophysical Research: Earth Surface*, 124(1), 21–37. doi:

10.1029/2018JF004654

- Yalin, M. S. (1972). *Mechanics of Sediment Transport*. Pergamon Press.
- Yang, C. T., & Sayre, W. W. (1971). Stochastic model for sand dispersion. *Journal of the Hydraulics Division, ASCE*, 97(HY2).
- Yano, K. (1969). Tracer Studies on the Movement of Sand and Gravel. In *Proceedings of the 12th congress iahr, vol 2*. (pp. 121–129). Kyoto, Japan.
- Zhang, Y., Meerschaert, M. M., & Packman, A. I. (2012). Linking fluvial bed sediment transport across scales. *Geophysical Research Letters*, 39(20), 1–6. doi: 10.1029/2012GL053476