

# Analytical derivation of three-stage bedload diffusion

Kevin Pierce

July 12, 2019

## 1 Two-state random walk formalism

The formalism of two-state random walks [e.g. *Weiss*, 1976, 1994; *Masoliver*, 2016; *Masoliver and Lindenberg*, 2017] provides new traction on the problem of 3-stage diffusion in bedload transport [e.g. *Nikora et al.*, 2001; *Nikora*, 2002; *Zhang et al.*, 2012; *Fan et al.*, 2016]. Assuming residence time distributions  $\psi_{\pm}(t)$  in each state with associated survival probabilities  $\Psi_{\pm}(t) = \int_t^{\infty} dt' \psi_{\pm}(t')$ , probabilities of moving a distance  $x$  up to time  $t$  within a sojourn  $f_{\pm}(x, t)$ , denoting  $g_{\pm}(x, t) = f_{\pm}(x, t)\psi_{\pm}(t)$  and  $G_{\pm}(x, t) = f_{\pm}(x, t)\Psi_{\pm}(x, t)$ , and letting  $\omega_{\pm}(x, t)$  be the probability that a sojourn in state  $\pm$  ends at  $x, t$  provides

$$p_{\pm}(x, t) = \theta_{\pm}G_{\pm}(x, t) + \int_0^{\infty} dx' \int_0^{\infty} dt' \omega_{\mp}(x', t') G_{\pm}(x - x', t - t') \quad (1)$$

and

$$\omega_{\pm}(x, t) = \theta_{\pm}g_{\pm}(x, t) + \int_0^{\infty} dx' \int_0^{\infty} dt' \omega_{\mp}(x', t') g_{\pm}(x - x', t - t'). \quad (2)$$

Here  $\theta_{\pm}$  are the initial probabilities of being in each state with  $\theta_{+} + \theta_{-} = 1$ . The probability of the two-state random walker being at position  $x$  at time  $t$  is

$$p(x, t) = p_{-}(x, t) + p_{+}(x, t). \quad (3)$$

Denoting the laplace transform with respect to the variable  $q$  as  $\mathcal{L}_q$  and associating variables  $\eta$  and  $s$  with  $\mathcal{L}_x$  and  $\mathcal{L}_t$ , taking  $\mathcal{L}_x \mathcal{L}_t$  of (2) provides a much simpler algebraic problem for the probability  $p$ :

$$\tilde{\omega}_{\pm} = \frac{[\theta_{\pm} + \theta_{\mp} \tilde{g}_{\mp}] \tilde{g}_{\pm}}{1 - \tilde{g}_{-} \tilde{g}_{+}} \quad (4)$$

and (c.f. *Masoliver* [2016] eq. 20)

$$\tilde{p}_{\pm} = (\theta_{\pm} + \tilde{\omega}_{\mp}) \tilde{G}_{\pm} = \frac{\theta_{\pm} + \theta_{\mp} \tilde{g}_{\mp}}{1 - \tilde{g}_{-} \tilde{g}_{+}} \tilde{G}_{\pm}. \quad (5)$$

Therefore the double transform of the joint PDF reads (c.f. *Masoliver* [2016] eq. )

$$\tilde{p}(\eta, s) = \frac{\{\theta_{-} + \theta_{+} \tilde{g}_{+}\} \tilde{G}_{-} + \{\theta_{+} + \theta_{-} \tilde{g}_{-}\} \tilde{G}_{+}}{1 - \tilde{g}_{-} \tilde{g}_{+}} \quad (6)$$

This is a direct generalization of the famous Montroll-Weiss formula for a single state continuous-time random walk [e.g. *Weiss*, 1994].

For the evaluation of this formula, a useful fact to take account of is

$$\mathcal{L}_t\{\Psi_{\pm}(t); s\} = \int_0^{\infty} dt e^{-st} \int_t^{\infty} dt' \psi_{\pm}(t') = \frac{1 - \tilde{\psi}(s)}{s}. \quad (7)$$

Finally, owing to the definition of the double Laplace transform of  $p(x, t)$ :

$$\tilde{p}(\eta, s) = \int_0^{\infty} dt e^{-st} \int_0^{\infty} dx e^{-\eta x} p(x, t) \quad (8)$$

we see the (double) inverse transform  $p(x, t)$  may not be necessary to study the scaling of the variance  $\langle x(t)^2 \rangle = \int_0^\infty x^2 p(x, t)$  of the bedload tracer cloud since its Laplace transform follows from the second derivative of the double transform of  $p$  wrt  $\eta$ :

$$\mathcal{L}_t\{\langle x(t)^2 \rangle; s\} = \partial_\eta^2 \tilde{p}(\eta, s) \Big|_{\eta=0}. \quad (9)$$

I hope this allows evaluation of three-stage diffusion. More generally,

$$\mathcal{L}_t\{\langle x(t)^k \rangle; s\} = (-)^k \partial_\eta^k \tilde{p}(\eta, s) \Big|_{\eta=0}. \quad (10)$$

Actually, it may be easier to work with a cumulant-type expression since unlike *Masoliver* [2016]; *Masoliver and Lindenberg* [2017] we consider asymmetric processes where  $\langle x(t) \rangle \neq 0$  necessarily.

## 2 The Einstein theory

Taking  $g_-(x, t) = \delta(x)k_-e^{-k_-t}$  (rest) and  $g_+(x, t) = k_+e^{-k_+x}\delta(t)$  (step) reproduces the *Einstein* [1937] diffusion theory. In this case the double transforms are:

$$\tilde{g}_-(\eta, s) = \frac{k_-}{k_- + s} \quad (11)$$

$$\tilde{g}_+(\eta, s) = \frac{k_+}{k_+ + \eta}, \quad (12)$$

and the survival functions are

$$\Psi_-(t) = \int_t^\infty dt' k_- e^{-k_-t'} = e^{-k_-t} \quad (13)$$

and

$$\Psi_+(t) = \int_t^\infty dt' \delta(t') = 0, \quad (14)$$

meaning  $G_-(x, t) = \delta(x)e^{-k_-t}$  and  $G_+(x, t) = 0$ , providing

$$\tilde{G}_-(\eta, s) = \frac{1}{k_- + s}. \quad (15)$$

Taking  $\theta_- = 1$  and  $\theta_+ = 0$ , so the dynamics start at rest, the MW generalization (6) is

$$\tilde{p}(\eta, s) = \frac{\tilde{G}_-}{1 - \tilde{g}_- \tilde{g}_+} = \frac{1}{s + \frac{k_- \eta}{k_+ + \eta}}. \quad (16)$$

The Laplace transform of the mean is

$$\mathcal{L}_t\{\langle x(t) \rangle; s\} = \frac{k_-}{s^2 k_+} \quad (17)$$

so in real space it's  $\langle x(t) \rangle = k_-t/k_+$  as expected [e.g. *Einstein*, 1937; *Nakagawa and Tsujimoto*, 1976]. The Laplace transform of the second moment is

$$\mathcal{L}_t\{\langle x(t)^2 \rangle; s\} = 2 \left( \frac{k_-}{k_+} \right)^2 \left[ \frac{1}{k_-} \frac{1}{s^2} + \frac{1}{s^3} \right], \quad (18)$$

so using the inversion formula

$$\mathcal{L}\left\{ \frac{1}{s^{k+1}}; s \right\} = \frac{t^k}{k!} \quad (19)$$

implies a second moment  $\langle x(t)^2 \rangle = (k_-/k_+)^2 [2t/k_- + t^2]$  and a variance exemplifying the normal diffusion of bedload:

$$\sigma_x^2 = \text{var}\{x(t)\} = \frac{2k_-}{k_+^2} t. \quad (20)$$

The diffusivity  $D$  is given by the square of the mean step distance  $1/k_+$  divided by the mean resting time  $1/k_-$ :

$$D_{\text{Einstein}} = \frac{k_-}{k_+^2}. \quad (21)$$

Of course, for the Einstein theory a closed form solution of the pdf  $p(x, t)$  is possible to obtain [e.g. *Einstein*, 1937; *Hubbell and Sayre*, 1964; *Daly and Porporato*, 2006; *Daly*, 2019]. The first transform in (16) inverts easily for [i.e. *Prudnikov et al.*, 1986, 1.1.1.2]

$$\tilde{p}(\eta, t) = \exp \left\{ -\frac{k_- \eta}{k_+ + \eta} t \right\}. \quad (22)$$

Incidentally this single Laplace transform of  $p$  provides the cumulant generating function  $c(\eta, t) = \log \tilde{p}(-\eta, t)$  from which the variance follows from a more simple computation:  $\sigma_x^2(t) = \partial_\eta^2 c(\eta, t)|_{\eta=0}$ . The second inversion follows from *Prudnikov et al.* [1986, 2.2.2.8] noting  $\mathcal{L}_x^{-1}\{1\} = \delta(x)$ . The result is [e.g. *Daly*, 2019]

$$p(x, t) = e^{-k_- t - k_+ x} \left\{ \sqrt{\frac{k_- k_+ t}{x}} I_1 \left( 2\sqrt{k_- k_+ x t} \right) + \delta(x) \right\}. \quad (23)$$

This exact solution has been the benchmark theory of bedload diffusion for over 100 years. I have only derived it within a more general framework of multi-state random walks [e.g. *Weiss*, 1994].

### 3 The Lisle Theory

Apart from formulations of *Einstein* [1937] using different step length and resting time distributions than exponential [e.g. *Sayre and Hubbell*, 1965], the first significant advancement from *Einstein* [1937] was due to *Lisle et al.* [1998]. I need to carefully investigate whether *Gordon et al.* [1972] did it too. They imparted a finite duration to bedload motions instead of considering them instantaneous like Einstein. In this way they derived two stages of bedload diffusion. This approach is closely related to the so-called persistent diffusion model [*Balakrishnan and Chaturvedi*, 1988; *Van Den Broeck*, 1990], the diffusion of a particle driven by dichotomous Markov noise [e.g. *Horsthemke and Lefever*, 1984; *Risken*, 1989; *Bena*, 2006]. The mathematics were essentially developed by Takacs (1957).

It is obtained by the choice  $g_-(x, t) = \delta(x)k_- e^{-k_- t}$  (rest) and  $g_+(x, t) = \delta(x - vt)k_+ e^{-k_+ t}$  (motion). Hence motions occur with velocity  $v$  for a duration characterized by an exponential distribution with mean  $1/k_+$ , while rests occur for a duration characterized by an exponential distribution with mean  $1/k_-$ . We consider each of the extreme initial conditions in turn.

#### 3.1 starting at rest

If the process starts from rest, this means  $\theta_- = 1$  and  $\theta_+ = 0$ . In this case the Laplace transforms are

$$\tilde{g}_-(\eta, s) = \frac{k_-}{k_- + s}, \quad (24)$$

$$\tilde{g}_+(\eta, s) = \frac{k_+}{k_+ + \eta v + s} \quad (25)$$

and

$$\tilde{G}_{\pm}(\eta, s) = \frac{1}{k_{\pm}} \tilde{g}_{\pm}(\eta, s). \quad (26)$$

Plugging these into (6) gives

$$\tilde{p}(\eta, s) = \frac{k + s + \eta v}{v(k_{-} + s)\eta + ks + s^2}, \quad (27)$$

where  $k = k_{-} + k_{+}$ . This inverts to

$$\tilde{p}(x, s) = \frac{k(k + s)}{v(k_{-} + s)^2} \exp \left[ -\frac{s(k + s)}{v(k_{-} + s)} x \right] + \frac{1}{k_{-} + s} \delta(x). \quad (28)$$

Using the property  $\mathcal{L}\{f(x + a); s\} = e^{as} \mathcal{L}\{f(x); s\}$  along with the transform of a modified Bessel function gives

$$\begin{aligned} p(x, t) = \delta(x) e^{-k_{-}t} + \frac{k}{v} \exp \left[ -\frac{k_{+}x}{v} - k_{-} \left( t - \frac{x}{v} \right) \right] \Theta(t - x/v) \\ \times \left\{ I_0 \left( 2\sqrt{\frac{k_{-}k_{+}x}{v}} \left( t - \frac{x}{v} \right) \right) \right. \\ \left. \sqrt{\frac{k_{+}v(t - x/v)}{k_{-}x}} I_1 \left( 2\sqrt{\frac{k_{-}k_{+}x}{v}} \left( t - \frac{x}{v} \right) \right) \right\}, \end{aligned} \quad (29)$$

a non-trivial result. The non-dimensionalization proposed by Lisle is  $\xi = k_{+}x/v$  and  $\tau = k_{-}(t - x/v)$ . In this notation the result appears as

$$p(\xi, \tau) = \frac{k_{+}}{v} \delta(\xi) e^{-\tau - k_{-}\xi/k_{+}} + \frac{k}{v} e^{-\xi - \tau} \Theta(\tau) \Theta(\xi) \left\{ I_0(2\sqrt{\xi\tau}) + \frac{k_{+}}{k_{-}} \sqrt{\frac{\tau}{\xi}} I_1(2\sqrt{\xi\tau}) \right\} \quad (30)$$

### 3.1.1 moments having started at rest

The first derivative gives

$$\mathcal{L}\{\langle x(t) \rangle; s\} = vk_{-} \frac{1}{s^2(s + k)}, \quad (31)$$

while the second gives

$$\mathcal{L}\{\langle x^2(t) \rangle; s\} = 2v^2k_{-} \frac{s + k_{-}}{s^3(s + k)^2}. \quad (32)$$

Therefore the mean is easy to obtain as

$$\frac{k^2}{vk_{-}} \langle x \rangle = e^{-kt} + kt - 1, \quad (33)$$

while the second moment is

$$\frac{k^4}{2v^2k_{-}^2} \langle x^2 \rangle = \frac{(kt)^2}{2} - 2kt - (kt + 3)e^{-kt} + 3 \quad (34)$$

$$+ \frac{k}{k_{-}} [kt - 2 + (kt + 2)e^{-kt}]. \quad (35)$$

Manipulating the mean provides

$$\frac{k^4}{2v^2k_{-}^2} \langle x \rangle^2 = \frac{1}{2} e^{-2kt} + \frac{(kt)^2}{2} + \frac{1}{2} - kt + (-1 + kt)e^{-kt}, \quad (36)$$

so the variance is

$$\frac{k^4}{2v^2k_-^2}\sigma_x^2(t) = -kt - (2kt + 2)e^{-kt} + \frac{5}{2} - \frac{1}{2}e^{-2kt} + \frac{k}{k_-}[kt - 2 + (kt + 2)e^{-kt}]. \quad (37)$$

In this case as well,  $t \rightarrow 0$  gives  $\sigma_x^2 \propto t^3$ . This is a non-trivial result. The terms cancel each other in a highly coordinated way. This result shows a sharp transition between two diffusion regimes when the mean time in motion is smaller than the mean time at rest. When the opposite is true, there are three stages of diffusion with a sub-diffusive intermediate regime. Therefore in Einstein's concept of bedload transport where resting periods are longer than motion periods, there should be two diffusion regimes when the distributions are light-tailed.

### 3.2 starting in motion

Now I'll try using the opposite initial condition (the one chosen by *Lisle et al.* [1998]):  $\theta_+ = 1$  and  $\theta_- = 0$ . In this case

$$\tilde{p}(\eta, s) = \frac{k + s}{v(k_- + s)\eta + (k + s)s}, \quad (38)$$

and a first inverse transform gives

$$p(x, s) = \frac{k + s}{v(k_- + s)} \exp \left[ -\frac{(k + s)s}{v(k_- + s)}x \right], \quad (39)$$

which is the same as the other case but without a delta function term at  $x = 0$ . This can be manipulated to

$$p(x, t) = \mathcal{L}^{-1} \left\{ \frac{k + s}{v(k_- + s)} \exp \left[ -\frac{(k + s)s}{v(k_- + s)}x \right]; t \right\} \quad (40)$$

$$= e^{-k_-t} \mathcal{L}^{-1} \left\{ \frac{k_+ + s}{vs} \exp \left[ -\frac{(k_+ + s)(s - k_-)}{vs}x \right]; t \right\} \quad (41)$$

$$= e^{-k_-t - (k_+ - k_-)x/v} \mathcal{L}^{-1} \left\{ \frac{k_+ + s}{vs} \exp \left[ \frac{k_-k_+}{vs}x - \frac{xs}{v} \right]; t \right\} \quad (42)$$

$$= e^{-k_-t - (k_+ - k_-)x/v} \mathcal{L}^{-1} \left\{ \frac{k_+ + s}{vs} \exp \left[ \frac{k_-k_+}{vs}x \right]; t - x/v \right\} \quad (43)$$

$$= e^{-k_-t - (k_+ - k_-)x/v} \left[ \mathcal{L}^{-1} \left\{ \frac{k_+}{vs} \exp \left[ \frac{k_-k_+}{vs}x \right]; t - x/v \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{v} \exp \left[ \frac{k_-k_+}{vs}x \right]; t - x/v \right\} \right] \quad (44)$$

$$= e^{-k_-t - (k_+ - k_-)x/v} \left[ \frac{k_+}{v} I_0 \left( 2\sqrt{\frac{k_-k_+x}{v} \left( t - \frac{x}{v} \right)} \right) + \frac{1}{v} \mathcal{L}^{-1} \left\{ \exp \left[ \frac{k_-k_+}{vs}x \right] - 1; t - x/v \right\} + \frac{1}{v} \delta(t - x/v) \right] \quad (45)$$

$$= e^{-k_-t - (k_+ - k_-)x/v} \left[ \frac{k_+}{v} I_0 \left( 2\sqrt{\frac{k_-k_+x}{v} \left( t - \frac{x}{v} \right)} \right) + \frac{1}{v} \sqrt{\frac{k_-k_+x}{v(t - x/v)}} I_1 \left( 2\sqrt{\frac{k_-k_+x}{v} \left( t - \frac{x}{v} \right)} \right) + \frac{1}{v} \delta(t - x/v) \right]. \quad (46)$$

A key property here was  $e^{-ax}f(x) = \mathcal{L}^{-1}\{f(s + a); x\}$ , and those in the appendix. This type of math is not easy for me. In the non-dimensional variables this becomes

$$p(\xi, \tau) = e^{-\tau - \xi} \left[ \frac{k_+}{v} I_0(2\sqrt{\xi\tau}) + \frac{k_-}{v} \sqrt{\frac{\xi}{\tau}} I_1(2\sqrt{\xi\tau}) + \frac{k_-}{v} \delta(\tau) \right]. \quad (47)$$

This appears totally aligned with *Lisle et al.* [1998].

### 3.2.1 moments having started in motion

In this case the mean is

$$\frac{k^2}{v} \langle x(t) \rangle = k_- kt + k_+ (1 - e^{-kt}). \quad (48)$$

This result seems to make sense and agree with the asymptotic result of *Lisle et al.* [1998], especially since it has a constant term associated with starting in motion (so a displacement of order  $v/k_+$  is guaranteed). The second moment is

$$\frac{k^4}{2v^2} \langle x^2 \rangle = k_-^2 \frac{(kt)^2}{2} + k_- k_+ \{2kt - 2 + 2e^{-kt}\} + k_+^2 \{1 - (1 + kt)e^{-kt}\}. \quad (49)$$

I checked this result numerically. Performing more manipulations on the mean obtains

$$\frac{k^4}{2v^2} \langle x \rangle^2 = k_-^2 \frac{(kt)^2}{2} + k_- k_+ \{kt - kte^{-kt}\} + k_+^2 \left\{ \frac{1}{2} - e^{-kt} + \frac{1}{2} e^{-2kt} \right\}, \quad (50)$$

so the variance is

$$\frac{k^4}{2v^2} \sigma_x^2(t) = k_- k_+ \{kt - 2 + (2 + kt)e^{-kt}\} + k_+^2 \left\{ \frac{1}{2} - kte^{-kt} - \frac{1}{2} e^{-2kt} \right\}. \quad (51)$$

This variance describes short term ballistic diffusion  $\sigma_x^2 \propto t^3$  and long term normal diffusion. When the mean motion time is larger than the mean resting time, the cross-over is sharp. When the mean motion time is smaller than the mean resting time, there is a sub-diffusive cross-over zone. So in the zone envisioned by Einstein when motions are relatively short compared to rests, there is sharp cross-over between  $t^3$  super-diffusion and normal diffusion.

## 4 A new generalization: Randomly stopped Lisle process

Consider the choice  $\psi_1(t) = k_1 e^{-k_1 t}$ ,  $f_1(x, t) = \delta(x)$ ,  $\psi_2(t) = k_2 e^{-k_2 t}$ , and

$$f_2(x, t|T) = \delta(x - vt)\Theta(T - t) + \delta(x)\Theta(t - T). \quad (52)$$

This describes a *randomly stopped* variant of the Lisle process. When  $t > T$ , the motion state becomes a second rest state: the motion is turned off. In this case,  $g_1(\eta, s) = \frac{k_1}{k_1 + s}$  and  $G_1(\eta, s) = g_1(\eta, s)/k_1$  as before, while

$$g_2(\eta, s|T) = \frac{k_2}{k_2 + s} e^{-(k_2 + s)T} + \frac{k_2}{k_2 + \eta v + s} \left( 1 - e^{-(k_2 + \eta v + s)T} \right) \quad (53)$$

and  $G_2(\eta, s|T) = g_2(\eta, s|T)/k_2$ . You can maybe solve this conditional to  $T$  to obtain  $p(x, t|T)$ . Then in-putting a distribution for the trapping time  $T$ , the over-all distribution will be  $p(x, t) = \int dT p(x, t|T) f(T)$ .

## Appendix: Laplace transforms

This is just a reference of useful Laplace transforms for these types of studies.

$$\mathcal{L}^{-1} \left\{ p^{-2} (p + a)^{-1}; x \right\} = \frac{1}{a^2} (e^{-ax} + ax - 1), \quad (54)$$

[Prudnikov et al., 1986, 2.1.2.33];

$$\mathcal{L}^{-1} \left\{ p^{-2} (p + a)^{-2}; x \right\} = \frac{1}{a^3} [ax - 2 + (ax + 2)e^{-ax}], \quad (55)$$

[Prudnikov et al., 1986, 2.1.2.49];

$$\mathcal{L}^{-1}\left\{p^{-3}(p+a)^{-2}; x\right\} = \frac{1}{a^4} \left[ \frac{(ax)^2}{2} - 2ax - (ax+3)e^{-ax} + 3 \right], \quad (56)$$

derived from the previous result using the differentiation property;

$$\mathcal{L}^{-1}\left\{p^{-1}(p+a)^{-2}; x\right\} = \frac{1}{a^2} \left( 1 - (1+ax)e^{-ax} \right), \quad (57)$$

[Prudnikov et al., 1986, 2.1.2.47];

$$\mathcal{L}^{-1}\left\{p^{-1}(p+a)^{-1}; x\right\} = \frac{1}{a} (1 - e^{-ax}), \quad (58)$$

[Prudnikov et al., 1986, 2.1.2.31];

$$\mathcal{L}^{-1}\{e^{as}\tilde{f}(s); x\} = \mathcal{L}^{-1}\{\tilde{f}(s); x+a\}, \quad (59)$$

the shifting property;

$$\mathcal{L}^{-1}\left\{\frac{1}{s^\nu} \exp(a/s); t\right\} = \left(\frac{t}{a}\right)^{(\nu-1)/2} I_{\nu-1}(2\sqrt{at}), \quad (60)$$

valid for  $\nu > 1$  from [Prudnikov et al., 1986, 2.2.2.1];

$$\mathcal{L}^{-1}\{e^{a/p} - 1; x\} = \sqrt{\frac{a}{x}} I_1(2\sqrt{ax}), \quad (61)$$

and [Prudnikov et al., 1986, 2.2.2.8]. All of these Laplace transforms are verified from Wolfram Alpha so I expect no typos. Finally,

$$\mathcal{L}\left\{\frac{a}{bs+c}; s\right\} = \frac{a}{b} e^{-cx/b} \quad (62)$$

and

$$\mathcal{L}\left\{\frac{as}{bs+c}; s\right\} = \frac{a}{b} \left[ \delta(x) - \frac{c}{b} e^{-cx/b} \right]. \quad (63)$$

## References

- Balakrishnan, V., and S. Chaturvedi, Persistent Diffusion on a Line, *Physica A*, 148, 581–596, 1988.
- Bena, I., Dichotomous Markov noise: Exact results for out-of-equilibrium systems. A review, *International Journal of Modern Physics B*, 20(20), 2825–2888, doi:10.1142/S0217979206034881, 2006.
- Daly, E., Poisson process transient solution, *Personal Communication*, pp. 1–2, 2019.
- Daly, E., and A. Porporato, Probabilistic dynamics of some jump-diffusion systems, *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, 73(2), 1–7, doi:10.1103/PhysRevE.73.026108, 2006.
- Einstein, H. A., Bed-load transport as a probability problem, Ph.D. thesis, ETH Zurich, 1937.
- Fan, N., A. Singh, M. Guala, E. Foufoula-Georgiou, and B. Wu, Exploring a semimechanistic episodic Langevin model for bed load transport: Emergence of normal and anomalous advection and diffusion regimes, *Water Resources Research*, 52(4), 3787–3814, doi:10.1002/2016WR018704, Received, 2016.
- Gordon, R., J. B. Carmichael, and F. J. Isackson, Saltation of Plastic Balls in a 'One-Dimensional' Flume, *Water Resources Research*, 8(2), 444–458, 1972.
- Horsthemke, W., and R. Lefever, *Noise-Induced Transitions: Theory and Applications in Physics, Chemistry, and Biology*, Springer-Verlag, 1984.

- Hubbell, D. W., and W. W. Sayre, Sand Transport Studies with Radioactive Tracers, *J. Hydr. Div.*, 90(HY3), 39–68, 1964.
- Lisle, I. G., C. W. Rose, W. L. Hogarth, P. B. Hairsine, G. C. Sander, and J.-Y. Parlange, Stochastic sediment transport in soil erosion, *Journal of Hydrology*, 204, 217–230, 1998.
- Masoliver, J., Fractional telegrapher’s equation from fractional persistent random walks, *Physical Review E*, 93(5), 1–10, doi:10.1103/PhysRevE.93.052107, 2016.
- Masoliver, J., and K. Lindenberg, Continuous time persistent random walk: a review and some generalizations, *European Physical Journal B*, 90(6), doi:10.1140/epjb/e2017-80123-7, 2017.
- Nakagawa, H., and T. Tsujimoto, On Probabilistic Characteristics of Motion of Individual Sediment Particles on Stream Beds, in *Hydraulic Problems Solved by Stochastic Methods: Second International IAHR Symposium on Stochastic Hydraulics*, pp. 293–320, Lund, Sweden, 1976.
- Nikora, V., On bed particle diffusion in gravel bed flows under weak bed load transport, *Water Resources Research*, 38(6), 1–9, doi:10.1029/2001WR000513, 2002.
- Nikora, V., J. Heald, D. Goring, and I. McEwan, Diffusion of saltating particles in unidirectional water flow over a rough granular bed, *Journal of Physics A: Mathematical and General*, 34(50), doi:10.1088/0305-4470/34/50/103, 2001.
- Prudnikov, A. P., Y. A. Brychov, and O. Marichev, *Integrals and Series*, vol. 5, Gordon and Breach, New York, 1986.
- Risken, H., *The Fokker-Planck Equation: Methods of Solution and Applications*, 2nd ed., Springer-Verlag, Ulm, 1989.
- Sayre, W. W., and D. W. Hubbell, Transport and Dispersion of Labeled Bed Material North Loup River, Nebraska, *Transport of Radionuclides by Streams*, 1965.
- Van Den Broeck, C., Taylor Dispersion Revisited, *Physica A*, 168, 677–696, 1990.
- Weiss, G. H., The two-state random walk, *Journal of Statistical Physics*, 15(2), 157–165, doi:10.1007/BF01012035, 1976.
- Weiss, G. H., *Aspects and applications of the random walk*, North Holland, Amsterdam, 1994.
- Zhang, Y., M. M. Meerschaert, and A. I. Packman, Linking fluvial bed sediment transport across scales, *Geophysical Research Letters*, 39(20), 1–6, doi:10.1029/2012GL053476, 2012.