

20260928 Quiz

for (MATH210) Applied Complex Variables

Jungwoo Kim

Department of Mathematics

Pohang University of Science and Technology

Exercise 26.4. In each case, determine the singular points of the function and state why the function is analytic everywhere else:

(a) $f(z) = \frac{2z + 1}{z(z^2 + 1)}.$

(b) $f(z) = \frac{z^3 + i}{z^2 - 3z + 2}.$

(c) $f(z) = \frac{z^2 + 1}{(z + 2)(z^2 + 2z + 2)}.$

Solution. Each function is rational, so by the differentiation rules it is analytic wherever its denominator is nonzero.

(a)

$$z(z^2 + 1) = z(z - i)(z + i),$$

so the singular points are

$$\boxed{z = 0, \pm i}.$$

(b)

$$z^2 - 3z + 2 = (z - 1)(z - 2),$$

so the singular points are

$$\boxed{z = 1, 2}.$$

(c) Since

$$z^2 + 2z + 2 = (z + 1)^2 + 1 = (z + 1 - i)(z + 1 + i),$$

the singular points are

$$\boxed{z = -2, -1 \pm i}.$$

In each case the function is analytic at every other point.

□

Exercise 26.6. Use results in Sec. 24 to verify that the function

$$g(z) = \ln r + i\theta \quad (r > 0, 0 < \theta < 2\pi)$$

is analytic in the indicated domain of definition, with derivative $g'(z) = 1/z$. Then show that the composite function

$$G(z) = g(z^2 + 1)$$

is analytic in the quadrant $x > 0, y > 0$, with derivative

$$G'(z) = \frac{2z}{z^2 + 1}.$$

Suggestion. Observe that $\text{Im}(z^2 + 1) > 0$ when $x > 0, y > 0$.

Solution. For

$$g(z) = u(r, \theta) + iv(r, \theta), \quad u = \ln r, \quad v = \theta,$$

we have

$$ru_r = 1 = v_\theta, \quad u_\theta = 0 = -rv_r.$$

The relevant partial derivatives are continuous for $r > 0, 0 < \theta < 2\pi$. Hence the theorem in Sec. 24 gives

$$g'(z) = e^{-i\theta}(u_r + iv_r) = \frac{e^{-i\theta}}{r} = \boxed{\frac{1}{z}}.$$

Now let $z = x + iy$ with $x > 0$ and $y > 0$. Since

$$z^2 + 1 = (x^2 - y^2 + 1) + 2xyi,$$

we have

$$\text{Im}(z^2 + 1) = 2xy > 0.$$

Thus $z^2 + 1$ lies in the domain of the chosen branch of g . The inner function $z^2 + 1$ is entire, so the composition is analytic in the quadrant. By the chain rule,

$$\boxed{G'(z) = g'(z^2 + 1) \cdot 2z = \frac{2z}{z^2 + 1}}.$$

□

Exercise 26.7. Let a function f be analytic everywhere in a domain D . Prove that if $f(z)$ is real-valued for all z in D , then $f(z)$ must be constant throughout D .

Solution. Write

$$f(z) = u(x, y) + iv(x, y).$$

Since f is real-valued throughout D ,

$$v(x, y) = 0$$

there, so $v_x = v_y = 0$. Because f is analytic, the Cauchy–Riemann equations give

$$u_x = v_y = 0, \quad u_y = -v_x = 0.$$

Consequently

$$f'(z) = u_x + iv_x = 0$$

throughout D . By the theorem in Sec. 25, an analytic function whose derivative vanishes throughout a domain is constant. Hence f is constant on D .

□

Exercise 27.2. Let the function $f(z) = u(x, y) + iv(x, y)$ be analytic in a domain D , and consider the families of level curves $u(x, y) = c_1$ and $v(x, y) = c_2$, where c_1 and c_2 are arbitrary real constants. Prove that these families are orthogonal. More precisely, show that if $z_0 = (x_0, y_0)$ is a point in D which is common to two particular curves $u(x, y) = c_1$ and $v(x, y) = c_2$ and if $f'(z_0) \neq 0$, then the lines tangent to those curves at (x_0, y_0) are perpendicular.

Suggestion. Note how it follows from the pair of equations $u(x, y) = c_1$ and $v(x, y) = c_2$ that

$$u_x + u_y \frac{dy}{dx} = 0 \quad \text{and} \quad v_x + v_y \frac{dy}{dx} = 0.$$

Solution. At z_0 , the Cauchy–Riemann equations give

$$u_x = v_y, \quad u_y = -v_x.$$

Also,

$$|f'(z_0)|^2 = u_x^2 + u_y^2,$$

so $f'(z_0) \neq 0$ implies that u_x and u_y are not both zero.

Suppose first that $u_x u_y \neq 0$. Implicit differentiation gives the slopes

$$m_u = -\frac{u_x}{u_y}, \quad m_v = -\frac{v_x}{v_y}.$$

Using the Cauchy–Riemann equations,

$$m_v = -\frac{-u_y}{u_x} = \frac{u_y}{u_x},$$

and therefore

$$m_u m_v = -1.$$

Thus the tangent lines are perpendicular.

If $u_y = 0$, then $u_x \neq 0$. The level curve of u has a vertical tangent, while the Cauchy–Riemann equations give $v_x = 0$ and $v_y = u_x \neq 0$, so the level curve of v has a horizontal tangent. The case $u_x = 0$ is analogous. Hence the two families are orthogonal wherever $f'(z_0) \neq 0$. □

Exercise 29.1. Use the theorem in Sec. 28 to show that if $f(z)$ is analytic and not constant throughout a domain D , then it cannot be constant throughout any neighborhood lying in D .

Suggestion. Suppose that $f(z)$ does have a constant value w_0 throughout some neighborhood in D .

Solution. Suppose, to the contrary, that $f(z) = w_0$ throughout some neighborhood contained in D . Let

$$g(z) = w_0$$

for all $z \in D$. Then g is analytic in D , and f and g agree throughout that neighborhood. By the uniqueness theorem in Sec. 28,

$$f(z) = g(z) = w_0$$

throughout D . This contradicts the assumption that f is not constant on D . Therefore f cannot be constant throughout any neighborhood lying in D . □

Exercise 29.5. Show that if the condition that $f(x)$ is real in the reflection principle (Sec. 29) is replaced by the condition that $f(x)$ is pure imaginary, then equation (1) in the statement of the principle is changed to

$$\overline{f(\bar{z})} = -f(z).$$

Solution. Define

$$h(z) = -if(z).$$

If $f(x)$ is pure imaginary on the segment of the real axis, then $h(x)$ is real there. Since h is analytic in the same domain, the reflection principle applied to h gives

$$\overline{h(\bar{z})} = h(z).$$

Substituting $h = -if$,

$$i \overline{f(\bar{z})} = -if(z),$$

so

$$\boxed{\overline{f(\bar{z})} = -f(z)}.$$

Conversely, if this identity holds, then for real x ,

$$\overline{f(x)} = -f(x),$$

which means that the real part of $f(x)$ is zero. Hence $f(x)$ is pure imaginary, as required. $\square$

Exercise 30.5. Write $|\exp(2z + i)|$ and $|\exp(iz^2)|$ in terms of x and y . Then show that

$$|\exp(2z + i) + \exp(iz^2)| \leq e^{2x} + e^{-2xy}.$$

Solution. For any complex number w ,

$$|e^w| = e^{\operatorname{Re} w}.$$

Since $z = x + iy$,

$$\operatorname{Re}(2z + i) = 2x,$$

so

$$|\exp(2z + i)| = e^{2x}.$$

Also,

$$z^2 = x^2 - y^2 + 2xyi,$$

and hence

$$iz^2 = -2xy + i(x^2 - y^2).$$

Therefore

$$|\exp(iz^2)| = e^{-2xy}.$$

The triangle inequality now gives

$$\boxed{|\exp(2z + i) + \exp(iz^2)| \leq e^{2x} + e^{-2xy}}.$$

$\square$

Exercise 30.13. Let the function $f(z) = u(x, y) + iv(x, y)$ be analytic in some domain D . State why the functions

$$U(x, y) = e^{u(x, y)} \cos v(x, y), \quad V(x, y) = e^{u(x, y)} \sin v(x, y)$$

are harmonic in D .

Solution. Since f is analytic in D and the exponential function is entire, the composition $e^{f(z)}$ is analytic in D . Moreover,

$$\begin{aligned} e^{f(z)} &= e^{u+iv} \\ &= e^u (\cos v + i \sin v) \\ &= U + iV. \end{aligned}$$

By the theorem in Sec. 27, the real and imaginary components of an analytic function are harmonic. Therefore U and V are harmonic throughout D . □

Exercise 33.2. Show that

- (a) $\log e = 1 + 2n\pi i$, $n = 0, \pm 1, \pm 2, \dots$;
- (b) $\log i = \left(2n + \frac{1}{2}\right) \pi i$, $n = 0, \pm 1, \pm 2, \dots$;
- (c) $\log(-1 + \sqrt{3}i) = \ln 2 + 2\left(n + \frac{1}{3}\right) \pi i$, $n = 0, \pm 1, \pm 2, \dots$

Solution. For $z \neq 0$,

$$\log z = \ln |z| + i(\operatorname{Arg} z + 2n\pi), \quad n \in \mathbb{Z}.$$

- (a) Since $|e| = e$ and $\operatorname{Arg} e = 0$,

$$\boxed{\log e = 1 + 2n\pi i}.$$

- (b) Since $|i| = 1$ and $\operatorname{Arg} i = \pi/2$,

$$\boxed{\log i = i \left(\frac{\pi}{2} + 2n\pi \right) = \left(2n + \frac{1}{2} \right) \pi i}.$$

- (c) The number $-1 + \sqrt{3}i$ has modulus 2 and principal argument $2\pi/3$. Thus

$$\begin{aligned} \log(-1 + \sqrt{3}i) &= \ln 2 + i \left(\frac{2\pi}{3} + 2n\pi \right) \\ &= \boxed{\ln 2 + 2 \left(n + \frac{1}{3} \right) \pi i}. \end{aligned}$$

□

Exercise 33.4. Show that

$$\log(i^2) \neq 2 \log i$$

when the branch

$$\log z = \ln r + i\theta \quad \left(r > 0, \frac{3\pi}{4} < \theta < \frac{11\pi}{4} \right)$$

is used. (Compare this with the example in Sec. 33.)

Solution. Let g denote the single-valued logarithmic branch specified in the exercise, written as $\log$ in the textbook. For i , the arguments are $\pi/2 + 2k\pi$. The unique one in the branch interval

$$\frac{3\pi}{4} < \theta < \frac{11\pi}{4}$$

is $5\pi/2$. Hence

$$g(i) = \frac{5\pi}{2}i, \quad 2g(i) = 5\pi i.$$

On the other hand, $i^2 = -1$, and the unique argument of -1 in the same interval is π . Thus

$$g(i^2) = g(-1) = \pi i.$$

Therefore

$$\boxed{g(i^2) = \pi i \neq 5\pi i = 2g(i)}.$$

□

Exercise 33.11. Show in two ways that the function

$$\ln(x^2 + y^2)$$

is harmonic in every domain that does not contain the origin.

Solution. Let

$$H(x, y) = \ln(x^2 + y^2).$$

First, direct differentiation gives

$$H_x = \frac{2x}{x^2 + y^2}, \quad H_y = \frac{2y}{x^2 + y^2},$$

and, for $(x, y) \neq (0, 0)$,

$$H_{xx} = \frac{2(y^2 - x^2)}{(x^2 + y^2)^2}, \quad H_{yy} = \frac{2(x^2 - y^2)}{(x^2 + y^2)^2}.$$

Hence

$$H_{xx} + H_{yy} = 0.$$

All first- and second-order partial derivatives are continuous away from the origin, so H is harmonic there.

For a second argument, fix any $z_0 \neq 0$. A sufficiently small disk about z_0 avoids the origin and admits an analytic branch g of the logarithm. For a continuous choice $\theta(z)$ of argument on that disk,

$$g(z) = \ln |z| + i\theta(z), \quad \operatorname{Re} g(z) = \ln |z| = \frac{1}{2} \ln(x^2 + y^2).$$

By the theorem in Sec. 27, the real part of an analytic function is harmonic. Consequently $H = 2 \operatorname{Re} g$ is harmonic near z_0 . Since z_0 was arbitrary, H is harmonic throughout every domain that excludes the origin; a global logarithm branch on that domain is not required.

□

Exercise 34.1. Show that for any two nonzero complex numbers z_1 and z_2 ,

$$\operatorname{Log}(z_1 z_2) = \operatorname{Log} z_1 + \operatorname{Log} z_2 + 2N\pi i,$$

where N has one of the values $0, \pm 1$. (Compare with Example 2 in Sec. 34.)

Solution. Let

$$\theta_1 = \text{Arg } z_1, \quad \theta_2 = \text{Arg } z_2, \quad S = \theta_1 + \theta_2.$$

Since $-\pi < \theta_j \leq \pi$,

$$-2\pi < S \leq 2\pi.$$

The principal argument of the product is obtained by bringing S into $(-\pi, \pi]$:

$$\text{Arg}(z_1 z_2) = S + 2N\pi,$$

where

$$N = \begin{cases} -1, & S > \pi, \\ 0, & -\pi < S \leq \pi, \\ 1, & S \leq -\pi. \end{cases}$$

Thus $N \in \{0, \pm 1\}$. Since $|z_1 z_2| = |z_1| |z_2|$,

$$\begin{aligned} \text{Log}(z_1 z_2) &= \ln |z_1 z_2| + i \text{Arg}(z_1 z_2) \\ &= \ln |z_1| + \ln |z_2| + i(\theta_1 + \theta_2 + 2N\pi) \\ &= \boxed{\text{Log } z_1 + \text{Log } z_2 + 2N\pi i}. \end{aligned}$$

□

Exercise 34.2. Verify expression (4), Sec. 34, for $\log(z_1/z_2)$ by

- (a) using the fact that $\arg(z_1/z_2) = \arg z_1 - \arg z_2$ (Sec. 9);
- (b) showing that $\log(1/z) = -\log z$, $z \neq 0$, in the sense that $\log(1/z)$ and $-\log z$ have the same set of values, and then referring to expression (1), Sec. 34, for $\log(z_1 z_2)$.

Solution. The identities here are identities of multiple-valued logarithms, so equality means equality of the corresponding sets of values, with compatible values selected.

- (a) Choose arguments $\theta_1 \in \arg z_1$ and $\theta_2 \in \arg z_2$. The argument identity in Sec. 9 states that all arguments of z_1/z_2 have the form $\theta_1 - \theta_2 + 2k\pi$, where $k \in \mathbb{Z}$. Therefore all logarithmic values of the quotient are

$$\ln \frac{|z_1|}{|z_2|} + i(\theta_1 - \theta_2 + 2k\pi), \quad k \in \mathbb{Z}.$$

These are precisely the differences of the values $\ln |z_1| + i(\theta_1 + 2m\pi)$ and $\ln |z_2| + i(\theta_2 + 2n\pi)$, because $m - n$ ranges over $\mathbb{Z}$. Hence, as sets of values,

$$\boxed{\log \left(\frac{z_1}{z_2} \right) = \log z_1 - \log z_2}.$$

- (b) Write $z = re^{i\theta}$. The values of $\log(1/z)$ are

$$-\ln r + i(-\theta + 2k\pi), \quad k \in \mathbb{Z}.$$

The values of $-\log z$ are

$$-\ln r - i(\theta + 2n\pi) = -\ln r + i(-\theta - 2n\pi), \quad n \in \mathbb{Z}.$$

These are the same set of values, so

$$\log(1/z) = -\log z.$$

Since

$$\frac{z_1}{z_2} = z_1 \frac{1}{z_2},$$

expression (1), Sec. 34, gives

$$\log \left(\frac{z_1}{z_2} \right) = \log z_1 + \log(1/z_2) = \boxed{\log z_1 - \log z_2}.$$

□

Exercise 34.5. Let z denote any nonzero complex number, written

$$z = re^{i\Theta} \quad (-\pi < \Theta \leq \pi),$$

and let n denote any fixed positive integer. Show that all of the values of $\log(z^{1/n})$ are given by

$$\log(z^{1/n}) = \frac{1}{n} \ln r + i \frac{\Theta + 2(pn + k)\pi}{n},$$

where $p = 0, \pm 1, \pm 2, \dots$ and $k = 0, 1, 2, \dots, n-1$. Then, after writing

$$\frac{1}{n} \log z = \frac{1}{n} \ln r + i \frac{\Theta + 2q\pi}{n}, \quad q = 0, \pm 1, \pm 2, \dots,$$

show that the set of values of $\log(z^{1/n})$ is the same as the set of values of $(1/n) \log z$. Thus show that

$$\log(z^{1/n}) = \frac{1}{n} \log z,$$

where, corresponding to a value of $\log(z^{1/n})$ taken on the left, the appropriate value of $\log z$ is to be selected on the right, and conversely. [The result in Exercise 5, Sec. 33, is a special case of this one.]

Suggestion. Use the fact that the remainder upon dividing an integer by a positive integer n is always an integer between 0 and $n-1$, inclusive; that is, any integer q can be written $q = pn + k$, where $p \in \mathbb{Z}$ and $k \in \{0, 1, \dots, n-1\}$.

Solution. The n distinct values of $z^{1/n}$ are

$$w_k = r^{1/n} \exp \left(i \frac{\Theta + 2k\pi}{n} \right), \quad k = 0, 1, \dots, n-1.$$

For each fixed k , all values of $\log w_k$ are

$$\begin{aligned} \log w_k &= \frac{1}{n} \ln r + i \left(\frac{\Theta + 2k\pi}{n} + 2p\pi \right) \\ &= \frac{1}{n} \ln r + i \frac{\Theta + 2(pn + k)\pi}{n}, \quad p \in \mathbb{Z}. \end{aligned}$$

Thus these formulas give all values of $\log(z^{1/n})$.

On the other hand, the values of $(1/n) \log z$ are

$$\frac{1}{n} \ln r + i \frac{\Theta + 2q\pi}{n}, \quad q \in \mathbb{Z}.$$

By the division algorithm, every integer q can be written uniquely as

$$q = pn + k, \quad p \in \mathbb{Z}, \quad 0 \leq k \leq n-1.$$

Hence the two displayed families of values are exactly the same. Therefore, in the multiple-valued sense described in Sec. 34,

$$\boxed{\log(z^{1/n}) = \frac{1}{n} \log z}.$$

□