

20261001 Quiz

for (MATH102) Calculus II

Jungwoo Kim

Department of Mathematics

Pohang University of Science and Technology

Problem 1.5. Find a linear function $\ell: \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying

$$\ell(1, 2) = 3, \quad \ell(2, 3) = 5.$$

Solution. By Theorem 1.2, every linear function $\ell: \mathbb{R}^2 \rightarrow \mathbb{R}$ has the form

$$\ell(x, y) = px + qy$$

for some $p, q \in \mathbb{R}$. The given values yield

$$p + 2q = 3, \quad 2p + 3q = 5.$$

Thus $p = q = 1$, and hence

$$\boxed{\ell(x, y) = x + y}.$$

□

Problem 1.6. Let $\mathbf{u} = (u_1, u_2)$, $\mathbf{v} = (v_1, v_2)$, and $\mathbf{w} = (w_1, w_2)$ be vectors in $\mathbb{R}^2$, and let $a, b, c \in \mathbb{R}$. Using the definitions

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2), \quad c\mathbf{u} = (cu_1, cu_2), \quad -\mathbf{u} = (-u_1, -u_2),$$

show the following properties.

- (a) $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$.
- (b) $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$.
- (c) $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$.
- (d) $(a + b)\mathbf{u} = a\mathbf{u} + b\mathbf{u}$.
- (e) $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$.

Solution. Each identity follows componentwise from the definitions.

(a)

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2) = (v_1 + u_1, v_2 + u_2) = \mathbf{v} + \mathbf{u}.$$

(b)

$$\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (u_1 + v_1 + w_1, u_2 + v_2 + w_2) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}.$$

(c)

$$c(\mathbf{u} + \mathbf{v}) = (cu_1 + cv_1, cu_2 + cv_2) = c\mathbf{u} + c\mathbf{v}.$$

(d)

$$(a + b)\mathbf{u} = (au_1 + bu_1, au_2 + bu_2) = a\mathbf{u} + b\mathbf{u}.$$

(e)

$$\mathbf{u} + (-\mathbf{u}) = (0, 0) = \mathbf{0}.$$

□

Problem 1.16. Let

$$\mathbf{u} = (1, 3), \quad \mathbf{v} = (3, 1).$$

- (a) Determine whether $\mathbf{u}$ and $\mathbf{v}$ are linearly independent.
- (b) Express $(4, 4)$ as a linear combination of $\mathbf{u}$ and $\mathbf{v}$.
- (c) Express $(4, 5)$ as a linear combination of $\mathbf{u}$ and $\mathbf{v}$.

Solution.

- (a) If $a\mathbf{u} + b\mathbf{v} = \mathbf{0}$, then

$$a + 3b = 0, \quad 3a + b = 0.$$

These equations give $a = b = 0$. Thus $\mathbf{u}$ and $\mathbf{v}$ are linearly independent.

- (b) Solving

$$a + 3b = 4, \quad 3a + b = 4$$

gives $a = b = 1$. Hence

$$\boxed{(4, 4) = \mathbf{u} + \mathbf{v}}.$$

- (c) Solving

$$a + 3b = 4, \quad 3a + b = 5$$

gives

$$a = \frac{11}{8}, \quad b = \frac{7}{8}.$$

Therefore

$$\boxed{(4, 5) = \frac{11}{8}\mathbf{u} + \frac{7}{8}\mathbf{v}}.$$

□

Problem 1.21. Let $\mathbf{u} = (u_1, u_2)$, $\mathbf{v} = (v_1, v_2)$, and $\mathbf{w} = (w_1, w_2)$. Prove the following properties of the dot product.

- (a) $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$.
- (b) $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$.

Solution. Using the definition of the dot product,

- (a)

$$\begin{aligned} \mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) &= u_1(v_1 + w_1) + u_2(v_2 + w_2) \\ &= (u_1v_1 + u_2v_2) + (u_1w_1 + u_2w_2) \\ &= \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}. \end{aligned}$$

- (b)

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 = v_1u_1 + v_2u_2 = \mathbf{v} \cdot \mathbf{u}.$$

□

Problem 1.25. Let $\ell: \mathbb{R}^2 \rightarrow \mathbb{R}$ be linear and satisfy

$$\ell(2, 1) = 3, \quad \ell(1, 1) = 2.$$

Find a vector $\mathbf{c} \in \mathbb{R}^2$ such that

$$\ell(\mathbf{u}) = \mathbf{c} \cdot \mathbf{u}$$

for every $\mathbf{u} \in \mathbb{R}^2$.

Solution. By Theorem 1.3, such a vector has the form $\mathbf{c} = (p, q)$. The given values yield

$$2p + q = 3, \quad p + q = 2.$$

Thus $p = q = 1$, so

$$\boxed{\mathbf{c} = (1, 1)}.$$

□

Problem 1.26. Find the cosine of the angle between the vectors

$$\mathbf{u} = (1, 2), \quad \mathbf{v} = (3, 1).$$

Solution. By Theorem 1.5,

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}.$$

Here

$$\mathbf{u} \cdot \mathbf{v} = 5, \quad \|\mathbf{u}\| = \sqrt{5}, \quad \|\mathbf{v}\| = \sqrt{10}.$$

Therefore

$$\boxed{\cos \theta = \frac{5}{\sqrt{50}} = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}}.$$

□

Problem 1.41. Show that the vectors

$$(1, 1, 1, 1), \quad (0, 1, 1, 1), \quad (0, 0, 1, 1), \quad (0, 0, 0, 1)$$

are linearly independent in $\mathbb{R}^4$.

Solution. Let these vectors be $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$, respectively, and suppose

$$c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3 + c_4 \mathbf{v}_4 = \mathbf{0}.$$

The first coordinate gives $c_1 = 0$. The second then gives $c_2 = 0$, the third gives $c_3 = 0$, and the fourth gives $c_4 = 0$. Hence the only linear combination equal to $\mathbf{0}$ is the trivial one, so the four vectors are linearly independent.

□

Problem 1.42. Determine whether the vectors

$$(1, 2, 1), \quad (1, 2, 2), \quad (1, 2, 3), \quad (1, 2, 4)$$

are linearly dependent or linearly independent. Identify the theorem of this section that is particularly applicable.

Solution. There are four vectors in $\mathbb{R}^3$. By Theorem 1.8, any $n + 1$ vectors in $\mathbb{R}^n$ are linearly dependent. Taking $n = 3$, these four vectors are therefore

linearly dependent

.

The applicable result is Theorem 1.8. □

Problem 1.47. Show that the vectors

$$(1, 1, 1), \quad (1, 2, 3), \quad (3, 2, 1)$$

are linearly dependent.

Solution. Let

$$\mathbf{v}_1 = (1, 1, 1), \quad \mathbf{v}_2 = (1, 2, 3), \quad \mathbf{v}_3 = (3, 2, 1).$$

We find the nontrivial relation

$$-4\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3 = \mathbf{0}.$$

Hence the vectors are linearly dependent. □

Problem 1.62. Let

$$\mathbf{c} = (c_1, \dots, c_n) \in \mathbb{R}^n, \quad \mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n.$$

Show that the function

$$f(\mathbf{x}) = x_1 + \mathbf{c} \cdot \mathbf{x}$$

is linear by finding $\mathbf{d} = (d_1, \dots, d_n)$ such that

$$f(\mathbf{x}) = \mathbf{d} \cdot \mathbf{x}.$$

Solution. Since

$$f(\mathbf{x}) = (1 + c_1)x_1 + c_2x_2 + \dots + c_nx_n,$$

take

$$\mathbf{d} = (1 + c_1, c_2, \dots, c_n) = \mathbf{e}_1 + \mathbf{c}.$$

Then $f(\mathbf{x}) = \mathbf{d} \cdot \mathbf{x}$. The dot product with a fixed vector is linear in $\mathbf{x}$, so f is linear. □

Problem 1.84. Evaluate each of the following products.

(a)

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 6 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

(b)

$$\begin{pmatrix} 3 & -4 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} 6 \\ 6 \end{pmatrix}.$$

(c)

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

(d)

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}^2.$$

Solution. Using the definition of matrix multiplication,

(a)

$$\begin{bmatrix} x_1 \\ 2x_2 \\ 6x_3 \end{bmatrix}.$$

(b)

$$\begin{pmatrix} 3 & -4 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} 6 \\ 6 \end{pmatrix} = \begin{bmatrix} -6 \\ 42 \end{bmatrix}.$$

(c)

$$\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}.$$

(d)

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}^2 = \begin{pmatrix} \cos^2 \theta - \sin^2 \theta & -2 \sin \theta \cos \theta \\ 2 \sin \theta \cos \theta & \cos^2 \theta - \sin^2 \theta \end{pmatrix}$$

$$= \begin{bmatrix} \cos(2\theta) & -\sin(2\theta) \\ \sin(2\theta) & \cos(2\theta) \end{bmatrix}.$$

□

Problem 1.95. Find an equation of each plane in the form

$$ax + by + cz = d.$$

- (a) The plane through the origin with normal vector $(1, 0, 0)$.
- (b) The plane through $(0, 0, 0)$, $(0, 1, 1)$, and $(-3, 0, 0)$.
- (c) The plane containing $(1, 1, 1)$ and parallel to the plane

$$x - 3y + 5z = 60.$$

Solution.

- (a) A normal vector is $(1, 0, 0)$, and the plane passes through the origin. Hence

$$\boxed{x = 0}.$$

- (b) Two vectors in the plane are

$$\mathbf{v} = (0, 1, 1), \quad \mathbf{w} = (-3, 0, 0).$$

Their cross product is

$$\mathbf{v} \times \mathbf{w} = (0, -3, 3),$$

so a normal vector is $(0, -1, 1)$. Since the plane passes through the origin,

$$\boxed{-y + z = 0}.$$

- (c) A parallel plane has the same normal vector $(1, -3, 5)$, so it has the form

$$x - 3y + 5z = d.$$

Substituting $(1, 1, 1)$ gives $d = 3$. Therefore

$$\boxed{x - 3y + 5z = 3}.$$

□

Problem 1.104. Using the formula

$$\mathbf{v} \times \mathbf{w} = (v_2w_3 - v_3w_2, v_3w_1 - v_1w_3, v_1w_2 - v_2w_1),$$

evaluate the following cross products.

- (a) $(1, 0, 0) \times (0, 1, 0)$.
- (b) $(0, 1, 0) \times (1, 0, 0)$.
- (c) $(0, 0, 1) \times (a, b, c)$.

Solution. Substituting directly into the given formula,

(a)

$$(1, 0, 0) \times (0, 1, 0) = (0, 0, 1).$$

(b)

$$(0, 1, 0) \times (1, 0, 0) = (0, 0, -1).$$

(c)

$$(0, 0, 1) \times (a, b, c) = (-b, a, 0).$$

□

Problem 1.105. Using

$$\mathbf{e}_1 \times \mathbf{e}_2 = \mathbf{e}_3, \quad \mathbf{e}_2 \times \mathbf{e}_3 = \mathbf{e}_1, \quad \mathbf{e}_3 \times \mathbf{e}_1 = \mathbf{e}_2,$$

and the distributive and anticommutative laws

$$\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = \mathbf{u} \times \mathbf{v} + \mathbf{u} \times \mathbf{w}, \quad \mathbf{v} \times \mathbf{w} = -\mathbf{w} \times \mathbf{v},$$

evaluate the following cross products.

- (a) $\mathbf{e}_1 \times \mathbf{e}_2$.
- (b) $\mathbf{e}_2 \times (\mathbf{e}_1 + \mathbf{e}_3)$.
- (c) $(2\mathbf{e}_1 + 3\mathbf{e}_3) \times (a\mathbf{e}_1 + b\mathbf{e}_2 + c\mathbf{e}_3)$.

Solution.

(a)

$$\mathbf{e}_1 \times \mathbf{e}_2 = \mathbf{e}_3.$$

(b)

$$\begin{aligned} \mathbf{e}_2 \times (\mathbf{e}_1 + \mathbf{e}_3) &= \mathbf{e}_2 \times \mathbf{e}_1 + \mathbf{e}_2 \times \mathbf{e}_3 \\ &= -\mathbf{e}_3 + \mathbf{e}_1. \end{aligned}$$

Thus

$$\mathbf{e}_2 \times (\mathbf{e}_1 + \mathbf{e}_3) = \mathbf{e}_1 - \mathbf{e}_3.$$

(c) Using distributivity and anticommutativity,

$$\begin{aligned} (2\mathbf{e}_1 + 3\mathbf{e}_3) \times (a\mathbf{e}_1 + b\mathbf{e}_2 + c\mathbf{e}_3) &= 2b\mathbf{e}_3 - 2c\mathbf{e}_2 + 3a\mathbf{e}_2 - 3b\mathbf{e}_1 \\ &= -3b\mathbf{e}_1 + (3a - 2c)\mathbf{e}_2 + 2b\mathbf{e}_3. \end{aligned}$$

Hence

$$-3b\mathbf{e}_1 + (3a - 2c)\mathbf{e}_2 + 2b\mathbf{e}_3.$$

□

Problem 2.4. Show that a constant function $\mathbf{F}: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear if and only if

$$\mathbf{F}(\mathbf{x}) = \mathbf{0}$$

for every $\mathbf{x} \in \mathbb{R}^n$.

Solution. Suppose first that $\mathbf{F}$ is linear. Since $\mathbf{F}$ is constant,

$$\mathbf{F}(\mathbf{x}) = \mathbf{F}(\mathbf{0})$$

for every $\mathbf{x}$. By linearity,

$$\mathbf{F}(\mathbf{0}) = \mathbf{F}(0\mathbf{x}) = 0\mathbf{F}(\mathbf{x}) = \mathbf{0}.$$

Thus $\mathbf{F}$ is the zero function.

Conversely, if $\mathbf{F}(\mathbf{x}) = \mathbf{0}$ for every $\mathbf{x}$, then

$$\mathbf{F}(c\mathbf{x}) = \mathbf{0} = c\mathbf{F}(\mathbf{x})$$

and

$$\mathbf{F}(\mathbf{x} + \mathbf{y}) = \mathbf{0} = \mathbf{F}(\mathbf{x}) + \mathbf{F}(\mathbf{y}).$$

Hence $\mathbf{F}$ is linear. □

Problem 2.5. The plane in $\mathbb{R}^3$ given by

$$z = 2x + 3y$$

is the 0 level set of a linear function $\ell: \mathbb{R}^3 \rightarrow \mathbb{R}$. Find such a function ℓ .

Solution. Rewrite the plane equation as

$$2x + 3y - z = 0.$$

Thus the linear function

$$\boxed{\ell(x, y, z) = 2x + 3y - z}$$

has exactly this plane as its 0 level set. □

Problem 2.13. Sketch or describe the c level sets $f(\mathbf{x}) = c$ in $\mathbb{R}^3$ for the following functions $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ and values of c .

(a) $f(\mathbf{x}) = \|\mathbf{x}\|^2$, for $c = \frac{1}{2}, 1, 2$.

(b) $f(\mathbf{x}) = \|\mathbf{x}\|$, for $c = \frac{1}{2}, 1, 2$.

(c) $f(\mathbf{x}) = \frac{1}{\|\mathbf{x}\|^2}$, for $c = \frac{1}{2}, 1, 2$.

(d) $f(\mathbf{x}) = \mathbf{u} \cdot \mathbf{x}$, for $c = \frac{1}{2}, 1, 2$, where $\mathbf{u}$ is a unit vector. Hint: write $\mathbf{x}$ as the sum of a vector parallel to $\mathbf{u}$ and a vector orthogonal to $\mathbf{u}$.

Solution.

(a) The equation $\|\mathbf{x}\|^2 = c$ describes a sphere centered at the origin with radius $\sqrt{c}$. Thus the radii are

$$\boxed{\frac{1}{\sqrt{2}}, 1, \sqrt{2}}.$$

- (b) The equation $\|\mathbf{x}\| = c$ describes a sphere centered at the origin with radius c . Thus the radii are

$$\left[\frac{1}{2}, 1, 2 \right].$$

- (c) The equation

$$\frac{1}{\|\mathbf{x}\|^2} = c$$

is equivalent to $\|\mathbf{x}\| = 1/\sqrt{c}$. Thus the level sets are spheres centered at the origin with radii

$$\left[\sqrt{2}, 1, \frac{1}{\sqrt{2}} \right].$$

- (d) Write

$$\mathbf{x} = t\mathbf{u} + \mathbf{y}, \quad \mathbf{y} \perp \mathbf{u}.$$

Since $\|\mathbf{u}\| = 1$,

$$\mathbf{u} \cdot \mathbf{x} = t.$$

Hence the level set $\mathbf{u} \cdot \mathbf{x} = c$ is the plane perpendicular to $\mathbf{u}$ through the point $c\mathbf{u}$. For $c = \frac{1}{2}, 1, 2$, the three planes pass through

$$\left[\frac{1}{2}\mathbf{u}, \mathbf{u}, 2\mathbf{u} \right],$$

respectively.

□

Problem 2.35. Show that the set

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 < 1\}$$

is open.

Solution. Let $\mathbf{p} = (a, b, c) \in S$ and set

$$\rho = \sqrt{a^2 + b^2} < 1, \quad r = 1 - \rho > 0.$$

If $\mathbf{q} = (x, y, z)$ satisfies $\|\mathbf{q} - \mathbf{p}\| < r$, then by the triangle inequality in the xy -plane,

$$\begin{aligned} \sqrt{x^2 + y^2} &\leq \sqrt{(x - a)^2 + (y - b)^2} + \sqrt{a^2 + b^2} \\ &\leq \|\mathbf{q} - \mathbf{p}\| + \rho \\ &< r + \rho = 1. \end{aligned}$$

Thus $\mathbf{q} \in S$, so $B_r(\mathbf{p}) \subseteq S$. Every point of S is therefore an interior point, and S is open by Definition 2.14.

□

Problem 2.36. Let

$$S = \mathbb{R}^2 \setminus \{\mathbf{0}\}.$$

Show that $\mathbf{0}$ is a boundary point of S .

Solution. Let $r > 0$. The open ball $B_r(\mathbf{0})$ contains $\mathbf{0}$, which is not in S , and it also contains, for example,

$$\left(\frac{r}{2}, 0\right) \in S.$$

Thus every ball centered at $\mathbf{0}$ contains a point of S and a point not in S . By Definition 2.15, $\mathbf{0}$ is a boundary point of S . □

Problem 2.37. Let

$$T = \{(x, y) \in \mathbb{R}^2 \mid x \geq 0, y \geq 0, x + y \leq 1\}.$$

- (a) Describe the boundary of T .
- (b) Show that the point $(0.0001, 0.9998)$ is an interior point of T .

Solution.

- (a) The boundary consists of the three line segments

$$\partial T = \{(0, y) \mid 0 \leq y \leq 1\} \cup \{(x, 0) \mid 0 \leq x \leq 1\} \cup \{(x, y) \mid x \geq 0, y \geq 0, x + y = 1\}.$$

- (b) Let

$$\mathbf{p} = (0.0001, 0.9998), \quad r = 0.00005.$$

If $\mathbf{q} = (x, y)$ satisfies $\|\mathbf{q} - \mathbf{p}\| < r$, then

$$|x - 0.0001| < r, \quad |y - 0.9998| < r.$$

Therefore

$$x > 0.00005 > 0, \quad y > 0.99975 > 0,$$

and

$$x + y < 0.0001 + 0.9998 + 2r = 1.$$

Hence $B_r(\mathbf{p}) \subseteq T$. By Definition 2.13, $\mathbf{p}$ is an interior point of T . □