

Contents

1	Introduction	1
1.1	(Algebraic?) Theory of Anyons	2
1.2	Symmetric Tensors	2
1.3	Review of Symmetric Tensor Networks.	3
1.4	Review of Anyonic Tensor Networks.	4
2	Decomposition of semisimple Categories	4
2.1	Q intertwiners	9
3	Categories	10
3.1	Tensor and Fusion Categories	10
4	Categories Needed	12
4.1	Group Theory	14
4.2	Fusion Trees	14
4.3	Monoidal data	15
5	Anyonic Hamiltonians	15
5.1	Review: The Golden Chain	17
5.2	Rydberg-blockade ladder	17
6	I dont know where to put this yet	17
7	implementation	18

1 Introduction

Lets list some definitions of Tensor. A Tensor T can be regarded as an element of a tensor product space

$$T \in \bigotimes_{j \in J} V^j$$

and we say T is a $|J|$ -legged tensor.

Given finite dimensional Vector spaces V and W , we have the identity that $V \otimes W^* \equiv \text{Hom}(W, V)$. Now, if we have a Tensor $T \in V_1 \otimes \cdots \otimes V_n$, and since for any Vector space V , $(V^*)^* = V$, we can peel off any number of V_i s, take their duals and represent T as a map

$$T : V_n^* \otimes \cdots \otimes V_{k+1}^* \rightarrow V_1 \otimes \cdots \otimes V_k$$

Note that we present the domain here in a contravariant fashion due to the contravariance of the dual.

Lets look at an example, let $t : V_1 \otimes V_2 \otimes W_2^* \otimes W_1^*$. We can represent as an operator in dirac notation:

$$t = \sum_{i_1, i_2, j_1, j_2} T_{j_1 j_2 i_1 i_2} |v_{i_1}\rangle |v_{i_2}\rangle \langle w_{j_1}| \langle w_{j_2}|$$

Graphically, we represent a general tensor

$$T : (W_n^* \otimes \cdots \otimes W_1) \rightarrow (V_1 \otimes V_2 \otimes \cdots \otimes V_n)$$

as follows, where we choose incoming arrows for duals in the domain and outgoings for normal spaces, and on the bottom we choose incoming for duals and outgoing for normals.

1.1 (Algebraic?) Theory of Anyons

1.2 Symmetric Tensors

Let G be a compact group. Consider a tensor T living in the tensor product space of its legs:

$$T \in V_1 \otimes \cdots \otimes V_n \otimes W_m^* \otimes \cdots \otimes W_1^*$$

Let $\rho_i : G \rightarrow GL(V_i)$ be the linear representation of G on each vector space V_i , and let $\sigma_j^* : G \rightarrow GL(W_j^*)$ be the dual representation on W_j^* .

The total action of the group G on the entire tensor product space is given by

$$\Pi : G \rightarrow GL \left(\bigotimes_i V_i \otimes \bigotimes_j W_j^* \right)$$

$$\Pi(g) = \rho_1(g) \otimes \cdots \otimes \rho_n(g) \otimes \sigma_m^*(g) \otimes \cdots \otimes \sigma_1^*(g)$$

A tensor T is defined to be a **symmetric tensor** (or G -invariant tensor) if it is invariant under simultaneous action on all legs for all $g \in G$ [SPV10]:

$$\Pi(g)T = T \quad \forall g \in G$$

By interpreting T as a map

$$T : W_1 \otimes \cdots \otimes W_m \rightarrow V_1 \otimes \cdots \otimes V_n$$


it follows directly from the definition of global invariance that the tensor must commute with the group action and we get the following definition of a symmetric tensor:

Definition 1.1 (symmetric tensor). A *symmetric tensor* is an intertwiner of the representations of the group G :

$$\bigotimes_{j=1}^m \sigma_j(g) T = T \bigotimes_{i=1}^n \rho_i(g) \quad \forall g \in G,$$

which we denote by

$$T \in \text{Hom}_G(W, V).$$



Corollary 1.1.1 (Networks of symmetric tensors are symmetric). Since a tensor network is built by composition and application of the (co)ev maps, both of which are intertwiners, the full network is also an intertwiner. Important there is that this entire discussion is only about FdVect or FdHilb for now, but i think the same applies to any braided category.

1.3 Review of Symmetric Tensor Networks.

Generally, we follow the notations of [SV12], [Sch+20], [Sch+20], [Sch20] which gave an overview for implementing $\text{SU}(2)$ invariant tensor networks. A general overview for (Non)abelian symmetries is [SPV10], and a detailed account for the abelian $\text{U}(1)$ in [SPV11]. There are a multitude of implementations given in major packages, such as ITensor, [FWS22], Cyten, TenPy, [Hau+24], QSpace, [Wei24] ChemTensor, UniTensor, YasTn (Abelian Only) [Ram+25] (MPSKit?), (Cytex (Abelian only i think.))). A general account of symmetries is given in [Unf24] with its respective implementation in [Hau+24] (We have to build cytens citation manually.)

1.4 Review of Anyonic Tensor Networks.

Category	Braiding / Twist	Tensor Network
Hilb_{fd} Monoidal category	No intrinsic braiding. Wire crossings are graphical only.	Ordinary tensor networks. Evaluation:
Braided monoidal category	Braiding $c_{X,Y} : X \otimes Y \rightarrow Y \otimes X$. Crossings carry data.	Crossings generate a braiding factor that needs to be absorbed by the structural tensors.
Symmetric fusion category Rep(G)	$c_{Y,X}c_{X,Y} = 1$. Double exchange is trivial.	Tensor contractions behave much like ordinary TNs. Example: Rep($\mathbb{Z}_2$)
Ribbon / Modular category	Nontrivial braiding: $c_{Y,X}c_{X,Y} \neq 1$. Twist: $\theta_X : X \rightarrow X$.	Anyonic tensor networks. Evaluation may absorb twist: $\widetilde{\text{ev}} = \text{ev} \circ (\theta_X \otimes \text{id})$. Examples: Fibonacci, Ising, $\text{SU}(2)_k$.

2 Decomposition of semisimple Categories

Lemma 2.1. For a vector space W , there is an isomorphism

$$\bigoplus^m W \simeq \mathbb{C}^m \otimes W.$$

Proof. Let $e_{i \in [m]}$ be the standard basis on $\mathbb{C}^m$. Let

$$\Phi : \mathbb{C}^m \otimes W \rightarrow \bigoplus^m W$$

be defined by

$$\phi(e_i \otimes w) = (0, \dots, w, \dots, 0)$$

and linear, thus

$$\Phi \left(\sum_i c_i e_i \otimes w_i \right) = (c_1 w_1, \dots, c_m w_m).$$

is clearly bijective, and we have our desired isomorphism. $\square$

Lemma 2.2. Let D be a finite dimensional k -vector space and let V be an object of a k -linear category C . Then

$$D \otimes V \cong V^{\oplus \dim D}$$

Proof. Let $n = \dim D$. Take a basis $\{e_i\}$ for D . This yields an isomorphism

$$D \cong \bigoplus_{i=1}^n k \cdot e_i \cong k^n$$

Since in a k -linear category the tensor product is additive (should we prove this?), we can distribute

$$D \otimes W \cong \bigoplus_{i=1}^n (k \cdot e_i \otimes W) \cong \bigoplus W = W^{\oplus n}$$

$\square$

Lemma 2.3 (Isotypic Decomposition). Let C be a semisimple k -linear category with simple objects $\{R_a\}$. Then every object $V \in C$ admits a canonical decomposition

$$V \cong \bigoplus_a D_a \otimes R_a. \quad (1)$$

where we refer to $D_a := \text{Hom}_C(R_a, V)$ as the *degeneracy* of the space.

Proof. Since C is semisimple, there exist integers $m_a \geq 0$ such that

$$V \cong \bigoplus_{a \in L} R_a^{\oplus m_a}.$$

Applying the additive functor $\text{Hom}_C(R_b, -)$ yields

$$\text{Hom}_C(R_b, V) \cong \bigoplus_{a \in L} \text{Hom}_C(R_b, R_a^{\oplus m_a}) \cong \bigoplus_{a \in L} \text{Hom}_C(R_b, R_a)^{\oplus m_a}.$$

Then, by Schurs lemma

$$\bigoplus_{a \in L} \text{Hom}_C(R_b, R_a)^{\oplus m_a} = k^{\oplus m_b}$$

$\square$

So we actually need this: Maybe this is simpler if we always let the degeneracy be C ?

Lemma 2.4. Let C be a k -linear category and let D, E be finite-dimensional k -vector spaces. For any objects $R, S \in C$, there is a natural isomorphism

$$\mathrm{Hom}_C(D \otimes R, E \otimes S) \cong \mathrm{Hom}_k(D, E) \otimes \mathrm{Hom}_C(R, S).$$

Proof. Let $m = \dim_k D$ and $n = \dim_k E$. Proof by isomorphism chains:

$$\begin{aligned} \mathrm{Hom}_C(D \otimes R, E \otimes S) &\cong \mathrm{Hom}_C(R^{\oplus m}, S^{\oplus n}) && \text{by 2.1} \\ &\cong \bigoplus_{i=1}^m \bigoplus_{j=1}^n \mathrm{Hom}_C(R, S) && \text{by additivity of } \mathrm{Hom}_C \\ &\cong k^{mn} \otimes \mathrm{Hom}_C(R, S) && \text{by 2.1} \\ &\cong \mathrm{Hom}_k(D, E) \otimes \mathrm{Hom}_C(R, S) && \text{since } \mathrm{Hom}_k(D, E) \cong k^{mn} \end{aligned}$$

(The last as s) □

Theorem 2.5 (Decomposition of Semisimple Categories). Let C be a semisimple, monoidal, k -linear category whose tensor product is k -linear in each variable, with simple objects $\{R_a\}_{a \in L}$ and let V_i, W_i be a collection of objects in C . Every morphism

$$T \in \mathrm{Hom}_C \left(\bigotimes_i V_i, \bigotimes_i W_i \right)$$

admit a decomposition as

$$T \in \bigoplus_{\hat{a}, \hat{b}} P^{(\hat{a}, \hat{b})} \otimes \mathrm{Hom}_C(R_{\hat{a}}, R_{\hat{b}}),$$

where $\hat{a} = (a_1, \dots, a_n)$ and P is a k -vector space of dimension ???

Proof.

$$\begin{aligned}
& \text{Hom}_C \left(\bigotimes_i V_i, \bigotimes_j W_j \right) \\
& \cong \text{Hom}_C \left(\bigotimes_i \left(\bigoplus_{a_i} D_{i,a_i} \otimes R_{a_i} \right), \bigotimes_j \left(\bigoplus_{b_j} E_{j,b_j} \otimes R_{b_j} \right) \right) \quad \text{by 2.3} \\
& \cong \text{Hom}_C \left(\bigoplus_{\hat{a}} D_{\hat{a}} \otimes R_{\hat{a}}, \bigoplus_{\hat{b}} E_{\hat{b}} \otimes R_{\hat{b}} \right) \quad \text{since } \otimes \text{ is } k\text{-linear in each variable} \\
& \cong \bigoplus_{\hat{a}, \hat{b}} \text{Hom}_C(D_{\hat{a}} \otimes R_{\hat{a}}, E_{\hat{b}} \otimes R_{\hat{b}}) \quad \text{by additivity of } \text{Hom}_C \\
& \cong \bigoplus_{\hat{a}, \hat{b}} \text{Hom}(D_{\hat{a}}, E_{\hat{b}}) \otimes \text{Hom}_C(R_{\hat{a}}, R_{\hat{b}}) \quad \text{by 2.4.}
\end{aligned}$$

□

Corollary 2.5.1 (Operators are block matrices). An operator

$$\text{Hom}(V, W) \cong \text{Hom}_C \left(\bigoplus_a D_a \otimes R_a, \bigoplus_b E_b \otimes R_b \right) \cong \bigoplus_{a,b} \text{Hom}(D_a, E_b) \otimes \text{Hom}_C(R_a, R_b)$$

and by Schurs lemma this is just

$$\bigoplus_{a \in \text{Irr}(V) \cap \text{Irr}(W)} \text{Hom}(D_a, E_a)$$

where $\text{Hom}(D_a, E_a)$ is a matrix. Lets say that a is in a collection of isomorphisms L . Then (show matrix here ...)

Corollary 2.5.2. Let $Q_{\mu, \eta}$ be a basis of $\text{Hom}_C(\bigotimes_{\hat{a}} R_{\hat{a}}, \bigotimes_{\hat{b}} R_{\hat{b}})$ where μ denotes the multiplicities and η the internal fusions. In a *multiplicity-free* Category we will omit μ as a label.

Example 1. If we let our monoidal semisimple category just be Vec_C , the only simple object is C . Thus there is also only the trivial biproduct over one sector.

Example 2. If we restrict to the category $\text{Rep}(\text{SU}(2))$ we can build the symmetric tensor networks as given in [SV12]. TODO: actual ly do this and convince yourself that the notations amd the triplet basis workd as desired.

Example 3. Take a 2 in 1 out tensor in $\text{Rep}(\text{SU}(2))$ as

$$T \in \text{Hom}_{\text{Rep}(\text{SU}(2))}(V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}, V_{\frac{1}{2}})$$

Since all parts are simple, nothing happens.

Example 4. However, now assign $V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}$ to each leg in the example above. Each leg then becomes $V_0 \oplus V_1 \cong \mathbb{C} \otimes V_0 \oplus \mathbb{C} \otimes V_1$. Then we can start with the decomposition.

$$\begin{aligned} & \text{Hom}_{\mathbb{C}}\left(\bigoplus_{a,b,c \in (0,1)} (\mathbb{C} \otimes \mathbb{C} \otimes \mathbb{C}) \otimes V_a \otimes V_b \otimes V_c, \bigoplus_{b \in (0,1)} \mathbb{C} \otimes V_b\right) \\ &= \bigoplus_{a,b,c,d \in (0,1)} \text{Hom}_{\mathbb{C}}((\mathbb{C} \otimes \mathbb{C} \otimes \mathbb{C}) \otimes (V_a \otimes V_b \otimes V_c), \mathbb{C} \otimes V_d) \\ &= \bigoplus_{a,b,c,d} \text{Hom}(\mathbb{C}^3, \mathbb{C}) \otimes \text{Hom}_{\text{Rep}(\text{SU}(2))}(V_a \otimes V_b \otimes V_c, V_d) \end{aligned}$$

Example 5. Easier. However, now assign $V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}$ to each leg in the example above. Each leg then becomes $V_0 \oplus V_1 \cong \mathbb{C} \otimes V_0 \oplus \mathbb{C} \otimes V_1$. Then we can start with the decomposition.

$$\begin{aligned} & \text{Hom}_{\mathbb{C}}\left(\bigoplus_{a,b \in (0,1)} (\mathbb{C} \otimes \mathbb{C}) \otimes V_a \otimes V_b, \bigoplus_{b \in (0,1)} \mathbb{C} \otimes V_b\right) \\ &= \bigoplus_{a,b,c \in (0,1)} \text{Hom}_{\mathbb{C}}((\mathbb{C} \otimes \mathbb{C}) \otimes (V_a \otimes V_b), \mathbb{C} \otimes V_c) \\ &= \bigoplus_{a,b,c} \text{Hom}(\mathbb{C}^2, \mathbb{C}) \otimes \text{Hom}_{\text{Rep}(\text{SU}(2))}(V_a \otimes V_b, V_c) \end{aligned}$$

Example 6. Consider the same operator as in example 5 but with its incoming legs fused together. So we have $V_{\text{in}} = V_{\frac{1}{2}}^4$ and $V_{\text{out}} = V_{\frac{1}{2}}^2$. This decomposes $V_2 \oplus V_1^{\oplus 3} \oplus V_0^{\oplus 2}$. Then we can directly use 2.5.1 to obtain

$$\text{Hom}(V_{\text{in}}, V_{\text{out}}) =$$

Example 7. Lets look at a tensor with in legs $(\tau \otimes \tau)$ and τ and out legs τ .

$$\begin{aligned} & \text{Hom}_{\mathbb{C}}(((k \otimes 1) \oplus (k \otimes \tau)) \otimes (k \otimes \tau), k \otimes \tau) = \text{Hom}_{\mathbb{C}}(\\ & \text{Hom}_{\mathbb{C}}(((k \otimes 1) \oplus (k \otimes \tau)) \otimes (k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}_{\mathbb{C}}((k \otimes 1 \otimes k \otimes \tau) \oplus (k \otimes \tau \otimes k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}_{\mathbb{C}}(k \otimes 1 \otimes k \otimes \tau, k \otimes \tau) \oplus \text{Hom}_{\mathbb{C}}(k \otimes \tau \otimes (k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}(k \otimes k, k) \otimes \text{Hom}_{\mathbb{C}}(1 \otimes \tau, \tau) \oplus \text{Hom}(k \otimes k, k) \otimes \text{Hom}_{\mathbb{C}}(\tau \otimes \tau, \tau). \end{aligned}$$

Example 8 (Anyonic matrix operator). Finally, consider a $4 \rightarrow 4$ anyon operator with one leg on each side, representing the space $\text{Hom}((\tau \otimes \tau \otimes \tau \otimes \tau), (\tau \otimes \tau \otimes \tau \otimes \tau))$. The full decomposition is $2 + 3\tau$. Hence we obtain

$$\begin{aligned} & \text{Hom}_{\mathbb{C}}\left((k^2 \otimes 1) \oplus (k^3 \otimes \tau), (k^2 \otimes 1) \oplus (k^3 \otimes \tau)\right). \\ &= \bigoplus_{a,b \in (1,\tau)} \text{Hom}_{\mathbb{C}}(k^{m_a} \otimes a, k^{m_b} \otimes b) = \bigoplus_{a,b \in (1,\tau)} \text{Hom}(k^{m_a}, k^{m_b}) \otimes \text{Hom}_{\mathbb{C}}(a, b) \end{aligned}$$

where

$$m_1 = 2, \quad m_\tau = 3.$$

For this case, Schurs lemma applies. So we have

$$= \text{Hom}(k^2, k^2) \oplus \text{Hom}(k^3, k^3)$$

which is just a block matrix with indexing $(1_1, 1_2, \tau_1, \tau_2, \tau_3)$.

Example 9 ($U(1)$ Symmetry). Technically we would want to show here that this is simpler in general. TOOD. Make a theorem? Or a corollary?

Example 10 (Ground state on 4 anyons). We can be given a ground state in the operator form as $\text{Hom}(\tau^4, \tau)$.

2.1 Q intertwiners

Lets look at 2 paths for the 4 index tensor, $(a, b, c \rightarrow d)$ and $(a, b \rightarrow c, d)$.

$$(Q_{j_a j_b j_c}^{j_e})_{m_1, m_2, m_3, m_4} = \sum_{m_e} \langle j_1 m_1; j_2 m_2 | j_e m_e \rangle \langle j_e m_e; j_3 m_3 | j_4 m_4 \rangle$$

while the other topology would look like

$$Q \dots = \sum_{m_e} \langle j_1 m_1; j_2 m_2 | j_e m_e \rangle \langle j_e m_e | j_3 m_3; j_4 m_4 \rangle$$

and as an operator this would have to live in $\text{Hom}(V_{j_1} \otimes V_{j_2}, V_{j_3} \otimes V_{j_4})$

The yoga tree $(-1, 1, -3), (-2, 1, -4)$ Gives us a good mixture of those. This Q would then have to look like

$$Q_{-1-2-3-4}^1 = \sum_1 \langle j_{-1} m_{-1}; j_1 m_1 | j_{-3} m_{-3} \rangle \langle j_{-2} m_{-2} | j_1 m_1; j_{-4} m_{-4} \rangle = \sum_{m_1} C_{-1,1 \rightarrow -3} \cdot C_{-2,1 \rightarrow -4}$$

How would we do this when we have a contraction of two simultaneous indices:

$$Q_{-1-2}^{1,2} = \sum_{1,2} \langle -1|1;2 \rangle \langle 1;2|-2 \rangle$$

If we define τ to be the fusion trees topology, then we can define ${}_{\tau}Q_j^i$

What would this look like for general symmetry groups G ? Lets choose a basis of its invariant decomp $|\lambda_a \mu_a \rangle$. Then the generalized CG coefficient would look like

$$(C_{\lambda_a \lambda_b \rightarrow \lambda_c}^{\alpha})_{\mu_a \mu_b}^{\mu_c} = \langle \lambda_a \mu_a; \lambda_b \mu_b | \lambda_c \mu_c \rangle_{\alpha}$$

where alpha denotes the singular fusion multiplicity at μ_c .

Now to fully specify Q here for k sectors (λ_i, μ_i) , we need to give it $k - 3$ internal edges $(\lambda_{e_1}, \dots, \lambda_{e_{k-3}})$ along with $k - 2$ mulitplicity symbols, $(\alpha_1, \dots, \alpha_{k-2})$

Can we write down a general formula for Q ?

This would then be

$$Q_{\lambda}$$

3 Categories

3.1 Tensor and Fusion Categories

Let C be a category with bifunctor

$$\otimes : C \times C \rightarrow C, \quad (X, Y) \rightarrow X \otimes Y.$$

and the isomporhism $F : (- \otimes -) \otimes - \simeq - \otimes (- \otimes -)$

... introduce fusion categories here, with associativity and commutativity restrictions, I think this would be a good start.

Definition 3.1 (Monoidal Category). A monoidal category is a category C with a bifunctor $\otimes : C \times C \rightarrow C$

Let C and D be linear abelian categories. Then the *Deligne Tensor Product* $C \boxtimes D$ is a linear abelian category with a bifunctor

$$\boxtimes : C \times D \rightarrow C \boxtimes D$$

with morphisms given by

$$\text{Hom}_{C \boxtimes D}(X \boxtimes Y, W \boxtimes V) = \text{Hom}_C(X, W) \otimes \text{Hom}_D(Y, V)$$

Definition 3.2 (The Deligne Tensor Product of Categories). Let k be a field, and let $\mathcal{C}$ and $\mathcal{D}$ be two k -linear, finite abelian categories. The **Deligne tensor product**, denoted by $\mathcal{C} \boxtimes \mathcal{D}$, is a k -linear finite abelian category equipped with a k -bilinear functor

$$\boxtimes : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{C} \boxtimes \mathcal{D}, \quad (X, Y) \mapsto X \boxtimes Y$$

TODOTOD!!

Definition 3.3 (The Category Vec_k). Let k be an algebraically closed field.

- **Objects:** The objects are finite dimensional vectors spaces over k .
- **Morphisms:** Linear maps between vector spaces.

Definition 3.4 (The Category $\text{Rep}_k(G)$). Let G be a group and k a field. The *category of linear representations of G over k* consists of the following data:

- **Objects:** The objects $\text{Rep}_k(G)$ are pairs (V, ρ_V) , where V is a k -vector space and $\rho_V : G \rightarrow \text{GL}(V)$.
- **Morphisms:** For any two objects (V, ρ_V) and (W, ρ_W) in $\text{Rep}_k(G)$, a morphism from V to W is a k -linear transformation $f : V \rightarrow W$ that is G -invariant (also called an *intertwiner*). That is, for every element $g \in G$, the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\rho_V(g)} & V \\ f \downarrow & & \downarrow f \\ W & \xrightarrow{\rho_W(g)} & W \end{array}$$

In algebraic terms, f must satisfy the condition:

$$f \circ \rho_V(g) = \rho_W(g) \circ f \quad \forall g \in G$$

The set of all such morphisms is denoted by $\text{Hom}_G(V, W)$.

We can further endow $\text{Rep}_k(G)$ with a monoidal structure: For any two objects $(V, \rho_V), (W, \rho_W)$, their tensor product is in $\text{Rep}_k(G)$ by the diagonal action

$$\rho_{(V \otimes W)}(g)(v \otimes w) = \rho_V(g)v \otimes \rho_W(g)w$$

Definition 3.5 (Direct product of representations). A representation of $G_1 \times G_2$ is a space where the pair (g_1, g_2) acts linearly. SO this is a pair

$$(V, \rho), \rho : (G_1 \times G_2) \rightarrow \text{GL}(V)$$

Now, let $(\text{Rep}_{\mathbb{C}}(G), \otimes, \mathbb{C})$ be the monoidal category of representations. We equip the category with a tensor product that acts as follows:

TODO....

Theorem 3.6 (Fusion Representation Category of finite groups). Let G be a finite group. Then $\text{Rep}(G)$ is a fusion category.

Theorem 3.7. Let G_1 and G_2 be finite groups and let $\text{Rep}(G_1), \text{Rep}(G_2)$ be their respective symmetry categories. Then there is an equivalence of categories

$$\text{Rep}(G_1 \times G_2) \simeq \text{Rep}(G_1) \boxtimes \text{Rep}(G_2)$$

Big issue here: $\text{Rep}(\text{SU}(2))$ is not a fusion category as it is not finite. But if i read it correctly, the deligne product only requires abelianness of its groups. Now, $\text{Rep}(G)$ is monoidal for any G but is it abelian?

So this is actually true, it is abelian. We could obtain a restricted versions by for example always restricting $\text{SU}(2)$ to some $\text{SU}(2)_k$ for some finite k . But i do not yet know how to present this nicely.

A second advantage would be that we can always represent fusion trees as monoidal categories (is this true or is fusion required here again?), and we can write down a unified theory for the fusion trees. Then I hope to write a nice method to view the "eval" of such a fusion tree, i.e. its clebsch gordan tensors by contracting down the topology of the tree.

4 Categories Needed

The motivation for this section is to introduce the necessary category theory background, and provide all the necessary theorems for a format that includes both representations and fusion categories.

Definition 4.1 (Monoidal Category). A monoidal category is a category C equipped with

1. A bifunctor $\otimes : C \times C \rightarrow C$.
2. A unit element $1 \in C$
3. An isomorphism

$$\alpha : - \otimes (- \otimes -) \rightarrow (- \otimes -) \otimes -$$

4. Left and right unitors $\gamma : 1 \otimes - \rightarrow -$ and $\delta : - \otimes 1 \rightarrow -$.

We denote this by the triple $(C, \otimes, 1)$ where we omit the unitors.

Definition 4.2 (Semisimple Category). An abelian category $\mathcal{C}$ is semisimple if every object is either simple or a direct sum of simple objects. An object is *simple* if its only subobjects are 0 or X itself. Any object can be written as

$$X = \bigoplus_{i \in I} S_i^{\oplus n_i}$$

Definition 4.3 (Abelian Category).

Lemma 4.4 (Schurs 1st lemma (which one?)). If S and T are simple objects in a semisimple category over an algebraically closed field (this is always $\mathbb{C}$ in our case), then it holds that

$$\text{Hom}(S, T) = \begin{cases} k, & \text{if } S \cong T; \\ 0, & \text{if } S \not\cong T. \end{cases}$$

Proof. Step 1. Let $f : S \rightarrow T$ be a morphisms. Since T is simple

$$\text{im}(f) \cong 0 \quad \text{or} \quad \text{im}(f) \cong T$$

□

Lets give a coordinatized approach to fusion categories:

Definition 4.5 (Fusion Category). A *fusion category* is a finite set of labels (sectors?) L and vector spaces V_c^{ab} for all $a, b, c \in L$, as well as the isomorphisms

$$F_d^{abc} : \bigoplus_e V_e^{ab} \otimes V_d^{ec} \rightarrow \bigoplus_e V_d^{ae} \otimes V_e^{bc}$$

such that the pentagon holds.

Lemma 4.6. Let $\mathcal{C}$ be a k -linear monoidal category whose tensor product is k -linear in each variable. Then

$$\bigotimes_{r=1}^n \left(\bigoplus_{a_r} X_{r, a_r} \right) \cong \bigoplus_{\hat{a}} \bigotimes_{r=1}^n X_{r, a_r},$$

where

$$\hat{a} = (a_1, \dots, a_n).$$

Lemma 4.7 (Finite Groups admit Fusion Category). Let G be a finite group, the $\text{Rep}(G)$ is a fusion category.

Let $\text{Vec} := (\text{FDCVec}, \otimes, \mathbb{C})$ be the monoidal category of finite dimensional complex vector spaces where the last tuple entry $\mathbb{C}$ denotes the unit object, since there are isomorphisms $V \otimes \mathbb{C} \cong V \cong \mathbb{C} \otimes V$.

Let $(\mathbb{C}\text{Rep}(G), \otimes, ?)$ denote the representation Category for a compact group G . Here $\otimes$ is the tensor product of representations $(\pi \otimes \rho, V \otimes W)$.

Definition 4.8 (Direct sum). Given a category C with objects $\{a_n\}$, an object $\bigoplus_{i=1}^n a_i \in C$ is called the *direct sum*. This is equipped with the injections and projections

$$\iota_i : a_i \rightarrow \bigoplus a_i \quad \text{and} \quad \pi_i : \bigoplus a_i \rightarrow a_i$$

such that

$$\pi_i \circ \iota_i = \text{id}_{a_i} \forall i$$

Lemma 4.9 (Peter Weyl Theorem). Let G be a compact group. Then all its unitary representations are completely reducible.

4.1 Group Theory

Definition 4.10 (Compact Group). A compact group is a group G equipped with a topology such that

$$m : G \times G \rightarrow G, \quad (g, h) \mapsto gh,$$

is continuous.

$$i : G \rightarrow G, \quad g \mapsto g^{-1}$$

is continuous and G is compact as a topological space.

Example 11. 1. Every finite group with the discrete topology is compact.

2.

$$U(n) = \{A \in M_n(\mathbb{C}) : A^* A = I\}$$

is compact for all integer n .

3. All

$$SU(n) = \{A \in U(n) : \det A = 1\}$$

are compact.

4.2 Fusion Trees

We consider a fusion tree as the basis that spans $\text{Hom}(R_a \otimes R_b \dots \otimes R_m, \otimes R_{a_j} \dots)$. Since $R_a \otimes R_b$ is semisimple, we have the relationship

$$R_a \otimes R_b \cong \bigoplus_c R_c^{\oplus N_{ab}^c}$$

with embedding (make this formal)

$$g_{c,\mu}^{ab} : R_c \rightarrow R_a \otimes R_b$$

and projections

$$f_{ab}^{c,\mu} : R_a \otimes R_b \rightarrow R_c$$

$$\text{id}_{R_a \otimes R_b} = \sum_{c \in L} \sum_{\mu=1}^{N_{ab}^c} g_{c,\mu}^{ab} f_{ab}^{c,\mu}$$

Lemma 4.11.

$$\text{Hom}(R_{a_1} \otimes \cdots \otimes R_{a_N}, R_b)$$

is spanned by ordinary fusion trees.

Proof. Base case: $N = 1$. Nothing to prove here.

INDuctive case:

Under semisimplicity we get that

$$\text{Hom}(R_{a_1} \otimes \cdots \otimes R_{a_n}, R_b) \cong \bigoplus_{c,\mu} \text{Hom}(R_c \otimes R_{a_3} \otimes \cdots \otimes R_{a_n}, R_b)$$

□

4.3 Monoidal data

Definition 4.12 ($\text{Rep}(\text{SU}(2))$). Let (V, ρ_V) be a representation.

Definition 4.13 (Twists from R moves). Suppose we are in a braided monoidal category with F and R symbols. Then we compute the *twist* (a self braiding) as follows:

$$R_c^{ab} R_c^{ba} = \frac{\theta_c}{\theta_a \theta_b}.$$

(state more possibilities here.)

Example 12. Take $s\text{Vect}$. Our simples are 1 and f . This *braiding* is in fact symmetric. Thus we have that

$$1 = \frac{\theta_1}{\theta_f^2} =$$

5 Anyonic Hamiltonians

Definition 5.1 (XXZ-Hamiltonian). The XXZ Hamiltonian is

$$H = J \cdot \sum_{i=1}^N (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_{i+1}^z)$$




Figure 1: XXZ Hamiltonian in both bases.

We can plot H and realize the sparsity structure thereof. Furthermore, a suitable rearrangement induces a block structure [1](#).

Now, we can realize about this hamiltonian is that there is a symmetry with the representations of $U(1)$. Lets review this shortly. A representation of $U(1)$ is a map

$$\rho : U(1) \rightarrow GL(V), \quad \rho(e^{i\theta_1})\rho(e^{i\theta_2}) = \rho(e^{i(\theta_1+\theta_2)})$$

The XXZ model lives in the Hilbert Space $\mathcal{H} = (C^2)^n$ and the relevant representation is generated by the total magnetization operator

$$S_{\text{tot}}^z = \frac{1}{2} \sum_{i=1}^N \sigma_i^z$$

The group action is given by

$$U(\theta) = e^{-i\theta S_{\text{tot}}^z}.$$

We now claim that H is invariant under $U(\theta)$.

$$U(\theta) H U(\theta)^\dagger = H, \quad \forall \theta \in \mathbb{R}.$$

Equivalently,

$$[H, S_{\text{tot}}^z] = 0.$$

Recall that this last condition entails a block diagonal basis for H , as H and S^z are simultaneously diagonalizable.

5.1 Review: The Golden Chain

For the golden chain, we consider $n + 2$ Fib anyons as our ground state. Their combined state yields a Fusion Tree $\text{Hom}(\tau^{\otimes n+1}, \tau)$ with intermediate charges $x_1, \dots, x_n$. These intermediate charges are further restricted as in that there are no possibilities to assign adjacent 1s to them (In fact this is what causes the dimension of the hilbert space to reduce to ϕ^n).

There is a three local hamiltonian for the golden chain defined as

$$H_i = - \sum_{x'_i} [F_{x_{i+1}}^{x_{i-1}\tau\tau}]_{x_i}^{x'_i} E_{x'_i} [F_{x_{i+1}}^{x_{i-1}\tau\tau}]_{x_i'}^{x''_i}$$

Write this three local hamiltonian in an operator form

$$H_i = -J \begin{pmatrix} 1 & & & & \\ & 0 & & & \\ & & 0 & & \\ & & & \varphi^{-2} & \varphi^{-\frac{3}{2}} \\ & & & \varphi^{-\frac{3}{2}} & \varphi^{-1} \end{pmatrix}$$

Note that this is block diagonal over the fusion channel (TODO: change notation).

5.2 Rydberg-blockade ladder

This is given by the fusion category $\text{so}(3)_2$. Here we have fusion rules

$$1 \otimes 1 = 0 \oplus 1 \oplus 2, \quad 2 \otimes 1 = 1, \quad 2 \otimes 2 = 0.$$

6 I dont know where to put this yet

If we start out on anyonic chains, like the $\text{SU}(2)_3$ case, this provideds a nice generalization for chains with $C = \text{Rep}(\text{SU}(2)_k)$ with fusion rules

$$j_1 \times j_2 = \sum_{j=|j_1-j_2|}^{\min(j_1+j_2, k-j_1-j_2)} j.$$

However now, if we approach this in the case of $\text{SU}(2)_\infty$ these fusion rules read

$$j_1 \times j_2 = \sum_{j=|j_1-j_2|}^{j_1+j_2}$$

but this chain would have an infinite Hilber Space? Also, lifting to representation operators reveals that there are infinitely many that are not independent.

7 implementation

We want to be able to represent a traditional tensor op such as

```
1 np.einsum("ijklmn,nkol -> ijmo", a, b)
```

for symmetric tensors. As we want to reuse code form existing librariries that is written using numpy, it would be ideal to write our classes in a way that would only require substituting np for our own libraries snp. Towards this goal we define a abstractclass Tensor

```
1 class Tensor(abc.ABC):
2     ...
3     @abstractmethod
4     def transpose(cls, new_shape:list[int]): ...
5     ...
```

and we implement both an np wrapper NPTensor and our symmetric Tensor as STensor, i.e.

```
1 class STensor(Tensor):
2
3     def __init__(self, ...):
4         self.open_legs:list[int|str]
5         self.fusion_tree:FusionTree(self.open_legs, ...)
6         ...
7     @abstractmethod
8     def transpose(cls, new_shape:list[int]): ...
9     ...
```

References

- [FWS22] Matthew Fishman, Steven R. White, and E. Miles Stoudenmire. *The ITensor Software Library for Tensor Network Calculations*. In: *SciPost Phys. Codebases* (2022), p. 4. DOI: [10.21468/SciPostPhysCodeb.4](https://doi.org/10.21468/SciPostPhysCodeb.4). URL: <https://arxiv.org/abs/2007.14822>.
- [Hau+24] Johannes Hauschild et al. *Tensor Network Python (TeNPy) Version 1*. In: *SciPost Phys. Codebases* (2024), p. 41. DOI: [10.21468/SciPostPhysCodeb.41](https://doi.org/10.21468/SciPostPhysCodeb.41). URL: <https://scipost.org/10.21468/SciPostPhysCodeb.41>.
- [Ram+25] Marek M. Rams, Gabriela Wójtowicz, Aritra Sinha, and Juraj Hasik. *YASTN: Yet Another Symmetric Tensor Networks; A Python Library for Abelian Symmetric Tensor Network Calculations*. In: *SciPost Phys. Codebases* (2025), p. 52. DOI: [10.21468/SciPostPhysCodeb.52](https://doi.org/10.21468/SciPostPhysCodeb.52). URL: <https://scipost.org/10.21468/SciPostPhysCodeb.52>.

- [Sch+20] Philipp Schmoll, Sukhbinder Singh, Matteo Rizzi, and Román Orús. *A Programming Guide for Tensor Networks with Global $SU(2)$ Symmetry*. In: *Annals of Physics* 419 (Aug. 1, 2020), p. 168232. DOI: [10.1016/j.aop.2020.168232](https://doi.org/10.1016/j.aop.2020.168232). URL: <https://www.sciencedirect.com/science/article/pii/S0003491620301664> (visited on 05/07/2026).
- [Sch20] Philipp Schmoll. *Tensor Network Simulations with Global $SU(2)$ Symmetry*. Mainz: Johannes Gutenberg University, July 2020. 150 pp. URL: <https://openscience.ub.uni-mainz.de/server/api/core/bitstreams/d9d9ca4e-46c8-4a8e-a831-7611475d2504/content> (visited on 05/08/2026).
- [SPV10] Sukhwinder Singh, Robert N. C. Pfeifer, and Guifre Vidal. *Tensor Network Decompositions in the Presence of a Global Symmetry*. In: *Physical Review A* 82.5 (Nov. 8, 2010), p. 050301. DOI: [10.1103/PhysRevA.82.050301](https://doi.org/10.1103/PhysRevA.82.050301). arXiv: [0907.2994](https://arxiv.org/abs/0907.2994) [cond-mat.str-el]. URL: <http://arxiv.org/abs/0907.2994> (visited on 05/11/2026).
- [SPV11] Sukhwinder Singh, Robert N. C. Pfeifer, and Guifre Vidal. *Tensor Network States and Algorithms in the Presence of a Global $U(1)$ Symmetry*. In: *Physical Review B* 83.11 (Mar. 15, 2011), p. 115125. DOI: [10.1103/PhysRevB.83.115125](https://doi.org/10.1103/PhysRevB.83.115125). URL: <https://arxiv.org/abs/1008.4774> (visited on 05/06/2026).
- [SV12] Sukhwinder Singh and Guifre Vidal. *Tensor Network States and Algorithms in the Presence of a Global $SU(2)$ Symmetry*. In: *Physical Review B* 86.19 (Nov. 9, 2012), p. 195114. DOI: [10.1103/PhysRevB.86.195114](https://doi.org/10.1103/PhysRevB.86.195114). URL: <https://arxiv.org/abs/1208.3919v1> (visited on 05/11/2026).
- [Unf24] Jakob Unfried. *Advancing Tensor Network Simulations* -. PhD thesis. Technische Universität München, 2024, p. 208. URL: <https://mediatum.ub.tum.de/1755273>.
- [Wei24] Andreas Weichselbaum. *QSpace - An Open-Source Tensor Library for Abelian and Non-Abelian Symmetries*. In: *SciPost Physics Codebases* (Nov. 14, 2024), p. 40. DOI: [10.21468/SciPostPhysCodeb.40](https://doi.org/10.21468/SciPostPhysCodeb.40). URL: <https://scipost.org/10.21468/SciPostPhysCodeb.40> (visited on 05/08/2026).