

Contents

1	Overview	3
1.1	Symmetries in Physics	3
1.2	Review: Landau Theory and it's limitations	4
1.3	Introduction	4
1.3.1	Some basic Category Theory	4
1.3.2	Tensors (Everything here is fdVect_k)	5
1.3.3	(Algebraic?) Theory of Anyons	6
1.3.4	Symmetric Tensors	6
1.3.5	Some Important Facts about Hamiltonians	7
1.3.6	Review of Symmetric Tensor Networks.	7
2	Decompositions for Categorical Networks	11
2.1	Decomposition of semisimple Categories	12
2.1.1	Q intertwiners	16
2.2	Categories	17
2.2.1	Tensor and Fusion Categories	17
2.3	Categories Needed	18
2.3.1	Representation Theory	20
2.3.2	Fusion Trees	20
2.3.3	Monoidal data	21
2.4	Anyonic Hamiltonians	21
2.4.1	1D Lattice Models	22
2.4.2	Rydberg-blockade ladder	23
2.5	I dont know where to put this yet	23
2.6	MPO-symmetries	24
2.7	implementation	24

Chapter 1

Overview

1.1 Symmetries in Physics

What do i even want to do here? I want to give an overview on the role of symmetries in condensed matter and many body physics that is also of interest to computer scientists, i.e. describe some sort of system to compute with. However also, there should be an overview about the flavours of (topological) phases, orders and entanglements. Most important foremost would be a paragraph on what "symmetry" as a concept refers to. Then we would want to classify the symmetries we can (or want to!) address with tensor networks (and string (spin?) networks!). Landau theory and its failure would be a great starting point for a condensed matter approach, this is directly correlated with all of topological order, symmetric hamiltonians and dualities.

So for now im thinking to treat this as a collection of independent chapters. First on global symmetries in tensor networks. Work backwards from the singh papers (however these are actually post-anyon networks (c.f. [Pfe+10])) and establish the extent that these papers deal with.

Then write one section on the anyonic networks with sparse mentions of symmetries. Of course we could give representations as a special case here if we take the representations to form an anyon theory (really just a fusion category) as given in [Sim23, § 20] but I do not know if this is desired. Well, the general theory is handled in [DH25] with utmost care, however I would like to give some duly needed proofs (in really all of these papers, I believe the word *proof* only appears once in all of them!) i.e. as done already with our decomposition theorem 2.1.5.

And then it really is just open ended. I think a nice description of dualities is duly manageable but this of course requires the discussion of module categories as in [LDV24a] as well as a restriction on the type of systems we deal with. Another option would be to explore MPO-symmetries, as these share quite a lot of language with our decompositions (in fact, the symmetries they generate are a direct superset of ours!). There is also the consideration of virtual only symmetries, and I currently remember a description of them here [dGro22]. Both the MPO-case and the virtual symmetry case require a discussion of projective symmetries as well.

Sometime I currently know nothing about is the theory on *symmetry protected topological* (SPT) phases, but it does sound relevant. [dGro22] also gives some tensor network methods in this context.

The topic of topological order would require a discussion of string nets, and inthereof the toric

code. I need to understand this anyways, so why not write it down nicely?

1.2 Review: Landau Theory and it's limitations

All matter is made of (what exactly?), but can display many different structural properties called *phases*, such as liquid, crystal, solid, (what else? condensations? is there a nice diagram?). These arrangements of matter can be manipulated, should the structural property change we say that a *phase transition* has occurred.

Landaus *symmetry breaking* theory [**<empty citation>**] was thought to classify all phases of matter (What are the 230 Groups/Crystals here?). However, as shown by [Wen95], there are phases that are not described by spontaneous symmetry breaking but rather a new quantity, the dependence on the topology of the ground state.

There is a nice review here [Wen13].

1.3 Introduction

1.3.1 Some basic Category Theory

Here we introduce some basic category concepts. For a review of categories for quantum systems see [Sel11], for a use in tensor networks see [VW18] or [Bia20](Check this!).

Similarly to the concepts above, we introduce categories because there are some that mirror the graphical language used by tensor networks exactly. Furthermore, both our later treatment of anyons and symmetries in Chapters makes use of these concepts. Here we mostly follow the language used by [Vak25].

Definition 1.3.1 (Category). A category C is built by a collection of objects, which we usually denote by $\text{obj}(C)$, and a set of morphisms $\text{Hom}_C(A, B)$ for each pair A, B of objects, where we refer to A as the domain and B as the codomain. If we want to refer to an object in C , we usually just write $A \in C$.

Furthermore, there is a binary operator $\circ : \text{Hom}_C(A, B) \times \text{Hom}_C(B, C) \rightarrow \text{Hom}(A, C)$ such that this is associative. Also, we have an identity morphism for each object A denoted by id_A s.t. $\text{id}_A \circ f = f$ and $g \circ \text{id}_B = g$ for all appropriate morphisms.

We say there is an *isomorphism* between two objects if there exists an inverse morphism (Just copy vakil here.).

As an example Hilb_{fd} forms a category where the objects are f.d. Hilber spaces and the morphisms are the linear maps between them.

Realize that this encodes all the needed information about sequential composition.

Definition 1.3.2 (Functor). A *Functor* is a map from category A to category B and is given by the data

$$F : \text{obj}(A) \rightarrow \text{obj}(B) \quad \text{and} \quad F : \text{Hom}(A, B) \rightarrow \text{Hom}(F(A), F(B))$$

such that F preserves the identity and composition

$$F(\text{id}_A) = \text{id}_{F(A)} \quad \text{and} \quad F(f \circ g) = F(f) \circ F(g)$$

Now, in a quantum system we want the option to also describe *parallel composition*. If we have two systems A and B we want to apply processes f and g on the combined system $A \otimes B$ as $f \otimes g$. So we introduce extra structure to our categories in the form of a functor $-\otimes-$. This acts between

$$\otimes : C \times C \rightarrow C$$

where

$$\otimes : \text{obj}(C) \times \text{obj}(C) \rightarrow C : (a, b) \mapsto a \otimes b$$

and

$$\otimes : \text{Mor}(A, A') \times \text{Hom}_C(B, B') \rightarrow \text{Hom}_C(A \otimes A', B \otimes B'), (f, g) \mapsto f \otimes g.$$

Now, the tensor product in general is not associative, i.e. the spaces $A \otimes (B \otimes C)$ and $(A \otimes B) \otimes C$ are only isomorphic. So we further define a natural isomorphism

$$\alpha : (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C)$$

called the associator.

We can now make this construction a formal definition:

Definition 1.3.3 (Monoidal Category). A monoidal category is a category C equipped with

1. A bifunctor $\otimes : C \times C \rightarrow C$.
2. A unit element $1 \in C$
3. An isomorphism

$$\alpha : - \otimes (- \otimes -) \rightarrow (- \otimes -) \otimes -$$

4. Left and right unitors $\gamma : 1 \otimes - \rightarrow -$ and $\delta : - \otimes 1 \rightarrow -$.

We denote this by the triple $(C, \otimes, 1)$ where we omit the unitors. Otherwise this is a 5 or 6 tuple
TODO.

TODO: graphical calculus.

1.3.2 Tensors (Everything here is fdVect_k)

Lets list some definitions of Tensor. A Tensor T can be regarded as an element of a tensor product space

$$T \in \bigotimes_{j \in J} V^j$$

and we say T is a $|J|$ -legged tensor.

Given finite dimensional Vector spaces V and W , we have the identity that $V \otimes W^* \equiv \text{Hom}(W, V)$. Now, if we have a Tensor $T \in V_1 \otimes \cdots \otimes V_n$, and since for any Vector space V , $(V^*)^* = V$, we can peel off any number of V_i s, take their duals and represent T as a map

$$T : V_n^* \otimes \cdots \otimes V_{k+1}^* \rightarrow V_1 \otimes \cdots \otimes V_k$$

Note that we present the domain here in a contravariant fashion due to the contravariance of the dual.

Lets look at an example, let $t : V_1 \otimes V_2 \otimes W_2^* \otimes W_1^*$. We can represent as an operator in dirac notation:

$$t = \sum_{i_1, i_2, j_1, j_2} T_{j_1 j_2 i_1 i_2} |v_{i_1}\rangle |v_{i_2}\rangle \langle w_{j_1}| \langle w_{j_2}|$$

Graphically, we represent a general tensor

$$T : (W_n^* \otimes \cdots \otimes W_1) \rightarrow (V_1 \otimes V_2 \otimes \cdots \otimes V_n)$$

as follows, where we choose incoming arrows for duals in the domain and outgoings for normal spaces, and on the bottom we choose incoming for duals and outgoing for normals.

1.3.3 (Algebraic?) Theory of Anyons

1.3.4 Symmetric Tensors

Let G be a compact group. Consider a tensor T living in the tensor product space of its legs:

$$T \in V_1 \otimes \cdots \otimes V_n \otimes W_m^* \otimes \cdots \otimes W_1^*$$

Let $\rho_i : G \rightarrow GL(V_i)$ be the linear representation of G on each vector space V_i , and let $\sigma_j^* : G \rightarrow GL(W_j^*)$ be the dual representation on W_j^* .

The total action of the group G on the entire tensor product space is given by

$$\begin{aligned} \Pi : G &\rightarrow GL \left(\bigotimes_i V_i \otimes \bigotimes_j W_j^* \right) \\ \Pi(g) &= \rho_1(g) \otimes \cdots \otimes \rho_n(g) \otimes \sigma_m^*(g) \otimes \cdots \otimes \sigma_1^*(g) \end{aligned}$$

A tensor T is defined to be a **symmetric tensor** (or G -invariant tensor) if it is invariant under simultaneous action on all legs for all $g \in G$ [SPV10]:

$$\Pi(g)T = T \quad \forall g \in G$$

By interpreting T as a map

$$T : W_1 \otimes \cdots \otimes W_m \rightarrow V_1 \otimes \cdots \otimes V_n$$

it follows directly from the definition of global invariance that the tensor must commute with the group action and we get the following definition of a symmetric tensor:


Definition 1.3.4 (symmetric tensor). A *symmetric tensor* is an intertwiner of the representations of the group G :

$$\bigotimes_{j=1}^m \sigma_j(g)T = T \bigotimes_{i=1}^n \rho_i(g) \quad \forall g \in G,$$

which we denote by

$$T \in \text{Hom}_G(W, V).$$

TODO: what is the definition of an MPO representation? This could include this as well, maybe as $\text{Hom}_{\text{MPO}}(W, V)$.



Corollary 1.3.4.1 (Networks of symmetric tensors are symmetric). Since a tensor network is built by composition and application of the (co)ev maps, both of which are intertwiners, the full network is also an intertwiner. Important there is that this entire discussion is only about FdVect or FdHilb for now, but i think the same applies to any braided category.

1.3.5 Some Important Facts about Hamiltonians

Definition 1.3.5 (Gapped Hamiltonian). Let H be a hamiltonian acting on a finite system of size L . Let

$$E_0 \leq E_1 \leq \dots$$

be the eigenvalues. The spectral gap is then

$$\Delta(L) = E_1 - E_0.$$

The Hamiltonian is gapped if there is a constant Δ

1.3.6 Review of Symmetric Tensor Networks.

Generally, we follow the notations of [SV12], [Sch+20], [Sch+20], [Sch20] which gave an overview for implementing $SU(2)$ invariant tensor networks. A general overview for (Non)abelian symmetries is [SPV10], and a detailed account for the abelian $U(1)$ in [SPV11]. There are a multitude of implementations given in major packages, such as ITensor, [FWS22], Cyten, TenPy, [Hau+24], QSpace, [Wei24] ChemTensor, UniTensor, YasTn (Abelian Only) [Ram+25] (MPSKit?), (Cytnx (Abelian only i think.))). A general account of symmetries is given in [Unf24] with its respective implementation in [Hau+24] (We have to build cytns citation manually.)

The motivation for constructing symmetric tensor networks is as follows: Assume we have a (unitary) representation of a group G denoted $U(g)$. Furthermore assume we have a hamiltonian that commutes with $[U(g)^{\otimes L}, H] = 0$. If the ground state is unique, we can obtain

$$U(g)^{\otimes L} H |\psi_0\rangle = H(U(g)^{\otimes L}) |\psi_0\rangle = E_0 U(g)^{\otimes L} |\psi_0\rangle$$

and thus $U(g) |\psi_0\rangle$ is a ground state as well. By uniqueness

$$U(g) |\psi_0\rangle = e^{i\theta(g)} |\psi_0\rangle$$




Figure 1.1: Three different presentations of the XXZ Hamiltonian. (i): In the default computational basis of spin vectors. (ii) In a reordering of the computational basis into equal spin up (down) classes, i.e. no spin up, one spin up etc... (iii) After changing the basis into the S^2 basis.

Definition 1.3.6 (XXZ-Hamiltonian). The XXZ Hamiltonian on N sites is

$$H = J \cdot \sum_{i=1}^N (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_{i+1}^z). \quad (1.1)$$

As given in 1.3.6, this acts on the *spin basis* made of states $|s_1 s_2 \dots s_n\rangle$ where $s_i \in \{\uparrow, \downarrow\}$.

We can plot H and realize the sparsity structure thereof. Furthermore, a suitable rearrangement induces a block structure 1.1.

Now, we can realize about this hamiltonian is that there is a symmetry with the representations of $U(1)$. Lets review this shortly. A representation of $U(1)$ is a group homomorphism

$$\rho : U(1) \rightarrow GL(V), \quad \rho(e^{i\theta_1})\rho(e^{i\theta_2}) = \rho(e^{i(\theta_1+\theta_2)}).$$

The XXZ model lives in the Hilbert Space $\mathcal{H} = (\mathbb{C}^2)^n$ and the relevant representation is generated by the total magnetization operator

$$S_{\text{tot}}^z = \frac{1}{2} \sum_{i=1}^N \sigma_i^z$$

The group action is given by

$$U(\theta) = e^{-i\theta S_{\text{tot}}^z}.$$

We now claim that H is invariant under $U(\theta)$.

$$U(\theta) H U(\theta)^\dagger = H, \quad \forall \theta \in \mathbb{R}.$$

Equivalently,

$$[H, S_{\text{tot}}^z] = 0.$$

Recall that when two operators commute, we can *simultaneously diagonalize* them [HJ12, Theorem 1.3.12].

$$[H, S_{\text{tot}}^z] = \sum_i^N [\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_i^z, \sum_{j=1}^N \sigma_j^z]$$

and since operators on different sites commute, only $j = i$ and $j = i + 1$ contribute. So we ought to write

$$\sum_i^N [\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_i^z, \sigma_i^z + \sigma_{i+1}^z]$$

and now using the identity $[AB, C] = A[B, C] + [A, C]B$:

$$[\sigma_i^x \sigma_{i+1}^x, \sigma_i^z + \sigma_{i+1}^z] = -2i(\sigma_i^y \sigma_{i+1}^x + \sigma_i^x \sigma_{i+1}^y)$$

$$[\sigma_i^y \sigma_{i+1}^y, \sigma_i^z + \sigma_{i+1}^z] = 2i(\sigma_i^x \sigma_{i+1}^y + \sigma_i^y \sigma_{i+1}^x)$$

which cancel out. Furthermore, the z spins commute as well. Thus the whole sum is 0 and we are done.

MPS

Definition 1.3.7 (Matrix Product State). A *Matrix Product State* is a quantum state $|\psi\rangle$ with n legs written as the contraction of n trivalent tensors $\{A^{[i]}\}_{i \in [n]}$. Formally this reads

$$|\psi[A]\rangle = \sum_{i_1, \dots, i_n} \text{Tr}(A_{i_1}^{[1]} A_{i_2}^{[2]} \cdots A_{i_n}^{[n]}) |i_1 i_2 \cdots i_n\rangle,$$

where $A_{i_k}^{[i]} \in \mathbb{C}^{D_i \times D_{i+1}}$. The elements of the collection $\{D_i\}_{i \in [n]}$ is referred to as the *bond-dimensions*.

Note that this is a state with closed boundaries. If we want open boundaries we have to replace the trace with appropriate end vectors (nice source?).

Example 1. We can represent each product state $|\Psi\rangle = |\psi\rangle^{\otimes n}$ as a MPS with bond dimension 1.


Theorem 1.3.8 (Fundamental Theorem of MPS). If two MPS states generated by A and B respectively. Figure out what we can do with this.

Proof [Cir+17], [VFW18], [Per+08].

Chapter 2

Decompositions for Categorical Networks

This chapter generalizes the decomposition in [SV12, Equation 27] to both Fusion Categories and the Representation categories of compact groups. Note that in [DH25], such a generalization is given for representations of finite groups, which does not include $SU(2)$. However I do not yet know if we should do so, since I not yet sure how to argue the F moves here. If we instead take the deformation $U_q(sl_2)$ we get finiteness (and fusion cats?), unfortunately $U_q(sl_2)$ is not a group.



2.1 Decomposition of semisimple Categories

Lemma 2.1.1. For a vector space W , there is an isomorphism

$$\bigoplus^m W \simeq \mathbb{C}^m \otimes W.$$

Proof. Let $e_{i \in [m]}$ be the standard basis on $\mathbb{C}^m$. Let

$$\Phi : \mathbb{C}^m \otimes W \rightarrow \bigoplus^m W$$

be defined by

$$\phi(e_i \otimes w) = (0, \dots, w, \dots, 0)$$

and linear, thus

$$\Phi \left(\sum_i c_i e_i \otimes w_i \right) = (c_1 w_1, \dots, c_m w_m).$$

is clearly bijective, and we have our desired isomorphism. $\square$

Lemma 2.1.2. Let D be a finite dimensional k -vector space and let V be an object of a k -linear category C . Then

$$D \otimes V \cong V^{\oplus \dim D}$$

Proof. Let $n = \dim D$. Take a basis $\{e_i\}$ for D . This yields an isomorphism

$$D \cong \bigoplus_{i=1}^n k \cdot e_i \cong k^n$$

Since in a k -linear category the tensor product is additive (should we prove this?), we can distribute

$$D \otimes W \cong \bigoplus_1^n (k \cdot e_i \otimes W) \cong \bigoplus W = W^{\oplus n}$$

$\square$

Lemma 2.1.3 (Isotopic Decomposition). Let C be a semisimple k -linear category with simple objects $\{R_a\}$. Then every object $V \in C$ admits a canonical decomposition

$$V \cong \bigoplus_a D_a \otimes R_a. \tag{2.1}$$

where we refer to $D_a := \text{Hom}_C(R_a, V)$ as the *degeneracy* of the space.

Proof. Since C is semisimple, there exist integers $m_a \geq 0$ such that

$$V \cong \bigoplus_{a \in L} R_a^{\oplus m_a}.$$

Applying the additive functor $\text{Hom}_C(R_b, -)$ yields

$$\text{Hom}_C(R_b, V) \cong \bigoplus_{a \in L} \text{Hom}_C(R_b, R_a^{\oplus m_a}) \cong \bigoplus_{a \in L} \text{Hom}_C(R_b, R_a)^{\oplus m_a}.$$

Then, by Schurs lemma

$$\bigoplus_{a \in L} \text{Hom}_C(R_b, R_a)^{\oplus m_a} = k^{\oplus m_b}$$

□

So we actually need this: Maybe this is simpler if we always let the degeneracy be C ?

Lemma 2.1.4. Let C be a k -linear category and let D, E be finite-dimensional k -vector spaces. For any objects $R, S \in C$, there is a natural isomorphism

$$\text{Hom}_C(D \otimes R, E \otimes S) \cong \text{Hom}_k(D, E) \otimes \text{Hom}_C(R, S).$$

Proof. Let $m = \dim_k D$ and $n = \dim_k E$. Proof by isomorphism chains:

$$\begin{aligned} \text{Hom}_C(D \otimes R, E \otimes S) &\cong \text{Hom}_C(R^{\oplus m}, S^{\oplus n}) && \text{by 2.1.1} \\ &\cong \bigoplus_{i=1}^m \bigoplus_{j=1}^n \text{Hom}_C(R, S) && \text{by additivity of } \text{Hom}_C \\ &\cong k^{mn} \otimes \text{Hom}_C(R, S) && \text{by 2.1.1} \\ &\cong \text{Hom}_k(D, E) \otimes \text{Hom}_C(R, S) && \text{since } \text{Hom}_k(D, E) \cong k^{mn} \end{aligned}$$

(The last as s)

□

Theorem 2.1.5 (Decomposition of Semisimple Categories). Let C be a semisimple, monoidal, k -linear category whose tensor product is k -linear in each variable, with simple objects $\{R_a\}_{a \in L}$ and let V_i, W_i be a collection of objects in C . Every morphism

$$T \in \text{Hom}_C \left(\bigotimes_i V_i, \bigotimes_i W_i \right)$$

admit a decomposition as

$$T \in \bigoplus_{\hat{a}, \hat{b}} P^{(\hat{a}, \hat{b})} \otimes \text{Hom}_C(R_{\hat{a}}, R_{\hat{b}}),$$

where $\hat{a} = (a_1, \dots, a_n)$ and P is a k -vector space of dimension ???

Proof.

$$\begin{aligned}
& \text{Hom}_C \left(\bigotimes_i V_i, \bigotimes_j W_j \right) \\
& \cong \text{Hom}_C \left(\bigotimes_i \left(\bigoplus_{a_i} D_{i,a_i} \otimes R_{a_i} \right), \bigotimes_j \left(\bigoplus_{b_j} E_{j,b_j} \otimes R_{b_j} \right) \right) \quad \text{by 2.1.3} \\
& \cong \text{Hom}_C \left(\bigoplus_{\hat{a}} D_{\hat{a}} \otimes R_{\hat{a}}, \bigoplus_{\hat{b}} E_{\hat{b}} \otimes R_{\hat{b}} \right) \quad \text{since } \otimes \text{ is } k\text{-linear in each variable} \\
& \cong \bigoplus_{\hat{a}, \hat{b}} \text{Hom}_C (D_{\hat{a}} \otimes R_{\hat{a}}, E_{\hat{b}} \otimes R_{\hat{b}}) \quad \text{by additivity of } \text{Hom}_C \\
& \cong \bigoplus_{\hat{a}, \hat{b}} \text{Hom}(D_{\hat{a}}, E_{\hat{b}}) \otimes \text{Hom}_C(R_{\hat{a}}, R_{\hat{b}}) \quad \text{by 2.1.4.}
\end{aligned}$$

□

Corollary 2.1.5.1 (Operators are block matrices). An operator

$$\text{Hom}(V, W) \cong \text{Hom}_C \left(\bigoplus_a D_a \otimes R_a, \bigoplus_b E_b \otimes R_b \right) \cong \bigoplus_{a,b} \text{Hom}(D_a, E_b) \otimes \text{Hom}_C(R_a, R_b)$$

and by Schurs lemma this is just

$$\bigoplus_{a \in \text{Irr}(V) \cap \text{Irr}(W)} \text{Hom}(D_a, E_a)$$

where $\text{Hom}(D_a, E_a)$ is a matrix. Lets say that a is in a collection of isomorphisms L . Then (show matrix here ...)

Corollary 2.1.5.2. Let $Q_{\mu, \eta}$ be a basis of $\text{Hom}_C(\bigotimes_{\hat{a}} R_{\hat{a}}, \bigotimes_{\hat{b}} R_{\hat{b}})$ where μ denotes the multiplicities and η the internal fusions. In a *multiplicity-free* Category we will omit μ as a label.

Example 2. If we let our monoidal semisimple category just be Vec_C , the only simple object is C . Thus there is also only the trivial biproduct over one sector.

Example 3. If we restrict to the category $\text{Rep}(\text{SU}(2))$ we can build the symmetric tensor networks as given in [SV12]. TODO: actual ly do this and convince yourself that the notations amd the triplet basis workd as desired.

Example 4. Take a 2 in 1 out tensor in $\text{Rep}(\text{SU}(2))$ as

$$T \in \text{Hom}_{\text{Rep}(\text{SU}(2))}(V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}, V_{\frac{1}{2}})$$

Since all parts are simple, nothing happens.

Example 5. However, now assign $V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}$ to each leg in the example above. Each leg then becomes $V_0 \oplus V_1 \cong \mathbb{C} \otimes V_0 \oplus \mathbb{C} \otimes V_1$. Then we can start with the decomposition.

$$\begin{aligned} & \text{Hom}_{\mathbb{C}} \left(\bigoplus_{a,b,c \in (0,1)} (\mathbb{C} \otimes \mathbb{C} \otimes \mathbb{C}) \otimes V_a \otimes V_b \otimes V_c, \bigoplus_{b \in (0,1)} \mathbb{C} \otimes V_b \right) \\ &= \bigoplus_{a,b,c,d \in (0,1)} \text{Hom}_{\mathbb{C}}((\mathbb{C} \otimes \mathbb{C} \otimes \mathbb{C}) \otimes (V_a \otimes V_b \otimes V_c), \mathbb{C} \otimes V_d) \\ &= \bigoplus_{a,b,c,d} \text{Hom}(\mathbb{C}^3, \mathbb{C}) \otimes \text{Hom}_{\text{Rep}(\text{SU}(2))}(V_a \otimes V_b \otimes V_c, V_d) \end{aligned}$$

Example 6. Easier. However, now assign $V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}$ to each leg in the example above. Each leg then becomes $V_0 \oplus V_1 \cong \mathbb{C} \otimes V_0 \oplus \mathbb{C} \otimes V_1$. Then we can start with the decomposition.

$$\begin{aligned} & \text{Hom}_{\mathbb{C}} \left(\bigoplus_{a,b \in (0,1)} (\mathbb{C} \otimes \mathbb{C}) \otimes V_a \otimes V_b, \bigoplus_{b \in (0,1)} \mathbb{C} \otimes V_b \right) \\ &= \bigoplus_{a,b,c \in (0,1)} \text{Hom}_{\mathbb{C}}((\mathbb{C} \otimes \mathbb{C}) \otimes (V_a \otimes V_b), \mathbb{C} \otimes V_c) \\ &= \bigoplus_{a,b,c} \text{Hom}(\mathbb{C}^2, \mathbb{C}) \otimes \text{Hom}_{\text{Rep}(\text{SU}(2))}(V_a \otimes V_b, V_c) \end{aligned}$$

Example 7. Consider the same operator as in example 5 but with its incoming legs fused together. So we have $V_{\text{in}} = V_{\frac{1}{2}}^4$ and $V_{\text{out}} = V_{\frac{1}{2}}^2$. This decomposes $V_2 \oplus V_1^{\oplus 3} \oplus V_0^{\oplus 2}$. Then we can directly use [2.1.5.1](#) to obtain

$$\text{Hom}(V_{\text{in}}, V_{\text{out}}) =$$

Example 8. Lets look at a tensor with in legs $(\tau \otimes \tau)$ and τ and out legs τ .

$$\begin{aligned} & \text{Hom}_{\mathbb{C}}(((k \otimes 1) \oplus (k \otimes \tau)) \otimes (k \otimes \tau), k \otimes \tau) = \text{Hom}_{\mathbb{C}}() \\ & \text{Hom}_{\mathbb{C}}(((k \otimes 1) \oplus (k \otimes \tau)) \otimes (k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}_{\mathbb{C}}((k \otimes 1 \otimes k \otimes \tau) \oplus (k \otimes \tau \otimes k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}_{\mathbb{C}}(k \otimes 1 \otimes k \otimes \tau, k \otimes \tau) \oplus \text{Hom}_{\mathbb{C}}(k \otimes \tau \otimes (k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}(k \otimes k, k) \otimes \text{Hom}_{\mathbb{C}}(1 \otimes \tau, \tau) \oplus \text{Hom}(k \otimes k, k) \otimes \text{Hom}_{\mathbb{C}}(\tau \otimes \tau, \tau). \end{aligned}$$

Example 9 (Anyonic matrix operator). Finally, consider a $4 \rightarrow 4$ anyon operator with one leg on each side, representing the space $\text{Hom}((\tau \otimes \tau \otimes \tau \otimes \tau), (\tau \otimes \tau \otimes \tau \otimes \tau))$. The full decomposition is $2 + 3\tau$. Hence we obtain

$$\begin{aligned} & \text{Hom}_{\mathbb{C}}((k^2 \otimes 1) \oplus (k^3 \otimes \tau), (k^2 \otimes 1) \oplus (k^3 \otimes \tau)) \\ &= \bigoplus_{a,b \in (1,\tau)} \text{Hom}_{\mathbb{C}}(k^{m_a} \otimes a, k^{m_b} \otimes b) = \bigoplus_{a,b \in (1,\tau)} \text{Hom}(k^{m_a}, k^{m_b}) \otimes \text{Hom}_{\mathbb{C}}(a, b) \end{aligned}$$

where

$$m_1 = 2, \quad m_\tau = 3.$$

For this case, Schurs lemma applies. So we have

$$= \text{Hom}(k^2, k^2) \oplus \text{Hom}(k^3, k^3)$$

which is just a block matrix with indexing $(1_1, 1_2, \tau_1, \tau_2, \tau_3)$.

Example 10 ($U(1)$ Symmetry). Technically we would want to show here that this is simpler in general. TOOD. Make a theorem? Or a corollary?

Example 11 (Ground state on 4 anyons). We can be given a ground state in the operator form as $\text{Hom}(\tau^4, \tau)$.

2.1.1 Q intertwiners

Lets look at 2 paths for the 4 index tensor, $(a, b, c \rightarrow d)$ and $(a, b \rightarrow c, d)$.

$$(Q_{j_a j_b j_c}^{j_e})_{m_1, m_2, m_3, m_4} = \sum_{m_e} \langle j_1 m_1; j_2 m_2 | j_e m_e \rangle \langle j_e m_e; j_3 m_3 | j_4 m_4 \rangle$$

while the other topology would look like

$$Q \dots = \sum_{m_e} \langle j_1 m_1; j_2 m_2 | j_e m_e \rangle \langle j_e m_e | j_3 m_3; j_4 m_4 \rangle$$

and as an operator this would have to live in $\text{Hom}(V_{j_1} \otimes V_{j_2}, V_{j_3} \otimes V_{j_4})$

The yoga tree $(-1, 1, -3), (-2, 1, -4)$ Gives us a good mixture of those. This Q would then have to look like

$$Q_{-1-2-3-4}^1 = \sum_1 \langle j_{-1} m_{-1}; j_1 m_1 | j_{-3} m_{-3} \rangle \langle j_{-2} m_{-2}; j_1 m_1; j_{-4} m_{-4} \rangle = \sum_{m_1} C_{-1,1 \rightarrow -3} \cdot C_{-2,1 \rightarrow -4}$$

How would we do this when we have a contraction of two simultaneous indices:

$$Q_{-1-2}^{1,2} = \sum_{1,2} \langle -1 | 1; 2 \rangle \langle 1; 2 | -2 \rangle$$

If we define τ to be the fusion trees topology, then we can define ${}_\tau Q_j^i$

What would this look like for general symmetry groups G ? Lets choose a basis of its invariant decomp $|\lambda_a \mu_a\rangle$. Then the generalized CG coefficient would look like

$$(C_{\lambda_a \lambda_b \rightarrow \lambda_c}^\alpha)_{\mu_a \mu_b}^{\mu_c} = \langle \lambda_a \mu_a; \lambda_b \mu_b | \lambda_c \mu_c \rangle_\alpha$$

where alpha denotes the singular fusion multiplicity at μ_c .

Now to fully specify Q here for k sectors (λ_i, μ_i) , we need to give it $k - 3$ internal edges $(\lambda_{e_1}, \dots, \lambda_{e_{k-3}})$ along with $k - 2$ mulitplicity symbols, $(\alpha_1, \dots, \alpha_{k-2})$

Can we write down a general formula for Q?

This would then be

$$Q_\lambda$$

2.2 Categories

2.2.1 Tensor and Fusion Categories

Let C be a category with bifunctor

$$\otimes : C \times C \rightarrow C, \quad (X, Y) \mapsto X \otimes Y.$$

and the isomorphism $F : (- \otimes -) \otimes - \simeq - \otimes (- \otimes -)$

... introduce fusion categories here, with associativity and commutativity restrictions, I think this would be a good start.

Definition 2.2.1 (Monoidal Category). A monoidal category is a category C with a bifunctor $\otimes : C \times C \rightarrow C$

Let C and D be linear abelian categories. Then the *Deligne Tensor Product* $C \boxtimes D$ is a linear abelian category with a bifunctor

$$\boxtimes : C \times D \rightarrow C \boxtimes D$$

with morphisms given by

$$\text{Hom}_{C \boxtimes D}(X \boxtimes Y, W \boxtimes V) = \text{Hom}_C(X, W) \otimes \text{Hom}_D(Y, V)$$

Definition 2.2.2 (The Deligne Tensor Product of Categories). Let k be a field, and let $\mathcal{C}$ and $\mathcal{D}$ be two k -linear, finite abelian categories. The **Deligne tensor product**, denoted by $\mathcal{C} \boxtimes \mathcal{D}$, is a k -linear finite abelian category equipped with a k -bilinear functor

$$\boxtimes : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{C} \boxtimes \mathcal{D}, \quad (X, Y) \mapsto X \boxtimes Y$$

TODOTOD!!

Definition 2.2.3 (The Category Vec_k). Let k be an algebraically closed field.

- **Objects:** The objects are finite dimensional vector spaces over k .
- **Morphisms:** Linear maps between vector spaces.

Definition 2.2.4 (The Category $\text{Rep}_k(G)$). Let G be a group and k a field. The *category of linear representations of G over k* consists of the following data:

- **Objects:** The objects $\text{Rep}_k(G)$ are pairs (V, ρ_V) , where V is a k -vector space and $\rho_V : G \rightarrow \text{GL}(V)$.
- **Morphisms:** For any two objects (V, ρ_V) and (W, ρ_W) in $\text{Rep}_k(G)$, a morphism from V to W is a k -linear transformation $f : V \rightarrow W$ that is G -invariant (also called an *intertwiner*). That is, for every element $g \in G$, the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\rho_V(g)} & V \\ f \downarrow & & \downarrow f \\ W & \xrightarrow{\rho_W(g)} & W \end{array}$$

In algebraic terms, f must satisfy the condition:

$$f \circ \rho_V(g) = \rho_W(g) \circ f \quad \forall g \in G$$

The set of all such morphisms is denoted by $\text{Hom}_G(V, W)$.

We can further endow $\text{Rep}_k(G)$ with a monoidal structure: For any two objects $(V, \rho_V), (W, \rho_W)$, their tensor product is in $\text{Rep}_k(G)$ by the diagonal action

$$\rho_{(V \otimes W)}(g)(v \otimes w) = \rho_V(g)v \otimes \rho_W(g)w$$

Definition 2.2.5 (Direct product of representations). A representation of $G_1 \times G_2$ is a space where the pair (g_1, g_2) acts linearly. SO this is a pair

$$(V, \rho), \rho : (G_1 \times G_2) \rightarrow \text{GL}(V)$$

Now, let $(\text{Rep}_{\mathbb{C}}(G), \otimes, \mathbb{C})$ be the monoidal category of representations. We equip the category with a tensor product that acts as follows:

TODO....

Theorem 2.2.6 (Fusion Representation Category of finite groups). Let G be a finite group. Then $\text{Rep}(G)$ is a fusion category.

Theorem 2.2.7. Let G_1 and G_2 be finite groups and let $\text{Rep}(G_1), \text{Rep}(G_2)$ be their respective symmetry categories. Then there is an equivalence of categories

$$\text{Rep}(G_1 \times G_2) \simeq \text{Rep}(G_1) \boxtimes \text{Rep}(G_2)$$

Big issue here: $\text{Rep}(\text{SU}(2))$ is not a fusion category as it is not finite. But if i read it correctly, the deligne product only requires abelianness of its groups. Now, $\text{Rep}(G)$ is monoidal for any G but is it abelian?

So this is actually true, it is abelian. We could obtain a restricted versions by for example always restricting $\text{SU}(2)$ to some $\text{SU}(2)_k$ for some finite k . But i do not yet know how to present this nicely.

A second advantage would be that we can always represent fusion trees as monoidal categories (is this true or is fusion required here again?), and we can write down a unified theory for the fusion trees. Then I hope to write a nice method to view the "eval" of such a fusion tree, i.e. its clebsch gordan tensors by contracting down the topology of the tree.

2.3 Categories Needed

The motivation for this section is to introduce the necessary category theory background, and provide all the necessary theorems for a format that includes both representations and fusion categories.

Definition 2.3.1 (Semisimple Category). An abelian category $\mathcal{C}$ is semisimple if every object is either simple or a direct sum of simple objects. An object is *simple* if its only subobjects are 0 or X itself. Any object can be written as

$$X = \bigoplus_{i \in I} S_i^{\oplus n_i}$$

Definition 2.3.2 (Abelian Category).

Lemma 2.3.3 (Schurs 1st lemma (which one?)). If S and T are simple objects in a semisimple category over an algebraically closed field (this is always $\mathbb{C}$ in our case), then it holds that

$$\text{Hom}(S, T) = \begin{cases} k, & \text{if } S \cong T; \\ 0, & \text{if } S \not\cong T. \end{cases}$$

Proof. Step 1. Let $f : S \rightarrow T$ be a morphism. Since T is simple

$$\text{im}(f) \cong 0 \quad \text{or} \quad \text{im}(f) \cong T$$

□

Lets give a coordinatized approach to fusion categories:

Definition 2.3.4 (Fusion Category). A *fusion category* is a finite set of labels (sectors?) L and vector spaces V_c^{ab} for all $a, b, c \in L$, as well as the isomorphisms

$$F_d^{abc} : \bigoplus_e V_e^{ab} \otimes V_d^{ec} \rightarrow \bigoplus_e V_d^{ae} \otimes V_e^{bc}$$

such that the pentagon holds.

Lemma 2.3.5. Let C be a k -linear monoidal category whose tensor product is k -linear in each variable. Then

$$\bigotimes_{r=1}^n \left(\bigoplus_{a_r} X_{r, a_r} \right) \cong \bigoplus_{\hat{a}} \bigotimes_{r=1}^n X_{r, a_r},$$

where

$$\hat{a} = (a_1, \dots, a_n).$$

Lemma 2.3.6 (Finite Groups admit Fusion Category). Let G be a finite group, the $\text{Rep}(G)$ is a fusion category.

Let $\text{Vec} := (\text{FDCVec}, \otimes, \mathbb{C})$ be the monoidal category of finite dimensional complex vector spaces where the last tuple entry $\mathbb{C}$ denotes the unit object, since there are isomorphisms $V \otimes \mathbb{C} \cong V \cong \mathbb{C} \otimes V$.

Let $(\text{CRep}(G), \otimes, ?)$ denote the representation Category for a compact group G . Here $\otimes$ is the tensor product of representations $(\pi \otimes \rho, V \otimes W)$.

Definition 2.3.7 (Direct sum). Given a category C with objects $\{a_n\}$, an object $\bigoplus_{i=1}^n a_i \in C$ is called the *direct sum*. This is equipped with the injections and projections

$$\iota_i : a_i \rightarrow \bigoplus a_i \quad \text{and} \quad \pi_i : \bigoplus a_i \rightarrow a_i$$

such that

$$\pi_i \circ \iota_i = \text{id}_{a_i} \forall i$$

Lemma 2.3.8 (Peter Weyl Theorem). Let G be a compact group. Then all its unitary representations are completely reducible.

2.3.1 Representation Theory

IG remove this since we will only be dealing with finite groups? discuss!

Definition 2.3.9 (Compact Group). A compact group is a group G equipped with a topology such that

$$m : G \times G \rightarrow G, \quad (g, h) \mapsto gh,$$

is continuous.

$$i : G \rightarrow G, \quad g \mapsto g^{-1}$$

is continuous and G is compact as a topological space.

Example 12. 1. Every finite group with the discrete topology is compact.

2.

$$U(n) = \{A \in M_n(\mathbb{C}) : A^*A = I\}$$

is compact for all integer n .

3. All

$$SU(n) = \{A \in U(n) : \det A = 1\}$$

are compact.

Definition 2.3.10 (Representation). Let G be a group and V a vector space over a field $\mathbb{F}$ (In our cases $\mathbb{C}$) suffices. A *representation* of G on V is a group homomorphism

$$\rho : G \longrightarrow GL(V),$$

where $GL(V)$ denotes the group of invertible linear transformations on V . Equivalently

$$\rho(gh) = \rho(g)\rho(h), \quad \rho(e) = I, \quad \rho(g^{-1}) = \rho(g)^{-1}.$$

The vector space V is called the *representation space*.

Definition 2.3.11 (Projective Representation). Given a group G , a linear representation of G over $\mathbb{C}$ of dimension n is a map $U : g \rightarrow \mathbb{C}^{n \times n}$.

2.3.2 Fusion Trees

We consider a fusion tree as the basis that spans $\text{Hom}(R_a \otimes R_b \dots \otimes R_m, \otimes R_{a_j} \dots)$. Since $R_a \otimes R_b$ is semisimple, we have the relationship

$$R_a \otimes R_b \cong \bigoplus_c R_c^{\oplus N_{ab}^c}$$

with embedding (make this formal)

$$g_{c,\mu}^{ab} : R_c \rightarrow R_a \otimes R_b$$

and projections

$$f_{ab}^{c,\mu} : R_a \otimes R_b \rightarrow R_c$$

$$\text{id}_{R_a \otimes R_b} = \sum_{c \in L} \sum_{\mu=1}^{N_{ab}^c} g_{c,\mu}^{ab} f_{ab}^{c,\mu}$$

Lemma 2.3.12.

$$\text{Hom}(R_{a_1} \otimes \cdots \otimes R_{a_N}, R_b)$$

is spanned by ordinary fusion trees.

Proof. Base case: $N = 1$. Nothing to prove here.

Inductive case:

Under semisimplicity we get that

$$\text{Hom}(R_{a_1} \otimes \cdots \otimes R_{a_n}, R_b) \cong \bigoplus_{c,\mu} \text{Hom}(R_c \otimes R_{a_3} \otimes \cdots \otimes R_{a_n}, R_b)$$

□

2.3.3 Monoidal data

Definition 2.3.13 ($\text{Rep}(\text{SU}(2))$). Let (V, ρ_V) be a representation.

Definition 2.3.14 (Twists from R moves). Suppose we are in a braided monoidal category with F and R symbols. Then we compute the *twist* (a self braiding) as follows:

$$R_c^{ab} R_c^{ba} = \frac{\theta_c}{\theta_a \theta_b}.$$

(state more possibilities here.)

Example 13. Take $s\text{Vect}$. Our simples are 1 and f . This *braiding* is in fact symmetric. Thus we have that

$$1 = \frac{\theta_1}{\theta_f^2} =$$

2.4 Anyonic Hamiltonians

Symmetric States

A state $|\psi\rangle$ is symmetric if $U^n(g)\psi = \psi$. Similarly, its density need require that $U^n(g)\rho U^n(g) = \rho$.

2.4.1 1D Lattice Models

So from what i can read here (also the nlab or wikipedia) a spin chain specifically only refers to a per site construction with a tensor product hilbert space. An anyon chain is, well, an anyonic chain. These do not admit such a decomposition. So we should call these simply 1D lattice models.

In [Fei+07], anyonic chains, specifically in the Fib model, are introduced simply as the fusion space of n anyons. Further, the authors define a primitive hamiltonian that assigns -1 to the interaction $\tau \otimes \tau = 1$ and 0 to the interaction $\tau \otimes \tau = \tau$. This is in turn realized on the basis of the intermediate states, and evaluated using the method of exact diagonalization [empty citation]. This is extended in [Tre+08], with longer ranger interaction. It will become apparent that the construction given in these papers is a manualization of the tensor network constructions we will build here. Generalizations to arbitrary models are given in [Gil+13] as well as [BG17]. Generalized spin chains and dualities using module categories are given in [Eck25].

The tensor network formalism is handled in [Pfe+10], [Pfe12] and [Aye17], and in terms of categories this is done in [Unf24] and [DH25].

For the golden chain, we consider $n + 2$ Fib anyons as our ground state. Their combined state yields a Fusion Tree $\text{Hom}(\tau^{\otimes n+1}, \tau)$ with intermediate charges $x_1, \dots, x_n$. These intermediate charges are further restricted as in that there are no possibilities to assign adjacent 1s to them (In fact this is what causes the dimension of the hilbert space to reduce to ϕ^n).

Generally, we can define an analogue of a general state of this N site lattice.

There is a three local hamiltonian for the golden chain defined as

$$H_i = - \sum_{x'_i} [F_{x_{i+1}}^{x_{i-1}\tau\tau}]_{x_i}^{x'_i} E_{x'_i} [F_{x_{i+1}}^{x_{i-1}\tau\tau}]_{x'_i}^{x_i}$$

Write this three local hamiltonian in an operator form

$$H_i = -J \begin{pmatrix} 1 & & & \\ & 0 & & \\ & & 0 & \\ & & \varphi^{-2} & \varphi^{-\frac{3}{2}} \\ & & \varphi^{-\frac{3}{2}} & \varphi^{-1} \end{pmatrix}$$

We can build this as an anyonic operator in $\text{Hom}(\tau \otimes \tau, \tau \otimes \tau)$ as

$$H_i = E_1 \begin{array}{c} \tau \quad \tau \\ \diagdown \quad \diagup \\ \bullet \\ | \\ \bullet \\ \diagup \quad \diagdown \\ \tau \quad \tau \end{array} 1 + E_\tau \begin{array}{c} \tau \quad \tau \\ \diagdown \quad \diagup \\ \bullet \\ | \\ \bullet \\ \diagup \quad \diagdown \\ \tau \quad \tau \end{array} \tau$$

In this case we can view the golden chain approach chosen above as a manualization of the tensor network process.

We can also write H_i in its operator form matrix as

$$H_i \in \bigoplus_{a \in 1, \tau} \text{Hom}(\mathbb{C}, \mathbb{C})$$

as

$$H_i = -J \begin{array}{c} 1 \quad \tau \\ 1 \quad \left(\begin{array}{cc} E_1 & 0 \\ 0 & E_\tau \end{array} \right) \\ \tau \end{array}$$

$$H_i = -J \begin{array}{c} |x_0, \mathbf{1}, \tau\rangle \\ |x_0, \tau, \mathbf{1}\rangle \\ |x_0, \tau, \tau\rangle_1 \\ |x_0, \tau, \tau\rangle_2 \\ |x_0, \tau, \tau\rangle_3 \end{array} \begin{array}{ccccc} |1\tau 1\rangle & |\tau\tau 1\rangle & |1\tau\tau\rangle_1 & |\tau 1\tau\rangle_2 & |\tau\tau, \tau\rangle_3 \\ \left(\begin{array}{ccccc} 1 & & & & \\ & 0 & & & \\ & & 0 & & \\ & & & \varphi^{-2} & \varphi^{-\frac{3}{2}} \\ & & & \varphi^{-\frac{3}{2}} & \varphi^{-1} \end{array} \right) \end{array}$$

2.4.2 Rydberg-blockade ladder

This is given by the fusion category $\text{so}(3)_2$. Here we have fusion rules

$$1 \otimes 1 = 0 \oplus 1 \oplus 2, \quad 2 \otimes 1 = 1, \quad 2 \otimes 2 = 0.$$

2.5 I dont know where to put this yet

If we start out on anyonic chains, like the $\text{SU}(2)_3$ case, this provides a nice generalization for chains with $C = \text{Rep}(\text{SU}(2)_k)$ with fusion rules

$$j_1 \times j_2 = \sum_{j=|j_1-j_2|}^{\min(j_1+j_2, k-j_1-j_2)} j.$$

However now, if we approach this in the case of $\text{SU}(2)_\infty$ these fusion rules read

$$j_1 \times j_2 = \sum_{j=|j_1-j_2|}^{j_1+j_2}$$

Is this the same as the Q deformations? It is for $q = 1$ and $k = \infty$ but what about the intermediates?

2.6 MPO-symmetries

So im not sure how this works completely but there seems to be way more literature on this than on (globally) symmetric tensors. However this also seems to be connected to anyonic networks. Also to the Category of MPS and MPO?

There is an account of (projective) representation on a peps that act on all levels in [Bri14]. Actually there are more such as chapter IV section 2 of [CWV26], [Wil+16]

Some sources here: On Lattice Models with Module Categories: [Eck25], [Loo+23] [LDV24b] [LDV25]. Also chapters IV and V of [CWV26]. On the structure see [Liu+25]

I think the original motivation for this is to have the following constraint on MPOs:

$$\text{MPO}_a \circ \text{MPO}_b = \sum_c N_{ab}^c \text{MPO}_c$$

which we can obtain by a fusion tensor X_{ab}^c that obeys an intertwining property.

We immediately obtain our case of global symmetry by encoding

$$\rho(g) = \rho_1(g) \otimes \cdots \otimes \rho_n(g)$$

as a MPO with bond dimension 1 by setting

$$W_{11,ij}^{[a]} = \rho_a(g)_{ij}$$

Call $\rho(g) = \text{MPO}_g$. Then $\rho(g)\rho(h) = \rho(hg)$

If we want $\text{MPO}_g \text{MPO}_h = \text{MPO}_{hg}$ we also have MPO as a unitary representation with bond dimensions > 1 . So this generalizes the symmetry approach.

TODO: check if we can give symmetries such as in the wanniers kramer dual that are given by $Z_i Z_{i+1}$ by not a $\text{Rep}(G)$ but instead a $\text{Rep}(G)$ where G is a groupoid.

2.7 implementation

Implementations that are available:

We want to be able to represent a traditional tensor op such as

```
1 np.einsum("ijklmn,nkol -> ijmo", a, b)
```

for symmetric tensors. As we want to reuse code form existing librariries that is written using numpy, it would be ideal to write our classes in a way that would only require substituting np for our own libraries snp. Towards this goal we define a abstractclass Tensor

```
1 class Tensor(abc.ABC):
2     ...
3     @abstractmethod
4     def transpose(cls, new_shape:list[int]): ...
5     ...
```

and we implement both an np wrapper NPTensor and our symmetric Tensor as STensor, i.e.

Name	Symmetry	Citation
ITensor	Nonabelian (Fermion?)	[FWS22]
Cyten	Categorical	[UH]
TeNPy	Abelian	[Hau+24]
QSpace	Nonabelian	[Wei24] [Wei12]
ChemTensor	SU(2)	[Men26]
UniTensor	?	[<empty citation>]
YasTn	Abelian	[Ram+25]
TensorKit.jl	Categorical	[DH25]
Cytnx	Abelian only (i think)	[<empty citation>]
quimb	Fermion?	[<empty citation>]
MPSKit.jl	like TensorKit?	[<empty citation>]
TeMFpy	Fermion?	[HS25]
abeliantensor	U(1) and $\mathbb{Z}_n$	[Hau26]
symmray	Fermion	[Gao+25] [Gra26]
grassmanntn	Grassmann	[Yos26]

Table 2.1: Overview of tensor network libraries and symmetry support. (There are some closed source things here as well, also what is the software used by singh et al.?)

```

1 class STensor(Tensor):
2
3     def __init__(self, ...):
4         self.open_legs: list[int | str]
5         self.fusion_tree: FusionTree(self.open_legs, ...)
6         ...
7     @abstractmethod
8     def transpose(cls, new_shape: list[int]): ...
9     ...

```


Bibliography

- [Aye17] Babatunde Moses Ayeni. “Studies of Braided Non-Abelian Anyons Using Anyonic Tensor Networks”. PhD thesis. Macquarie University, Dec. 3, 2017. DOI: [10.25949/19435697.v1](https://doi.org/10.25949/19435697.v1). arXiv: [1708.06476 \[quant-ph\]](https://arxiv.org/abs/1708.06476). URL: https://figshare.mq.edu.au/articles/thesis/Studies_of_braided_non-Abelian_anyons_using_anyonic_tensor_networks/19435697/1 (visited on 06/28/2026).
- [BG17] Matthew Buican and Andrey Gromov. “Anyonic Chains, Topological Defects, and Conformal Field Theory”. In: *Communications in Mathematical Physics* 356.3 (Dec. 1, 2017), pp. 1017–1056. DOI: [10.1007/s00220-017-2995-6](https://doi.org/10.1007/s00220-017-2995-6). arXiv: [1701.02800v2 \[hep-th\]](https://arxiv.org/abs/1701.02800v2). (Visited on 06/28/2026).
- [Bia20] Jacob Biamonte. *Lectures on Quantum Tensor Networks*. Jan. 4, 2020. DOI: [10.48550/arXiv.1912.10049](https://doi.org/10.48550/arXiv.1912.10049). arXiv: [1912.10049 \[quant-ph\]](https://arxiv.org/abs/1912.10049). (Visited on 05/11/2026). Pre-published.
- [Bri14] Jacob Bridgeman. “Effective Edge States of Symmetry Protected Topological Systems”. MA thesis. Perimeter Institute and University of Waterloo, 2014. URL: <https://jcbbridgeman.github.io/assets/pdf/MastersThesis.pdf>.
- [Cir+17] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete. “Matrix Product Density Operators: Renormalization Fixed Points and Boundary Theories”. In: *Annals of Physics* 378 (Mar. 1, 2017), pp. 100–149. DOI: [10.1016/j.aop.2016.12.030](https://doi.org/10.1016/j.aop.2016.12.030). arXiv: [1606.00608v4 \[quant-ph\]](https://arxiv.org/abs/1606.00608v4).
- [CWV26] Bram Vancraeynest-De Cuiper, Weronika Wiesiolek, and Frank Verstraete. “Les Houches Lecture Notes on Tensor Networks”. In: *SciPost Physics Lecture Notes* (June 23, 2026), p. 128. DOI: [10.21468/SciPostPhysLectNotes.128](https://doi.org/10.21468/SciPostPhysLectNotes.128). arXiv: [2512.24390 \[cond-mat.str-el\]](https://arxiv.org/abs/2512.24390). (Visited on 06/27/2026).
- [dGro22] Caroline I de Groot. “Topological Phases, Symmetries and Open Systems”. Technische Universität München, 2022. URL: https://mediatum.ub.tum.de/1653795?query=Topological+Phases%2C+Symmetries+and+Open+Systems&show_id=1658844&srcnodeid=1653795 (visited on 05/09/2026).
- [DH25] Lukas Devos and Jutho Haegeman. *TensorKit.Jl: A Julia Package for Large-Scale Tensor Computations, with a Hint of Category Theory*. Version 1. 2025. DOI: [10.48550/ARXIV.2508.10076](https://doi.org/10.48550/ARXIV.2508.10076). (Visited on 06/01/2026). Pre-published.

- [Eck25] Luisa Eck. “Quantum Lattice Models from Fusion Categories”. PhD thesis. University of Oxford, 2025. URL: <https://ora.ox.ac.uk/objects/uuid:bcb39a31-1eaa-403d-bcf8-c7be6fb3d4d8> (visited on 06/28/2026).
- [Fei+07] Adrian Feiguin et al. “Interacting Anyons in Topological Quantum Liquids: The Golden Chain”. In: *Physical Review Letters* 98.16 (Apr. 20, 2007), p. 160409. doi: [10.1103/PhysRevLett.98.160409](https://doi.org/10.1103/PhysRevLett.98.160409). (Visited on 06/28/2026).
- [FWS22] Matthew Fishman, Steven R. White, and E. Miles Stoudenmire. “The ITensor Software Library for Tensor Network Calculations”. In: *SciPost Phys. Codebases* (2022), p. 4. doi: [10.21468/SciPostPhysCodeb.4](https://doi.org/10.21468/SciPostPhysCodeb.4).
- [Gao+25] Yang Gao et al. “Fermionic Tensor Network Contraction for Arbitrary Geometries”. In: *Physical Review Research* 7.2 (May 27, 2025), p. 023193. doi: [10.1103/PhysRevResearch.7.023193](https://doi.org/10.1103/PhysRevResearch.7.023193). (Visited on 05/04/2026).
- [Gil+13] C. Gils et al. “Anyonic Quantum Spin Chains: Spin-1 Generalizations and Topological Stability”. In: *Physical Review B* 87.23 (June 17, 2013), p. 235120. doi: [10.1103/PhysRevB.87.235120](https://doi.org/10.1103/PhysRevB.87.235120). (Visited on 06/28/2026).
- [Gra26] Johnnie Gray. *Jcmgray/Symmray*. June 21, 2026. URL: <https://github.com/jcmgray/symmray> (visited on 06/30/2026).
- [Hau+24] Johannes Hauschild et al. “Tensor Network Python (TeNPy) Version 1”. In: *SciPost Phys. Codebases* (2024), p. 41. doi: [10.21468/SciPostPhysCodeb.41](https://doi.org/10.21468/SciPostPhysCodeb.41).
- [Hau26] Markus Hauru. *Mhauru/Abeliantensors*. May 16, 2026. URL: <https://github.com/mhauru/abeliantensors> (visited on 06/30/2026).
- [HJ12] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. 2nd ed. Cambridge: Cambridge University Press, 2012. doi: [10.1017/CB09781139020411](https://doi.org/10.1017/CB09781139020411). URL: <https://www.cambridge.org/highereducation/books/matrix-analysis/FDA3627DC2B9F5C3DF2FD8C3CC136B48#overview> (visited on 06/30/2026).
- [HS25] Simon H. Hille and Attila Szabó. *TeMFPy: A Python Library for Converting Fermionic Mean-Field States into Tensor Networks*. 2025. arXiv: [2510.05227](https://arxiv.org/abs/2510.05227) [cond-mat.str-el]. URL: <https://arxiv.org/abs/2510.05227>.
- [LDV24a] Laurens Lootens, Clement Delcamp, and Frank Verstraete. “Dualities in One-Dimensional Quantum Lattice Models: Topological Sectors”. In: *PRX Quantum* 5.1 (Mar. 6, 2024), p. 010338. doi: [10.1103/PRXQuantum.5.010338](https://doi.org/10.1103/PRXQuantum.5.010338). arXiv: [2211.03777](https://arxiv.org/abs/2211.03777) [quant-ph]. (Visited on 06/08/2026).
- [LDV24b] Laurens Lootens, Clement Delcamp, and Frank Verstraete. “Dualities in One-Dimensional Quantum Lattice Models: Topological Sectors”. In: *PRX Quantum* 5.1 (Mar. 6, 2024), p. 010338. doi: [10.1103/PRXQuantum.5.010338](https://doi.org/10.1103/PRXQuantum.5.010338). (Visited on 06/25/2026).
- [LDV25] Laurens Lootens, Clement Delcamp, and Frank Verstraete. “Entanglement and the Density Matrix Renormalisation Group in the Generalised Landau Paradigm”. In: *Nature Physics* 21.10 (Oct. 2025), pp. 1657–1663. doi: [10.1038/s41567-025-02961-2](https://doi.org/10.1038/s41567-025-02961-2). arXiv: [2408.06334](https://arxiv.org/abs/2408.06334) [quant-ph]. (Visited on 06/24/2026).

- [Liu+25] Yuhan Liu et al. *Trading Mathematical for Physical Simplicity: Bialgebraic Structures in Matrix Product Operator Symmetries*. Sept. 3, 2025. doi: [10.48550/arXiv.2509.03600](https://doi.org/10.48550/arXiv.2509.03600). arXiv: [2509.03600](https://arxiv.org/abs/2509.03600) [[quant-ph](#)]. (Visited on 06/27/2026). Pre-published.
- [Loo+23] Laurens Lootens, Clement Delcamp, Gerardo Ortiz, and Frank Verstraete. “Dualities in One-Dimensional Quantum Lattice Models: Symmetric Hamiltonians and Matrix Product Operator Intertwiners”. In: *PRX Quantum* 4.2 (June 29, 2023), p. 020357. doi: [10.1103/PRXQuantum.4.020357](https://doi.org/10.1103/PRXQuantum.4.020357). (Visited on 05/16/2026).
- [Men26] Christian B. Mendl. *Qc-Tum/Chemtensor*. qc-tum, May 28, 2026. URL: <https://github.com/qc-tum/chemtensor> (visited on 06/30/2026).
- [Per+08] D. Perez-Garcia et al. “String Order and Symmetries in Quantum Spin Lattices”. In: *Physical Review Letters* 100.16 (Apr. 22, 2008), p. 167202. doi: [10.1103/PhysRevLett.100.167202](https://doi.org/10.1103/PhysRevLett.100.167202). arXiv: [0802.0447](https://arxiv.org/abs/0802.0447) [[cond-mat.str-el](#)]. (Visited on 06/28/2026).
- [Pfe+10] Robert N. C. Pfeifer et al. “Simulation of Anyons with Tensor Network Algorithms”. In: *Physical Review B* 82.11 (Sept. 29, 2010), p. 115126. doi: [10.1103/PhysRevB.82.115126](https://doi.org/10.1103/PhysRevB.82.115126). arXiv: [1006.3532](https://arxiv.org/abs/1006.3532) [[cond-mat.str-el](#)]. (Visited on 06/16/2026).
- [Pfe12] Robert N. C. Pfeifer. *Simulation of Anyons Using Symmetric Tensor Network Algorithms*. Feb. 10, 2012. doi: [10.48550/arXiv.1202.1522](https://doi.org/10.48550/arXiv.1202.1522). arXiv: [1202.1522](https://arxiv.org/abs/1202.1522) [[cond-mat](#)]. (Visited on 05/11/2026). Pre-published.
- [Ram+25] Marek M. Rams, Gabriela Wójtowicz, Aritra Sinha, and Juraj Hasik. “YASTN: Yet Another Symmetric Tensor Networks; A Python Library for Abelian Symmetric Tensor Network Calculations”. In: *SciPost Phys. Codebases* (2025), p. 52. doi: [10.21468/SciPostPhysCodeb.52](https://doi.org/10.21468/SciPostPhysCodeb.52).
- [Sch+20] Philipp Schmoll, Sukhbinder Singh, Matteo Rizzi, and Román Orús. “A Programming Guide for Tensor Networks with Global $SU(2)$ Symmetry”. In: *Annals of Physics* 419 (Aug. 1, 2020), p. 168232. doi: [10.1016/j.aop.2020.168232](https://doi.org/10.1016/j.aop.2020.168232). (Visited on 05/07/2026).
- [Sch20] Philipp Schmoll. “Tensor Network Simulations with Global $SU(2)$ Symmetry”. Mainz: Johannes Gutenberg University, July 2020. 150 pp. URL: <https://openscience.ub.uni-mainz.de/server/api/core/bitstreams/d9d9ca4e-46c8-4a8e-a831-7611475d2504/content> (visited on 05/08/2026).
- [Sel11] P. Selinger. “A Survey of Graphical Languages for Monoidal Categories”. In: *New Structures for Physics*. Ed. by Bob Coecke. Berlin, Heidelberg: Springer, 2011, pp. 289–355. doi: [10.1007/978-3-642-12821-9_4](https://doi.org/10.1007/978-3-642-12821-9_4). URL: <https://arxiv.org/abs/0908.3347> (visited on 06/24/2026).
- [Sim23] Steven H. Simon. *Topological Quantum*. Oxford University Press, Sept. 29, 2023. doi: [10.1093/oso/9780198886723.001.0001](https://doi.org/10.1093/oso/9780198886723.001.0001). URL: <https://doi.org/10.1093/oso/9780198886723.001.0001> (visited on 06/24/2026).
- [SPV10] Sukhwinder Singh, Robert N. C. Pfeifer, and Guifre Vidal. “Tensor Network Decompositions in the Presence of a Global Symmetry”. In: *Physical Review A* 82.5 (Nov. 8, 2010), p. 050301. doi: [10.1103/PhysRevA.82.050301](https://doi.org/10.1103/PhysRevA.82.050301). arXiv: [0907.2994](https://arxiv.org/abs/0907.2994) [[cond-mat.str-el](#)]. (Visited on 05/11/2026).

- [SPV11] Sukhwinder Singh, Robert N. C. Pfeifer, and Guifre Vidal. “Tensor Network States and Algorithms in the Presence of a Global $U(1)$ Symmetry”. In: *Physical Review B* 83.11 (Mar. 15, 2011), p. 115125. doi: [10.1103/PhysRevB.83.115125](https://doi.org/10.1103/PhysRevB.83.115125). arXiv: [1008.4774v1](https://arxiv.org/abs/1008.4774v1) [cond-mat.str-el]. (Visited on 05/06/2026).
- [SV12] Sukhwinder Singh and Guifre Vidal. “Tensor Network States and Algorithms in the Presence of a Global $SU(2)$ Symmetry”. In: *Physical Review B* 86.19 (Nov. 9, 2012), p. 195114. doi: [10.1103/PhysRevB.86.195114](https://doi.org/10.1103/PhysRevB.86.195114). arXiv: [1208.3919v1](https://arxiv.org/abs/1208.3919v1) [cond-mat.str-el]. (Visited on 05/11/2026).
- [Tre+08] Simon Trebst, Matthias Troyer, Zhenghan Wang, and Andreas W. W. Ludwig. “A Short Introduction to Fibonacci Anyon Models”. In: *Progress of Theoretical Physics Supplement* 176 (June 1, 2008), pp. 384–407. doi: [10.1143/PTPS.176.384](https://doi.org/10.1143/PTPS.176.384). arXiv: [0902.3275v1](https://arxiv.org/abs/0902.3275v1) [cond-mat.stat-mech]. (Visited on 06/28/2026).
- [UH] Jakob Unfried and Johannes Michael Hauschild. *Cyten*.
- [Unf24] Jakob Unfried. “Advancing Tensor Network Simulations -”. PhD thesis. Technische Universität München, 2024, p. 208. URL: <https://mediatum.ub.tum.de/1755273>.
- [Vak25] Ravi Vakil. *The Rising Sea: Foundations of Algebraic Geometry*. Princeton University Press, Oct. 21, 2025. 688 pp. URL: <https://press.princeton.edu/books/paperback/9780691268675/the-rising-sea> (visited on 06/24/2026).
- [VW18] Gert Vercleyen, Frank Verstraete, and Universiteit Gent. Faculteit Wetenschappen. “The Mathematical Structure of Tensor Networks”. MA thesis. Master of Science in de wiskunde, 2018. URL: https://lib.ugent.be/fulltxt/RUG01/002/479/664/RUG01-002479664_2018_0001_AC.pdf.
- [Wei12] Andreas Weichselbaum. “Non-Abelian Symmetries in Tensor Networks: A Quantum Symmetry Space Approach”. In: *Annals of Physics* 327.12 (Dec. 1, 2012), pp. 2972–3047. doi: [10.1016/j.aop.2012.07.009](https://doi.org/10.1016/j.aop.2012.07.009). (Visited on 05/07/2026).
- [Wei24] Andreas Weichselbaum. “QSpace - An Open-Source Tensor Library for Abelian and Non-Abelian Symmetries”. In: *SciPost Physics Codebases* (Nov. 14, 2024), p. 40. doi: [10.21468/SciPostPhysCodeb.40](https://doi.org/10.21468/SciPostPhysCodeb.40). arXiv: [2405.06632v1](https://arxiv.org/abs/2405.06632v1) [cond-mat.str-el]. (Visited on 05/08/2026).
- [Wen13] Xiao-Gang Wen. “Topological Order: From Long-Range Entangled Quantum Matter to an Unification of Light and Electrons”. In: *ISRN Condensed Matter Physics* 2013 (Mar. 27, 2013), pp. 1–20. doi: [10.1155/2013/198710](https://doi.org/10.1155/2013/198710). arXiv: [1210.1281](https://arxiv.org/abs/1210.1281) [cond-mat.str-el]. (Visited on 07/01/2026).
- [Wen95] Xiao-Gang Wen. *Topological Orders and Edge Excitations in FQH States*. Oct. 31, 1995. doi: [10.1080/00018739500101566](https://doi.org/10.1080/00018739500101566). arXiv: [cond-mat/9506066](https://arxiv.org/abs/cond-mat/9506066). (Visited on 06/30/2026). Pre-published.
- [Wil+16] Dominic J. Williamson et al. “Matrix Product Operators for Symmetry-Protected Topological Phases: Gauging and Edge Theories”. In: *Physical Review B* 94.20 (Nov. 30, 2016), p. 205150. doi: [10.1103/PhysRevB.94.205150](https://doi.org/10.1103/PhysRevB.94.205150). arXiv: [1412.5604](https://arxiv.org/abs/1412.5604) [quant-ph]. (Visited on 06/27/2026).

- [Yos26] Atis Yosprakob. *Ayosprakob/Grassmannntn*. June 7, 2026. URL: <https://github.com/ayosprakob/grassmannntn> (visited on 06/30/2026).