

Something about tensor networks, symmetries, anyons and categories.

Jeremias Rieser

advisors: C. Mendl and B. Decker

July 27, 2026

1 Introduction

Tensor Networks allow the efficient representation and computation of quantum states for large N using methods such as DMRG. Symmetric Tensor Networks yield a speedup over traditional methods for appropriate Hamiltonians. Furthermore, we would also like to represent *Anyonic* states and Hamiltonians by these operations, and since some anyon species are universal for quantum computation, we would like to represent circuits as well. It is folklore that tensor networks and quantum systems in general are governed by category-theoretic constructions.

In this thesis we give a formal overview how all these concepts fit into the same categorical framework while presenting a review of the existing literature, i.e. we give an interpretation of the results of Singh and Vidal [SV12], Weichselbaum [Wei12b], Pfeifer et al. [Pfe+10], Unfried [Unf24], and Devos and Haegeman [DH25], embed them with a rigorous categorical theory and provide missing proofs. Due to the interconnectedness with topological order, we phrase them in the language of [KZ22].

1.1 Thesis Outline

In Section 2 we review the technical literature which is shared by the chapters (sections?). We give accounts on symmetries, representation theory, fermionic networks, anyons and categories. Theorem 7.6 presents the main theorem that is the backend of our tensor system. While we will state the first 2 sections with scarce proofs, chapter ?? will be quite rigorous and supercede all statements from the previous sections.

Maybe we will present a outlook on Modular Categories

2 Background

Literature 2.1 (Tensor Networks). The modern theory of *tensor networks* shaped out of the study of *renormalization* and quantum many-body systems, initially with the *density-matrix-renormalization-group* of White [Whi92; Whi93] and later extended by Ostlund and Rommer [OR95] as a variational method (ansatz?) in the form of *matrix product states* (MPS) (Maybe this is actually due to [FNW92]). This was further generalized to 2 dimensions with the introduction of *projected entangled-pair states* (PEPS) [VC04]. *Multiscale-entanglement renormalization ansatz* (MERA) [Vid08] can be considered as an extension of MPS allowing for long-range entanglement (this is probably wrong). Whats missing? Holography [SMB18] but there are earlier ones.

In the purely mathematical view as a network of tensor contractions however, both concept and the notation we use today were introduced by [Pen71] under the name of *tensor-systems*.

While we introduce this concept classically and endow it with symmetry as a start, we will start concerning ourselves with a categorical embedding of tensor networks in what is commonly referred to as a *fusion-category* (Lit ???).

Literature 2.2 (Symmetric Tensor Network). Originally, the idea to exploit symmetries of the hamiltonian for a speedup was proposed to be used in dmrg by [MG02] (What is [SM98]? This sounds like it as well.). More

general theory for *globally invariant tensor networks*, i.e. networks that are built of tensors that are globally invariant is given by Singh, Pfeifer, and Vidal [SPV10] preceeding more detailed accounts on abelian symmetries [SPV11] and nonabelian (SU(2)) symmetries in [SV12]. A similar approach to (non)abelian symmetries is given by Weichselbaum [Wei12a; Wei20]. These rely on the general idea, that under representations of some compact group G , there are on site symmetries on each tensor, i.e. $[A, U_g^{\otimes k}] = 0$ there is a generalized *wigner-eckart-theorem* for tensors with n legs of the form

$$T \in \bigoplus_a D_a \otimes V_a$$

where D_a is a matrix and V_a is a *fusion-tree*, i.e. a clebsch-gordan tensor. Reviews and specific algorithms in the SU(2) case are given by [SO20; Sch20; Sch+20]. [Sch+20, § 8] proposes that this method is appropriate for Anyonic simulation as well (although this is done earlier by singh & pfeiffer.). Symmetries as categorified networks, as we will present in ??, are explained in [Unf24] and [DH25] and in this work in section 5.

For a list of software implementing symmetries see ??.

Literature 2.3 (Anyons). *Anyons* as quasiparticles with exotic exchange statistics were originally proposed by [Wil90] [LM77]. Anyons are usually described as *defects* in a (topologically ordered?) medium of dimension two (technically codimension after [SS23AnyonsInTedK]). The exchange statistics of N particles on this medium M is goverend by the *configuration space*

$$\text{Conf}_N(M) = \{(x_1, \dots, x_N) \in M^N : x_i \neq x_j\} \setminus S_N.$$

The statistics of the particle exchange is given by the representation of $\pi_1(\text{Conf}_N(M))$ (the fundamental (homology?) group). If $M = \mathbb{R}^2$, this corresponds to the *braid group* on N strands B_N . If the dimension of $M \geq 3$ this corresponds to the symmetric group S_N . If these representations are unitary and the (ground state?) is degenerate, this allows for *topological quantum computation* (c.f. Literature 2.7).

Literature 2.4 (Anyonic Spin Chains and Tensor Networks). (Check if this is the first). A first appearance of *Anyonic Tensor Networks* in terms of spin chains is given by Feiguin et al. [Fei+07] as the *golden chain*, which realizes a interacting 2-local hamiltonian on a n Fib-anyon lattice. This is reviewed in [Tre+08] with generalizations to 3-local and longer range hamiltonians. A general account outside of the Fib-model can be found in [Gil+13] and [BG17] (Check these again.). We can consider these approaches as a sort of manualization of the tensor network approach. Anyonic tensor networks are first introduced by [Pfe+10] and [Pfe12], developed out of the symmetry methods from section 3 and both the *isotopy* rules and anyonic operators defined in [BSS08]. Furthermore, stuctures such as an anyonic MERA are introduced.

We also need to discuss the topic of *anyonic entanglement entropy*, as given in [Pfe14; BKP17; Pfe12].

Literature 2.5 (Simulation of Anyons and String Nets). This should probably go in the previous section. [Liu+22] [Kir+23]

Literature 2.6 (Anyons as Categories). The correspondence between the *physical* treatment of anyons and the *categorical treatment* of anyons as a *braided fusion category* (modular functor?) was originally descibed by Kitaev [Kit06, § E, § 8] based on previous work in quantum field theory already making use of modular categories and their diagrammatic calculus such as [RT91]. Further treatments are found in [wang] and [simon-nayak-das-sarma]. A modern treatment of the realization through the Configuration space of n -anyons if given by [Val21] and in terms of modular functors by [Bur16].

Literature 2.7 (Topological Quantum Computation). Fault tolerant *Topological Quantum Computation* (TQC) using anabelian anyons was originally proposed by Kitaev [Kit03, § 7] (Originally published in 1997). Also [Fre98] and ??? has shown that there are nonabelian anyon models that are universal [DN05] [Moc03] [Moc04].

[MSS24] gives nicer literature. [Sim23] is a great textbook account from one of the original authors on quantum compiling and computing. [Nay+08] is also one of the original accounts. This does not need to be any longer. Maybe mention categorical computation in general? but idk what htis is yet [KZ23].

Literature 2.8 (Quantum Compiling). Some (Many?) anyon theories are universal for quantum computation. The *Solovay-Kitaev theorem* [Kit97][DN05] yields an efficient algorithm for approximating any quantum gate using only basic gates from a universal set. Here we can consider 1.) The representation of braid circuits as a tensor network operation and 2). A natural embedding of injection weaves as proposed by Simon et al. [Sim+06] where 2 anyons are braided in their composite fusion channel by realizing this operation as a tensor reshape.

Literature 2.9 (Categorified Quantum Mechanics). There are multiple approach to a diagrammatic and categorified approach to quantum mechanics. Generally one wants both spaces and maps represented by a framework that allows for parallel and entangled operations, i.e. has a tensor product structure. [Sel11] gives a review of this [VVW18] does for tensor networks specifically.

TODO: find the XZ-calculus literature by Kissinger? Specific examples such as XZ-calculus in the context of TQC is handled by [AK23].

TODO: How to mention that this results in or from Hilb_{fd} ? Does it? The work from selinger et.al does not yet include anyons. For this, one needs Literature 2.6.

Also, take [SS26, § 1] for a description of categorified mechanics. This stats that structure for *entangled* processes is given by the monoidal category of complex vector spaces.

$$\begin{array}{ccc}
 \begin{array}{l} \text{parallel quantum processes} \\ \text{entanglement, no-cloning} \end{array} & (\text{Mod}_{\mathbb{C}}, \otimes, \mathbb{C}) \xrightarrow{\dagger\text{-structure}} & (\text{Hilb}_{\text{fd}}, \otimes, ?) \\
 \uparrow & & \uparrow \\
 \begin{array}{l} \text{quantum processes} \\ \text{superpositions by } \oplus \end{array} & \text{Mod}_{\mathbb{C}} \xrightarrow{\dagger\text{-structure}} & \text{Hilb}_{\text{fd}}
 \end{array}$$

Additionally, many of the concepts of topological order are governed by category theory [KZ22], as well as the topic of higher symmetries (should this be its own Lit sect if relevant? It is for MPO symmetries [DT24] [Kon+20]).

2.1 Tensors (Everything here is $fdVect_k$)

Generally, we want to regard a tensor as a thing equipped with legs, where each legs holds a vector space. But we also want to be able to regard tensors as operators such as Hamiltonians.

Lets list some definitions of Tensor. A Tensor T can be regarded as an element of a tensor product space

$$T \in \bigotimes_{j \in J} V_j$$

where each V_j is a finite-dimensional vector space and we say T is a $|J|$ -legged tensor.

Given !finite! dimensional Vector spaces V and W , we have the identity that $V \otimes W^* \equiv \text{Hom}(W, V)$. This is called the *hom-tensor duality*. (TODO: is this canonical? What about the other way around?)

In a basis expansion, this would relate the expansion of $V \otimes W^*$ as

$$T = \sum_i v_i \otimes f_i$$

where $f_i \in W^* : W \rightarrow k$ with the map $T \in \text{Hom}(W, V)$ as

$$T = (w \mapsto \sum_i f_i(w) v_i).$$

Now, if we have a Tensor $T \in V_1 \otimes \cdots \otimes V_n$, and since for any Vector space V , $(V^*)^* = V$, we can peel off any number of V_i s, take their duals and represent T as a map

$$T : V_n^* \otimes \cdots \otimes V_{k+1}^* \rightarrow V_1 \otimes \cdots \otimes V_k$$

Note that we present the domain here in a contravariant fashion due to the contravariance of the dual.

Lets look at an example, let $t \in V_1 \otimes V_2 \otimes W_2^* \otimes W_1^*$. We can represent as an operator in dirac notation:

$$t = \sum_{i_1, i_2, j_1, j_2} T_{j_1 j_2}^{i_1 i_2} |v_{i_1}^1\rangle |v_{i_2}^2\rangle \langle w_{j_1}^1| \langle w_{j_2}^2|$$

where $|v_i\rangle$

Graphically, we represent a general tensor

$$T : (W_n^* \otimes \cdots \otimes W_1) \rightarrow (V_1 \otimes V_2 \otimes \cdots \otimes V_n)$$

as follows, where we choose incoming arrows for duals in the domain and outgoings for normal spaces, and on the bottom we choose incoming for duals and outgoing for normals.

3 Symmetric Tensor Networks

In the following, let G be a compact group and V, W be finite dimensional vector spaces.

Definition 3.1 (Representation). A *representation* of a compact group G over a vector space V is a group homomorphism

$$\rho : G \rightarrow \text{GL}(V), \quad \rho(g)\rho(h) = \rho(g \cdot h), \quad \rho(e) = id_V$$

The *representation space* is the tuple (ρ, V) but usually we shorten this to just V .

We can state this equivalently as the representation under the group action of G on V :

Definition 3.2 (Representation in action form). A representation is a map

$$\alpha : G \times V \rightarrow V, \quad (g, v) \mapsto g \cdot v$$

such that this is linear, todo. I think we get the equality to the hom def if we set $g \cdot v = \rho(g)v$

Definition 3.3 (Intertwiner). A linear map $A : V \rightarrow W$ is an *intertwiner*, if under representations $\rho_V : G \rightarrow \text{GL}(V)$ and $\rho_W : G \rightarrow \text{GL}(W)$ it holds that

$$A \circ \rho_V(g) = \rho_W(g) \circ A \quad \forall g \in G.$$

Denote the space of all intertwiners as $\text{Hom}_G(V, W)$.

Definition 3.4 (Unitary representation). If our vector space V is equipped with a (Hermitian?) inner product

Definition 3.5 (Irreducible representation (Woit 16)). A representation ρ is an *irrep* if there are no $W \subset V$ such that $(\rho|_W, W)$ is a representation.

Definition 3.6 (Tensor product representation). Given reps (ρ_V, V) and (ρ_W, W) , the *tensor product representation* $(\rho_{V \otimes W}, V \otimes W)$ is defined by its action on the elements

$$\rho_{V \otimes W}(g)(v \otimes w) = \rho_V(g)v \otimes \rho_W(g)w$$

Definition 3.7 (Direct sum representation). As above, the *direct sum representation* $(\rho_{V \oplus W}, V \oplus W)$

$$\rho_{V \oplus W}(g) = \begin{pmatrix} \rho_V(g) & \\ & \rho_W(g) \end{pmatrix}$$

Definition 3.8 (Isotypic / Semisimple decomposition). Each semisimple representation V decomposes as

$$V \cong \bigoplus_{i \in I} V^{\oplus n_i}$$

by grouping isomorphic simples.

Definition 3.9 (???). Let R_a and R_b be two irreducible representation (spaces). Their tensor product is again semisimple and there is a decomposition

$$R_a \otimes R_b \cong \bigoplus_{c \in I} R_c^{\oplus N_c^{ab}}$$

where we call N_c^{ab} the *fusion multiplicity* in G . If $N_c^{ab} \leq 1$ for all a, b, c , we say that the representation is *multiplicity-free*.

Example 3.10 (Clebsch-Gordan series). Let $G = \text{SU}(2)$ and let V_i and V_j be two spin irreps of charge i and j . These combine to

$$(\pi_{V_i \otimes V_j}, V_i \otimes V_j) = \bigoplus_{k=|i-j|}^{i+j} (\pi_{V_k}, V_k)$$

3.1 Symmetric Tensors

Let G be a compact group. Consider a tensor T living in the tensor product space of its legs:

$$T \in V_1 \otimes \cdots \otimes V_n \otimes W_m^* \otimes \cdots \otimes W_1^*$$

Let $\rho_i : G \rightarrow \text{GL}(V_i)$ be the linear representation of G on each vector space V_i , and let $\sigma_j^* : G \rightarrow \text{GL}(W_j^*)$ be the dual representation on W_j^* .

The total action of the group G on the entire tensor product space is given by

$$\begin{aligned} \Pi : G &\rightarrow \text{GL} \left(\bigotimes_i V_i \otimes \bigotimes_j W_j^* \right) \\ \Pi(g) &= \rho_1(g) \otimes \cdots \otimes \rho_n(g) \otimes \sigma_m^*(g) \otimes \cdots \otimes \sigma_1^*(g) \end{aligned}$$

A tensor T is defined to be a **symmetric tensor** (or G -invariant tensor) if it is invariant under simultaneous action on all legs for all $g \in G$ [SPV10]:

$$\Pi(g)T = T \quad \forall g \in G$$

Recall that by the isomorphism $\text{Hom}(V, W) \cong W \otimes V^*$ as

$$w \otimes f \mapsto (v \mapsto f(v)w).$$

We can push the product representation through this isomorphism as

$$\begin{aligned} \pi(g)(w \otimes f) &= \rho_W(g)w \otimes \rho_{V^*}(g)f = \rho_W(g)w \otimes (\rho_V(g)^{-1})^T f = \rho_W(g)w \otimes f(\rho_V(g)^{-1}) \\ &\cong v \mapsto f(\rho_V(g)^{-1}v) \rho_W(g)w \end{aligned}$$

By interpreting T as a map

$$T : W_1 \otimes \cdots \otimes W_m \rightarrow V_1 \otimes \cdots \otimes V_n$$


it follows directly from the definition of global invariance that the tensor must commute with the group action and we get the following definition of a symmetric tensor:

Definition 3.11 (symmetric tensor). A *symmetric tensor* is an intertwiner of the representations of the group G :

$$\bigotimes_{j=1}^m \sigma_j(g) \circ T = T \circ \bigotimes_{i=1}^n \rho_i(g) \quad \forall g \in G,$$

which we denote by

$$T \in \text{Hom}_G(W, V).$$



Corollary 3.12 (Networks of symmetric tensors are symmetric). *Since a tensor network is built by composition and application of the (co)ev maps, both of which are intertwiners, the full network is also an intertwiner. Important there is that this entire discussion is only about FdVect or FdHilb for now, but i think the same applies to any braided category.*

3.2 Review of Symmetric Tensor Networks.

The motivation for constructing symmetric tensor networks is as follows: Assume we have a (unitary) representation of a group G denoted $U(g)$. Furthermore assume we have a hamiltonian that commutes with $[U(g)^{\otimes L}, H] = 0$. If the ground state is unique, we can obtain

$$U(g)^{\otimes L} H |\psi_0\rangle = H(U(g)^{\otimes L}) |\psi_0\rangle = E_0 U(g)^{\otimes L} |\psi_0\rangle$$

and thus $U(g) |\psi_0\rangle$ is a ground state as well. By uniqueness

$$U(g) |\psi_0\rangle = e^{i\theta(g)} |\psi_0\rangle$$




Figure 1: Three different presentations of the XXZ Hamiltonian. (i): In the default computational basis of spin vectors. (ii) In a reordering of the computational basis into equal spin up (down) classes, i.e. no spin up, one spin up etc..., here U is simply a permutation matrix. (iii) After changing the basis into the (S^2, S_z) basis.

Definition 3.13 (XXZ-Hamiltonian). The XXZ Hamiltonian on N sites is

$$H = J \cdot \sum_{i=1}^N (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_{i+1}^z). \quad (1)$$

As given in definition 3.13, this acts on the *spin basis* made of states $|s_1 s_2 \dots s_n\rangle$ where $s_i \in \{\uparrow, \downarrow\}$.

We can plot H and realize the sparsity structure thereof. Furthermore, a suitable rearrangement induces a block structure fig. 1.

Now, we can realize about this hamiltonian is that there is a symmetry with the representations of $U(1)$. Lets review this shortly. A representation of $U(1)$ is a group homomorphism

$$\rho : U(1) \rightarrow \text{GL}(V)^n, \quad \rho(e^{i\theta_1}) \rho(e^{i\theta_2}) = \rho(e^{i(\theta_1 + \theta_2)}).$$

The XXZ model lives in the Hilbert Space $\mathcal{H} = (\mathbb{C}^2)^{\otimes n}$ and the relevant representation is generated by the total magnetization operator

$$S_{\text{tot}}^z = \frac{1}{2} \sum_{i=1}^N \sigma_i^z$$

The group action is given by

$$U(\theta) = e^{-i\theta S_{\text{tot}}^z}.$$

We now claim that H is invariant under $U(\theta)$.

$$U(\theta) H U(\theta)^\dagger = H, \quad \forall \theta \in \mathbb{R}.$$

Equivalently,

$$[H, S_{\text{tot}}^z] = 0.$$

Recall that when two operators commute, we can *simultaneously diagonalize* them [HJ12, Theorem 1.3.12].

$$[H, S_{\text{tot}}^z] = \sum_i^N [\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_i^z, \sum_{j=1}^N \sigma_j^z]$$

and since operators on different sites commute, only $j = i$ and $j = i + 1$ contribute. So we ought to write

$$\sum_i^N [\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_i^z, \sigma_i^z + \sigma_{i+1}^z]$$

and now using the identity $[AB, C] = A[B, C] + [A, C]B$:

$$[\sigma_i^x \sigma_{i+1}^x, \sigma_i^z + \sigma_{i+1}^z] = -2i(\sigma_i^y \sigma_{i+1}^x + \sigma_i^x \sigma_{i+1}^y)$$

$$[\sigma_i^y \sigma_{i+1}^y, \sigma_i^z + \sigma_{i+1}^z] = 2i(\sigma_i^x \sigma_{i+1}^y + \sigma_i^y \sigma_{i+1}^x)$$

which cancel out. Furthermore, the z spins commute as well. Thus the whole sum is 0 and we are done.

In the $SU(2)$ case, we have the generators

$$S_{\text{tot}}^\alpha = \frac{1}{2} \sum_{i=1}^N \sigma_i^\alpha, \quad \alpha \in \{x, y, z\}$$

The representation here is $U(R) = e^{i\theta n \cdot S_{\text{tot}}}$ and the Hamiltonian is invariant under

$$U(R) H U(R)^\dagger = H, \quad R \in SU(2)$$

Since

$$[S_{\text{tot}}^x, S_{\text{tot}}^y] = i S_{\text{tot}}^z$$

we need a new operator for the simultaneous diagonalization

$$S_{\text{tot}} = \sum_{a \in \{x, y, z\}} (S_{\text{tot}}^a)^2$$

and we need to verify this is indeed the correct operator TODO.

The eigenvalues of S_{tot}^2 are $S(S + 1)$ where S are the half integers.

Also we have that

$$(\mathbb{C}^2)^{\otimes 6} \cong V_{\frac{1}{2}}^{\otimes 6} \cong 5V_0 \oplus 9V_1 \oplus 5V_2 \oplus V_3$$

Now our operator is a morphism in $\text{End}(V_{\frac{1}{2}}^{\otimes 6})$ and thus by Schurs lemma¹

$$H \in \bigoplus_{S \in \{0, 1, 2, 3\}} \text{Mat}_{m_S \times m_S} \otimes I_{2S+1}(\mathbb{C}) \cong \bigoplus_{S \in \{0, 1, 2, 3\}} I_{2S+1} \otimes \text{Mat}_{m_S \times m_S}(\mathbb{C})$$

So we expect 20 blocks? 5 have dimension 1, 9 have dimension 3, 5 have dimension 5 and 1 has 7.

¹These are related by a permutation matrix. Proof? Is this the X from [DH25]?

Definition 3.14 (Matrix Product State). A *Matrix Product State* is a quantum state $|\psi\rangle$ with n legs written as the contraction of n trivalent tensors $\{A^{[i]}\}_{i \in [n]}$. Formally this reads

$$|\psi[A]\rangle = \sum_{i_1, \dots, i_n} \text{Tr} \left(A_{i_1}^{[1]} A_{i_2}^{[2]} \cdots A_{i_n}^{[n]} \right) |i_1 i_2 \cdots i_n\rangle,$$

where $A_{i_k}^{[i]} \in \mathbb{C}^{D_i \times D_{i+1}}$. The elements of the collection $\{D_i\}_{i \in [n]}$ is referred to as the *bond-dimensions*.

Note that this is a state with closed boundaries. If we want open boundaries we have to replace the trace with appropriate end vectors (nice source?).

Example 3.15. We can represent each product state $|\Psi\rangle = |\psi\rangle^{\otimes n}$ as a MPS with bond dimension 1.

Theorem 3.16 (Fundamental Theorem of MPS). *If two MPS states generated by A and B respectively. Figure out what we can do with this.*

Proof [Cir+17], [VWW18], [Per+08].

4 Fermionic Tensor Networks

Here we follow [Bul+17; Gao+25; Mor+25] primarily. These papers present the theory of fermionic networks over the category of *super vector spaces*, denoted sVect , which is also equal to a $\mathbb{Z}_2$ -graded Vector space $\text{Vec}_{\mathbb{Z}_2}$ as we will see in section ?? . Each $V \in \text{sVect}$ admits a decomposition into *odd* and *even* parity sectors

$$V = V_0 \oplus V_1.$$

Denote the tensor product in sVect by $\otimes_{\mathbb{Z}_2}$. Since fermions possess a non-trivial *exchange statistic*, i.e. exchanging two fermions incurs a phase factor of -1 on the wave function, the graded tensor product encodes this under its canonical braiding² also called the *Koszul sign rule*

$$|i\rangle \otimes_{\mathbb{Z}_2} |j\rangle = (-1)^{|i||j|} |j\rangle \otimes_{\mathbb{Z}_2} |i\rangle. \quad (2)$$



5 Anyonic Tensor Networks

We need to introduce anyons here, or somehow build it into the review of the golden chain.

Also we should do or have done isotopy before. Technically the discussion here also requires the completeness relations.


$$\begin{array}{c} a \quad b \\ \downarrow \quad \downarrow \\ \uparrow \quad \uparrow \end{array} = \sum_{c, \mu} \sqrt{\frac{d_c}{d_a d_b}} \begin{array}{c} a \quad b \\ \swarrow \quad \searrow \\ \uparrow c \\ \swarrow \quad \searrow \end{array} \quad (3)$$

²Check on the Nlab, the proof that this is canonical is quite complex?



5.1 1D Lattice Models & The Golden Chain

In [Fei+07], anyonic chains, specifically in the Fib model, are introduced as a 1D lattice of n τ anyons, defined as the fusion space thereof.



Here, the state is fully characterized by its intermediate indices $x_1 \dots x_{n-1}$, so we can choose a basis $|x_1 x_2 \dots x_{n-1}\rangle$.

Similar to the Heisenberg-chain³, the golden chain introduces a Hamiltonian that assigns -1 to the interaction $\tau \otimes \tau = 1$ and 0 to the interaction $\tau \otimes \tau = \tau$.

We can decompose the whole space $\text{Hom}(\tau^{\otimes n+1}, \tau)$ into the local states

$$\text{Hom}(\tau^{\otimes n+1}, \tau) = \bigoplus_{\substack{x_1, x_2, \dots, x_{n-1} \\ \text{all admissible paths}}} \text{Hom}(\tau \otimes \tau, x_1) \otimes \text{Hom}(x_1 \otimes \tau, x_2) \otimes \dots \otimes \text{Hom}(x_{n-1} \otimes \tau, \tau) \quad (4)$$

with the admissibility condition that there are not adjacent 1 in the sequence of x_i s.

This is evaluated as a 3 local hamiltonian, as we can decompose the spin chain into local parts where two τ anyons are adjacent (check how they call it in the paper.).

$$\begin{aligned} \begin{array}{c} \tau \quad \tau \\ \downarrow \quad \downarrow \\ x_{i-1} \quad x_i \quad x_{i+1} \end{array} &= \sum_{x_i} F_{x_{i+1}x_i}^{x_{i-1}\tau\tau} \begin{array}{c} \tau \quad \tau \\ \swarrow \quad \searrow \\ x_{i-1} \quad x_{i+1} \end{array} \\ &\in \text{Hom}(x_{i-1} \otimes \tau, x_i) \quad \in \text{Hom}(x_{i-1} \otimes x'_i, x_{i+1}) \\ &\otimes \text{Hom}(x_i \otimes \tau, x_{i+1}) \quad \otimes \text{Hom}(\tau \otimes \tau, x'_i) \end{aligned}$$

Now we can apply the interaction based on the intermediate state⁴ x'_i and apply another F -move back to the chain basis.

$$H_{[i,i+1]} : |x_{i-1}x_ix_{i+1}\rangle \rightarrow |x_{i-1}x'_ix_{i+1}\rangle := - \sum_{x'_i \in \{1, \tau\}} [F_{x_{i+1}}^{x_{i-1}\tau\tau}]_{x_i}^{x'_i} E_{x'_i} [F_{x_{i+1}}^{x_{i-1}\tau\tau}]_{x'_i}^{x_i''}$$

³Note that $\frac{1}{2} \otimes \frac{1}{2} = 0 \oplus 1$ in the Heisenberg $\text{SU}(2)$ model.

⁴Technically we need a projector here read up again.

Write this three local hamiltonian in an operator form

$$H_i = -J \begin{pmatrix} 1 & & & & \\ & 0 & & & \\ & & 0 & & \\ & & & \varphi^{-2} & \varphi^{-\frac{3}{2}} \\ & & & \varphi^{-\frac{3}{2}} & \varphi^{-1} \end{pmatrix}$$

This method is reviewed and extended in [Tre+08], with longer ranger interaction. It will become apparent that the construction given in these papers is a manualization of the tensor network constructions we will build here. Generalizations to arbitrary models are given in [Gil+13] as well as [BG17]. Generalized spin chains and dualities using module categories are given in [Eck25].

The tensor network formalism is handled in [Pfe+10], [Pfe12] and [Aye17], and in terms of categories this is done in [Unf24] and [DH25].

For the tensor network operations, we should consider the changing of the basis as an implicit operation that is performed when contracting, just like it would be in a normal einsum.

In a tensor network formalism can build this as an anyonic operator in $Hom(\tau \otimes \tau, \tau \otimes \tau)$ as

$$H_i = E_1 \begin{array}{c} \tau \quad \tau \\ \diagdown \quad \diagup \\ \bullet \\ | \\ \bullet \\ \diagup \quad \diagdown \\ \tau \quad \tau \end{array} + E_\tau \begin{array}{c} \tau \quad \tau \\ \diagdown \quad \diagup \\ \bullet \\ | \\ \bullet \\ \diagup \quad \diagdown \\ \tau \quad \tau \end{array}$$

(Note this is allowed due to $\mathbb{C}$ -linearity of Fib.)

In this case we can view the golden chain approach chosen above as a manualization of the tensor network process.

We can also write H_i in its operator form matrix as

$$H_i \in \bigoplus_{a \in 1, \tau} Hom(\mathbb{C}, \mathbb{C})$$

as

$$H_i = -J \begin{array}{c} |1\tau 1\rangle \\ |1\tau \tau\rangle \\ |\tau\tau 1\rangle \\ |\tau\tau \tau\rangle \end{array} \begin{pmatrix} 1 & & & & \\ & 0 & & & \\ & & 0 & & \\ & & & \varphi^{-2} & \varphi^{-\frac{3}{2}} \\ & & & \varphi^{-\frac{3}{2}} & \varphi^{-1} \end{pmatrix} \begin{array}{c} \langle 1\tau 1| \\ \langle 1\tau \tau| \\ \langle \tau\tau 1| \\ \langle \tau\tau \tau| \end{array}$$

$$H_i = -J \begin{array}{c} |1\tau 1\rangle \\ |\tau\tau 1\rangle \\ |1\tau \tau\rangle \\ |\tau\tau 1\rangle \\ |\tau\tau \tau\rangle \end{array} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \varphi^{-2} & \varphi^{-\frac{3}{2}} \\ 0 & 0 & 0 & \varphi^{-\frac{3}{2}} & \varphi^{-1} \end{pmatrix} \begin{array}{c} \langle 1\tau 1| \\ \langle \tau\tau 1| \\ \langle 1\tau \tau| \\ \langle \tau\tau 1| \\ \langle \tau\tau \tau| \end{array}$$

5.2 Anyonic States

For simple lattice states, [Sin+14] defines a bipartition of the lattice into anyons A and B . Then we consider the whole space under charge a_{tot} as

$$V_{a_{tot}} = \bigoplus_{N_{ab}^{a_{tot}}=1} (V_a^A \otimes V_b^B)$$

What i dont get here is why we are allowed to tensor both these sectors together without the space $V_{ab}^{a_{tot}}$ explicetly.

TODO delete all of this and introduce this over the Hom category.

Anyhow, the fundamental building blocks we are using are the vector spaces V_c^{ab} (where we refer to this as the fusion space from types a and b to c). This allows us to build the vector space of two anyonic types a, b as the sum over the fusion outcome:

$$V^{ab} = \bigoplus_{c \in Irr(C)} V_c^{ab}$$

We can directly introduce the R move (a braiding of the initial anyons) to obtain the isomorphic space V^{ba} as

$$R: \bigoplus_{c \in \text{Irr}(C)} V_c^{ab} \cong \bigoplus_{c \in \text{Irr}(C)} V_c^{ba}$$

We can also write the space $V_c^{ab} = \text{Hom}(a \otimes b, c)$ in the categorical language. Here we also need to realize that there is a difference between the isomorphism β and the action of R , which is really on the hom space $\text{Hom}(-, c)$. So this would be a yoneda perspective? Since we know R for all $a, b, c \in C$, we also know the action of R as an isomorphism. Well technically i believe that this allows us to determine for all simples c , and then semisimplicity extends this to all subjects (this would be nice to write down) and then the yoneda embedding is enough to know the whole space. Similarly, in our $\text{Rep}(G)$ categories (which i actually believe to be symmetric? Find this in etingof)

The algebra-category correspondence in the theory of anyons was first noted by [Kit06, App E]

$$\text{Hom} \left(\bigotimes_{k=1}^m X_{ik}, \bigotimes_{l=1}^n X_{jl} \right) = \text{span}_{\mathbb{C}} \left\{ \begin{array}{c} \begin{array}{ccccccc} & X_{i_1} & X_{i_2} & X_{i_3} & \dots & X_{i_{m-1}} & X_{i_m} \\ & \searrow & \swarrow & \downarrow & & \nearrow & \nwarrow \\ & e_1 & & & & & \\ & & e_2 & & & & \\ & & & \ddots & & & \\ & & & & e_{m-3} & & \\ & & & & & e_{m-2} & \\ & & & & & & b \\ & & f_{n-2} & & & & \\ & & & f_{n-3} & & & \\ & & & & \ddots & & \\ & & & & & & \\ & f_1 & & & & & \\ & \nearrow & \searrow & \downarrow & & & \\ X_{j_1} & X_{j_2} & X_{j_3} & \dots & X_{j_{n-1}} & X_{j_n} \end{array} \end{array} \right\} \quad (5)$$

$$\begin{array}{c}
\begin{array}{ccc}
& \bigoplus_{e,l} V_e^{ab} \otimes V_f^{cd} \otimes V_g^{el} & \\
id_{V_e^{ab}} \otimes F_g^{ecd} \nearrow & & \searrow F_g^{abl} \otimes id_{V_l^{cd}} \\
\bigoplus_{e,f} V_e^{ab} \otimes V_f^{ec} \otimes V_g^{fd} & & \bigoplus_{k,l} V_g^{ak} \otimes V_k^{bl} \otimes V_l^{cd} \\
F_f^{abc} \otimes id_{V_g^{fd}} \searrow & & \nearrow F_k^{bcd} \otimes id_{V_g^{ak}} \\
\bigoplus_{f,h} V_h^{af} \otimes V_f^{bc} \otimes V_g^{hd} & \xrightarrow{F_g^{afd} \otimes id_{V_f^{k,h}}} & \bigoplus_{f,h} V_g^{ak} \otimes V_h^{bc} \otimes V_k^{hd}
\end{array} \\
\\
\begin{array}{ccc}
& \bigoplus_{f,h} V_h^{af} \otimes V_f^{bc} \otimes V_g^{hd} & \\
\swarrow & & \searrow \\
\bigoplus_{e,f} V_e^{ab} \otimes V_f^{ec} \otimes V_g^{fd} & & \bigoplus_{f,h} V_h^{af} \otimes V_f^{bc} \otimes V_g^{hd} \\
F_f^{abc} \otimes id_{V_g^{fd}} \downarrow & & \uparrow \\
\bigoplus_{f,h} V_h^{af} \otimes V_f^{bc} \otimes V_g^{hd} & \xrightarrow{F_f^{abc} \otimes id_{V_g^{fd}}} & \bigoplus_{k,h} V_g^{ak} \otimes V_h^{bc} \otimes V_k^{hd}
\end{array} \\
\\
\begin{array}{ccc}
V \times W & \xrightarrow{\phi} & V \otimes W \\
& \searrow f & \downarrow \exists! \tilde{f} \\
& & U
\end{array}
\end{array} \tag{6}$$

$$\tag{7}$$

6 Categorical Networks

Definition 6.1 (Category). A *category* C consists of

- a set of *objects* $\text{ob}(C)$,
- for every pair $A, B \in \text{ob}(C)$, a *hom-set* $\text{Hom}_C(A, B)$ of *morphisms* $f : A \rightarrow B$,
- for every $A \in \text{ob}(C)$, an *identity morphism* $\text{id}_A : A \rightarrow A$,
- for every triple $A, B, C \in \text{ob}(C)$, a *composition map*

$$\circ : C(B, C) \times C(A, B) \longrightarrow C(A, C),$$

such that composition is *associative*, i.e. $(f \circ g) \circ h = f \circ (g \circ h)$ whenever defined, and *unital*, i.e. $f \circ \text{id}_A = f = \text{id}_B \circ f$ for all $f : A \rightarrow B$.

Note that some categories such as $\text{Rep}(G)$ and $\text{Vec}_{\mathbb{C}}$ admit more structures than ordinary categories. For fdVect , the hom spaces of linear maps $V \rightarrow W$ have a natural identification as a vector space as well. The composition map is $\mathbb{C}$ -bilinear. This immediately motivates the following definition:

Definition 6.2 ($\mathbb{C}$ -linear category). A $\mathbb{C}$ -linear category endows each of its hom-sets with a module structure, such that the composition is bilinear. That is

$$\circ : \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) \longrightarrow \text{Hom}_C(A, C)$$

is $\mathbb{C}$ -bilinear: for all $f, f' : B \rightarrow C$, $g, g' : A \rightarrow B$, and $c \in \mathbb{C}$,

$$(f + f') \circ g = f \circ g + f' \circ g, \quad f \circ (g + g') = f \circ g + f \circ g', \quad (cf) \circ g = c(f \circ g) = f \circ (cg).$$

We will now explore how to use category theory to model quantum processes. We want this to be able to do a list of things. First, given two systems A and B , we want the combined system to be an object as well. Given processes, i.e. operators on each individual system $f : A \rightarrow A', g : B \rightarrow B'$, there should be a combined process. Additionally, structures such as entanglement and no-cloning should be readily apparent in this theory.

Consider the tensor product $\otimes_{\mathbb{C}}$ of vector spaces V, W . We can define this using a *universal property* ...
TODO.

This should motivate the following definition:

Definition 6.3 (Monoidal Category). A *monoidal category* $\mathcal{C}$ is equipped with a unit $I \in \mathcal{C}$ and a tensor (functor) $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, as well as 3 structural isomorphisms

$$\rho_A : A \otimes I \cong A \quad \lambda_B : I \otimes B \cong B \quad \alpha_{ABC} : (A \otimes B) \otimes C \cong A \otimes (B \otimes C)$$


such that the following diagrams commute:

$$\begin{array}{ccc} (A \otimes I) \otimes B & \xrightarrow{\alpha_{A,I,B}} & A \otimes (I \otimes B) \\ \rho_A \otimes \text{id}_B \searrow & & \swarrow \text{id}_A \otimes \lambda_B \\ & A \otimes B & \end{array} \quad \begin{array}{ccc} ((A \otimes B) \otimes C) \otimes D & \xrightarrow{\alpha_{A,B,C} \otimes \text{id}_D} & (A \otimes (B \otimes C)) \otimes D \\ \alpha_{A \otimes B, C, D} \downarrow & & \downarrow \alpha_{A, B \otimes C, D} \\ (A \otimes B) \otimes (C \otimes D) & & A \otimes ((B \otimes C) \otimes D) \\ \alpha_{A, B, C \otimes D} \searrow & & \swarrow \text{id}_A \otimes \alpha_{B, C, D} \\ & A \otimes (B \otimes (C \otimes D)) & \end{array}$$

For simplicity we denote a monoidal category as the triple $(\mathcal{C}, \otimes, I)$.

6.1 Decompositions for Categorical Networks

This chapter generalizes the decomposition in [SV12, Equation 27] to both Fusion Categories and the Representation categories of compact groups. Note that in [DH25], such a generalization is given for representations of finite groups, which does not include $SU(2)$. However I do not yet know if we should do so, since I not yet sure how to argue the F moves here. If we instead take the deformation $U_q(sl_2)$ we get finiteness (and fusion cats?), unfortunately $U_q(sl_2)$ is not a group.



7 Decomposition of semisimple Categories

Lemma 7.1. For a vector space W , there is an isomorphism

$$\bigoplus^m W \simeq \mathbb{C}^m \otimes W.$$

Proof. Let $e_{ii \in [m]}$ be the standard basis on $\mathbb{C}^m$. Let

$$\Phi : \mathbb{C}^m \otimes W \rightarrow \bigoplus^m W$$

be defined by

$$\phi(e_i \otimes w) = (0, \dots, w, \dots, 0)$$

and linear, thus

$$\Phi \left(\sum_i c_i e_i \otimes w_i \right) = (c_1 w_1, \dots, c_m w_m).$$

is clearly bijective, and we have our desired isomorphism. $\square$

Lemma 7.2. Let D be a finite dimensional k -vector space and let V be an object of a k -linear category $\mathcal{C}$. Then

$$D \otimes V \cong V^{\oplus \dim D}$$

Proof. Let $n = \dim D$. Take a basis $\{e_i\}$ for D . This yields an isomorphism

$$D \cong \bigoplus_{i=1}^n k \cdot e_i \cong k^n$$

Since in a k -linear category the tensor product is additive (should we prove this?), we can distribute

$$D \otimes W \cong \bigoplus_1^n (k \cdot e_i \otimes W) \cong \bigoplus W = W^{\oplus n}$$

$\square$

Lemma 7.3 (Isotopic Decomposition). *Let C be a semisimple k -linear category with simple objects $\{R_a\}$. Then every object $V \in C$ admits a canonical decomposition*

$$V \cong \bigoplus_a D_a \otimes R_a. \quad (8)$$

where we refer to $D_a := \text{Hom}_C(R_a, V)$ as the degeneracy of the space.

Proof. Since C is semisimple, there exist integers $m_a \geq 0$ such that

$$V \cong \bigoplus_{a \in L} \bigoplus_{i=1}^{D_a^V} a \cong \bigoplus_{a \in L} R_a^{\oplus m_a}.$$

Applying the additive functor $\text{Hom}_C(R_b, -)$ yields

$$\text{Hom}_C(R_b, V) \cong \bigoplus_{a \in L} \text{Hom}_C(R_b, R_a^{\oplus m_a}) \cong \bigoplus_{a \in L} \text{Hom}_C(R_b, R_a)^{\oplus m_a}.$$

Then, by Schurs lemma

$$\bigoplus_{a \in L} \text{Hom}_C(R_b, R_a)^{\oplus m_a} = k^{\oplus m_b}$$

□

So we actually need this: Maybe this is simpler if we always let the degeneracy be C ?

Lemma 7.4. *Let C be a k -linear category and let D, E be finite-dimensional k -vector spaces. For any objects $R, S \in C$, there is a natural isomorphism*

$$\text{Hom}_C(D \otimes R, E \otimes S) \cong \text{Hom}_k(D, E) \otimes \text{Hom}_C(R, S).$$

Proof. Let $m = \dim_k D$ and $n = \dim_k E$. Proof by isomorphism chains:

$$\begin{aligned} \text{Hom}_C(D \otimes R, E \otimes S) &\cong \text{Hom}_C(R^{\oplus m}, S^{\oplus n}) && \text{by lemma 7.1} \\ &\cong \bigoplus_{i=1}^m \bigoplus_{j=1}^n \text{Hom}_C(R, S) && \text{by additivity of Hom}_C \\ &\cong k^{mn} \otimes \text{Hom}_C(R, S) && \text{by lemma 7.1} \\ &\cong \text{Hom}_k(D, E) \otimes \text{Hom}_C(R, S) && \text{since } \text{Hom}_k(D, E) \cong k^{mn} \end{aligned}$$

(The last as s)

□

Definition 7.5 (Copower). For a Vect_k -enriched (and tensored?) category C , and $E \in \text{Vect}_k, X \in C$ the power X^E is the object of C representing the natural isomorphism

$$\text{Hom}_C(Y, X^E) \cong \text{Hom}_k(E, \text{Hom}_C(Y, X))$$

Theorem 7.6 (Decomposition of Semisimple Categories). *Let C be a semisimple, monoidal, k -linear category whose tensor product is k -linear in each variable, with simple objects $\{R_a\}_{a \in L}$ and let V_i, W_i be a collection of objects in C . Every morphism*

$$T \in \text{Hom}_C \left(\bigotimes_{i=1}^n V_i, \bigotimes_{i=1}^m W_i \right)$$

admit a decomposition as

$$T \in \bigoplus_{\hat{a}, \hat{b}} P^{\hat{a}, \hat{b}} \otimes \text{Hom}_C(R_{\hat{a}}, R_{\hat{b}}),$$

where $\hat{a} = (a_1, \dots, a_n)$ and P is a k -vector space of dimension ???

Proof.

$$\begin{aligned}
& \text{Hom}_C \left(\bigotimes_i V_i, \bigotimes_j W_j \right) \\
& \cong \text{Hom}_C \left(\bigotimes_i \left(\bigoplus_{a_i} D_{i,a_i} \otimes R_{a_i} \right), \bigotimes_j \left(\bigoplus_{b_j} E_{j,b_j} \otimes R_{b_j} \right) \right) \quad \text{by lemma 7.3} \\
& \cong \text{Hom}_C \left(\bigoplus_{\hat{a}} D_{\hat{a}} \otimes R_{\hat{a}}, \bigoplus_{\hat{b}} E_{\hat{b}} \otimes R_{\hat{b}} \right) \quad \text{since } \otimes \text{ is } k\text{-linear in each variable} \\
& \cong \bigoplus_{\hat{a}, \hat{b}} \text{Hom}_C(D_{\hat{a}} \otimes R_{\hat{a}}, E_{\hat{b}} \otimes R_{\hat{b}}) \quad \text{by additivity of } \text{Hom}_C \\
& \cong \bigoplus_{\hat{a}, \hat{b}} \text{Hom}(D_{\hat{a}}, E_{\hat{b}}) \otimes \text{Hom}_C(R_{\hat{a}}, R_{\hat{b}}) \quad \text{by lemma 7.4.}
\end{aligned}$$

□

Corollary 7.7 (Operators are block matrices). *This is also given here: [KZ22, Remark 3.4.8]*

An operator

$$\text{Hom}(V, W) \cong \text{Hom}_C \left(\bigoplus_a R_a^{\oplus d_a}, \bigoplus_b R_b^{\oplus e_b} \right) \cong \bigoplus_{a,b} \text{Hom}(D_a, E_b) \otimes \text{Hom}_C(R_a, R_b)$$

and by Schurs lemma this is just

$$\bigoplus_{a \in \text{Irr}(V) \cap \text{Irr}(W)} \text{Hom}(D_a, E_a) \cong \bigoplus_{a \in \text{Irr}(V) \cap \text{Irr}(W)} \text{Mat}_{d_a \times e_a}(k)$$

Corollary 7.8. *Let $Q_{\mu,\eta}$ be a basis of $\text{Hom}_C(\bigotimes_{\hat{a}} R_{\hat{a}}, \bigotimes_{\hat{b}} R_{\hat{b}})$ where μ denotes the multiplicities and η the internal fusions. In a multiplicity-free Category we will omit μ as a label.*

Example 7.9. If we let our monoidal semisimple category just be $\text{Vec}_{\mathbb{C}}$, the only simple object is $\mathbb{C}$. Thus there is also only the trivial biproduct over one sector.

Example 7.10. If we restrict to the category $\text{Rep}(SU(2))$ we can build the symmetric tensor networks as given in [SV12]. TODO: actual ly do this and convince yourself that the notations amd the triplet basis workd as desired.

Example 7.11. Take a 2 in 1 out tensor in $\text{Rep}(SU(2))$ as

$$T \in \text{Hom}_{\text{Rep}(SU(2))}(V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}, V_{\frac{1}{2}})$$

Since all parts are simple, nothing happens.

Example 7.12. However, now assign $V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}$ to each leg in the example above. Each leg then becomes $V_0 \oplus V_1 \cong \mathbb{C} \otimes V_0 \oplus \mathbb{C} \otimes V_1$. Then we can start with the decomposition.

$$\begin{aligned}
& \text{Hom}_C \left(\bigoplus_{a,b,c \in (0,1)} (C \otimes C \otimes C) \otimes V_a \otimes V_b \otimes V_c, \bigoplus_{b \in (0,1)} C \otimes V_b \right) \\
& = \bigoplus_{a,b,c,d \in (0,1)} \text{Hom}_C((C \otimes C \otimes C) \otimes (V_a \otimes V_b \otimes V_c), C \otimes V_d) \\
& = \bigoplus_{a,b,c,d} \text{Hom}(C^3, C) \otimes \text{Hom}_{\text{Rep}(SU(2))}(V_a \otimes V_b \otimes V_c, V_d)
\end{aligned}$$

Example 7.13. Easier. However, now assign $V_{\frac{1}{2}} \otimes V_{\frac{1}{2}}$ to each leg in the example above. Each leg then becomes $V_0 \oplus V_1 \cong \mathbb{C} \otimes V_0 \oplus \mathbb{C} \otimes V_1$. Then we can start with the decomposition.

$$\begin{aligned} & \text{Hom}_C \left(\bigoplus_{a,b \in (0,1)} (C \otimes C) \otimes V_a \otimes V_b, \bigoplus_{b \in (0,1)} C \otimes V_b \right) \\ &= \bigoplus_{a,b,c \in (0,1)} \text{Hom}_C((C \otimes C) \otimes (V_a \otimes V_b), C \otimes V_c) \\ &= \bigoplus_{a,b,c} \text{Hom}(C^2, C) \otimes \text{Hom}_{\text{Rep}(SU(2))}(V_a \otimes V_b, V_c) \end{aligned}$$

Example 7.14. Consider the same operator as in example 5 but with its incoming legs fused together. So we have $V_{in} = V_{\frac{1}{2}}^4$ and $V_{out} = V_{\frac{1}{2}}^2$. This decomposes $V_2 \oplus V_1^{\oplus 3} \oplus V_0^{\oplus 2}$. Then we can directly use corollary 7.7 to obtain

$$\text{Hom}(V_{in}, V_{out}) =$$

Example 7.15. Lets look at a tensor with in legs $(\tau \otimes \tau)$ and τ and out legs τ .

$$\begin{aligned} & \text{Hom}_C(((k \otimes 1) \oplus (k \otimes \tau)) \otimes (k \otimes \tau), k \otimes \tau) = \text{Hom}_C() \\ & \text{Hom}_C(((k \otimes 1) \oplus (k \otimes \tau)) \otimes (k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}_C((k \otimes 1 \otimes k \otimes \tau) \oplus (k \otimes \tau \otimes k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}_C(k \otimes 1 \otimes k \otimes \tau, k \otimes \tau) \oplus \text{Hom}_C(k \otimes \tau \otimes (k \otimes \tau), k \otimes \tau) \\ & \cong \text{Hom}(k \otimes k, k) \otimes \text{Hom}_C(1 \otimes \tau, \tau) \oplus \text{Hom}(k \otimes k, k) \otimes \text{Hom}_C(\tau \otimes \tau, \tau). \end{aligned}$$

Example 7.16 (Anyonic matrix operator). Finally, consider a $4 \rightarrow 4$ anyon operator with one leg on each side, representing the space $\text{Hom}((\tau \otimes \tau \otimes \tau \otimes \tau), (\tau \otimes \tau \otimes \tau \otimes \tau))$. The full decomposition is $2 + 3\tau$. Hence we obtain

$$\begin{aligned} & \text{Hom}_C((k^2 \otimes 1) \oplus (k^3 \otimes \tau), (k^2 \otimes 1) \oplus (k^3 \otimes \tau)) \\ &= \bigoplus_{a,b \in (1,\tau)} \text{Hom}_C(k^{m_a} \otimes a, k^{m_b} \otimes b) = \bigoplus_{a,b \in (1,\tau)} \text{Hom}(k^{m_a}, k^{m_b}) \otimes \text{Hom}_C(a, b) \end{aligned}$$

where

$$m_1 = 2, \quad m_\tau = 3.$$

For this case, Schurs lemma applies. So we have

$$= \text{Hom}(k^2, k^2) \oplus \text{Hom}(k^3, k^3)$$

which is just a block matrix with indexing $(1_1, 1_2, \tau_1, \tau_2, \tau_3)$.

Example 7.17 ($U(1)$ Symmetry). Technically we would want to show here that this is simpler in general. TOOD. Make a theorem? Or a corollary?

Example 7.18 (Ground state on 4 anyons). We can be given a ground state in the operator form as $\text{Hom}(\tau^4, \tau)$.

7.1 Q intertwiners

Lets look at 2 paths for the 4 index tensor, $(a, b, c \rightarrow d)$ and $(a, b \rightarrow c, d)$.

$$(\mathcal{Q}_{j_a j_b j_c}^{j_e})_{m_1, m_2, m_3, m_4} = \sum_{m_e} \langle j_1 m_1; j_2 m_2 | j_e m_e \rangle \langle j_e m_e; j_3 m_3 | j_4 m_4 \rangle$$

while the other topology would look like

$$Q \dots = \sum_{m_e} \langle j_1 m_1; j_2 m_2 | j_e m_e \rangle \langle j_e m_e | j_3 m_3; j_4 m_4 \rangle$$

and as an operator this would have to live in $Hom(V_{j_1} \otimes V_{j_2}, V_{j_3} \otimes V_{j_4})$

The yoga tree $(-1, 1, -3), (-2, 1, -4)$ Gives us a good mixture of those. This Q would then have to look like

$$Q_{-1-2-3-4}^1 = \sum_1 \langle j_{-1}m_{-1}; j_1m_1 | j_{-3}m_{-3} \rangle \langle j_{-2}m_{-2} | j_1m_1; j_{-4}m_{-4} \rangle = \sum_{m_1} C_{-1,1 \rightarrow -3} \cdot C_{-2,1 \rightarrow -4}$$

How would we do this when we have a contraction of two simultaneous indices:

$$Q_{-1-2}^{1,2} = \sum_{1,2} \langle -1 | 1; 2 \rangle \langle 1; 2 | -2 \rangle$$

If we define τ to be the fusion trees topology, then we can define ${}_{\tau}Q_j^i$

What would this look like for general symmetry groups G ? Lets choose a basis of its invariant decomp $|\lambda_a t_a \mu_a\rangle$. Then the generalized CG coefficient would look like

$$(C_{\lambda_a \lambda_b \rightarrow \lambda_c}^{\alpha})_{\mu_a \mu_b}^{\mu_c} = \langle \lambda_a \mu_a; \lambda_b \mu_b | \lambda_c \mu_c \rangle_{\alpha}$$

where alpha denotes the singular fusion multiplicity at μ_c .

Now to fully specify Q here for k sectors (λ_i, μ_i) , we need to give it $k - 3$ internal edges $(\lambda_{e_1}, \dots, \lambda_{e_{k-3}})$ along with $k - 2$ mulitplicity symbols, $(\alpha_1, \dots, \alpha_{k-2})$

Can we write down a general formula for Q ?

This would then be

$$Q_{\lambda}$$

8 Categories

8.1 Tensor and Fusion Categories

Let C be a category with bifunctor

$$\otimes : C \times C \rightarrow C, \quad (X, Y) \rightarrow X \otimes Y.$$

and the isomorphism $F : (- \otimes -) \otimes - \simeq - \otimes (- \otimes -)$

... introduce fusion categories here, with associativity and commutativity restrictions, I think this would be a good start.

Definition 8.1 (Monoidal Category). A monoidal category is a category C with a bifunctor $\otimes : C \times C \rightarrow C$

Let C and D be linear abelian categories. Then the *Deligne Tensor Product* $C \boxtimes D$ is a linear abelian category with a bifunctor

$$\boxtimes : C \times D \rightarrow C \boxtimes D$$

with morphisms given by

$$Hom_{C \boxtimes D}(X \boxtimes Y, W \boxtimes V) = Hom_C(X, W) \otimes Hom_D(Y, V)$$

Definition 8.2 (The Deligne Tensor Product of Categories). Let k be a field, and let C and D be two k -linear, finite abelian categories. The **Deligne tensor product**, denoted by $C \boxtimes D$, is a k -linear finite abelian category equipped with a k -bilinear functor

$$\boxtimes : C \times D \longrightarrow C \boxtimes D, \quad (X, Y) \mapsto X \boxtimes Y$$

TODOTOD!!

Definition 8.3 (The Category Vec_k). Let k be an algebraically closed field.

- **Objects:** The objects are finite dimensional vectors spaces over k .

- **Morphisms:** Linear maps between vector spaces.

Definition 8.4 (The Category $\text{Rep}_k(G)$). Let G be a group and k a field. The *category of linear representations of G over k* consists of the following data:

- **Objects:** The objects $\text{Rep}_k(G)$ are pairs (V, ρ_V) , where V is a k -vector space and $\rho_V : G \rightarrow \text{GL}(V)$.
- **Morphisms:** For any two objects (V, ρ_V) and (W, ρ_W) in $\text{Rep}_k(G)$, a morphism from V to W is a k -linear transformation $f : V \rightarrow W$ that is G -invariant (also called an *intertwiner*). That is, for every element $g \in G$, the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\rho_V(g)} & V \\ f \downarrow & & \downarrow f \\ W & \xrightarrow{\rho_W(g)} & W \end{array}$$

In algebraic terms, f must satisfy the condition:

$$f \circ \rho_V(g) = \rho_W(g) \circ f \quad \forall g \in G$$

The set of all such morphisms is denoted by $\text{Hom}_G(V, W)$.

We can further endow $\text{Rep}_k(G)$ with a monoidal structure: For any two objects $(V, \rho_V), (W, \rho_W)$, their tensor product is in $\text{Rep}_k(G)$ by the diagonal action

$$\rho_{(V \otimes W)}(g)(v \otimes w) = \rho_V(g)v \otimes \rho_W(g)w$$

Definition 8.5 (Direct product of representations). A representation of $G_1 \times G_2$ is a space where the pair (g_1, g_2) acts linearly. SO this is a pair

$$(V, \rho), \rho : (G_1 \times G_2) \rightarrow \text{GL}(V)$$

Now, let $(\text{Rep}_{\mathbb{C}}(G), \otimes, \mathbb{C})$ be the monoidal category of representations. We equip the category with a tensor product that acts as follows:

TODO....

Theorem 8.6 (Fusion Representation Category of finite groups). *Let G be a finite group. Then $\text{Rep}(G)$ is a fusion category.*

Theorem 8.7. *Let G_1 and G_2 be finite groups and let $\text{Rep}(G_1), \text{Rep}(G_2)$ be their respective symmetry categories. Then there is an equivalence of categories*

$$\text{Rep}(G_1 \times G_2) \simeq \text{Rep}(G_1) \boxtimes \text{Rep}(G_2)$$

Big issue here: $\text{Rep}(\text{SU}(2))$ is not a fusion category as it is not finite. But if i read it correctly, the deligne product only requires abelianness of its groups. Now, $\text{Rep}(G)$ is monoidal for any G but is it abelian?

So this is actually true, it is abelian. We could obtain a restricted versions by for example always restricting $\text{SU}(2)$ to some $\text{SU}(2)_k$ for some finite k . But i do not yet know how to present this nicely.

A second advantage would be that we can always represent fusion trees as monoidal categories (is this true or is fusion required here again?), and we can write down a unified theory for the fusion trees. Then I hope to write a nice method to view the "eval" of such a fusion tree, i.e. its clebsch gordan tensors by contracting down the topology of the tree.

9 Categories Needed

The motivation for this section is to introduce the necessary category theory background, and provide all the necessary theorems for a format that includes both representations and fusion categories.

Definition 9.1 (Semisimple Category). An abelian category C is semisimple if every object is either simple or a direct sum of simple objects. An object is *simple* if its only subobjects are 0 or X itself. Any object can be written as

$$X = \bigoplus_{i \in I} S_i^{\oplus n_i}$$

Lemma 9.2 (Schur's lemma [Sch05]). If S and T are simple objects in a semisimple category over an algebraically closed field (this is always C in our case), then it holds that

$$\text{Hom}(S, T) = \begin{cases} k, & \text{if } S \cong T; \\ 0, & \text{if } S \not\cong T. \end{cases}$$

Proof [IR19, Theorem 7.5].

Lets give a coordinatized approach to fusion categories:

Definition 9.3 (Fusion Category). A *fusion category* is a finite set of labels (sectors?) L and vector spaces V_c^{ab} for all $a, b, c \in L$, as well as the isomorphisms

$$F_d^{abc} : \bigoplus_e V_e^{ab} \otimes V_d^{ec} \rightarrow \bigoplus_e V_d^{ae} \otimes V_e^{bc}$$

such that the pentagon holds.

Lemma 9.4. Let C be a k -linear monoidal category whose tensor product is k -linear in each variable. Then

$$\bigotimes_{r=1}^n \left(\bigoplus_{a_r} X_{r, a_r} \right) \cong \bigoplus_{\hat{a}} \bigotimes_{r=1}^n X_{r, a_r},$$

where

$$\hat{a} = (a_1, \dots, a_n).$$

Lemma 9.5 (Finite Groups admit Fusion Category). Let G be a finite group, the $\text{Rep}(G)$ is a fusion category.

Let $\text{Vec} := (\text{FDCVec}, \otimes, \mathbb{C})$ be the monoidal category of finite dimensional complex vector spaces where the last tuple entry $\mathbb{C}$ denotes the unit object, since there are isomorphisms $V \otimes \mathbb{C} \cong V \cong \mathbb{C} \otimes V$.

Let $(\mathbb{C}\text{Rep}(G), \otimes, ?)$ denote the representation Category for a compact group G . Here $\otimes$ is the tensor product of representations $(\pi \otimes \rho, V \otimes W)$.

Definition 9.6 (Direct sum). Given a category C with objects $\{a_n\}$, an object $\bigoplus_{i=1}^n a_i \in C$ is called the *direct sum*. This is equipped with the injections and projections

$$\iota_i : a_i \rightarrow \bigoplus a_i \quad \text{and} \quad \pi_i : \bigoplus a_i \rightarrow a_i$$

such that

$$\pi_i \circ \iota_i = \text{id}_{a_i} \forall i$$

Lemma 9.7 (Peter Weyl Theorem). Let G be a compact group. Then all its unitary representations are completely reducible.

9.1 Representation Theory

IG remove this since we will only be dealing with finite groups? discuss!

Definition 9.8 (Compact Group). A compact group is a group G equipped with a topology such that

$$m : G \times G \rightarrow G, \quad (g, h) \mapsto gh,$$

is continuous.

$$i : G \rightarrow G, \quad g \mapsto g^{-1}$$

is continuous and G is compact as a topological space.

Example 9.9. 1. Every finite group with the discrete topology is compact.

2.

$$U(n) = \{A \in M_n(\mathbb{C}) : A^* A = I\}$$

is compact for all integer n .

3. All

$$SU(n) = \{A \in U(n) : \det A = 1\}$$

are compact.

Definition 9.10 (Representation). Let G be a group and V a vector space over a field $\mathbb{F}$ (In our cases $\mathbb{C}$) suffices. A *representation* of G on V is a group homomorphism

$$\rho : G \longrightarrow \text{GL}(V),$$

where *operatornamename* $\text{GL}(V)$ denotes the group of invertible linear transformations on V . Equivalently

$$\rho(gh) = \rho(g)\rho(h), \quad \rho(e) = I, \quad \rho(g^{-1}) = \rho(g)^{-1}.$$

The vector space V is called the *representation space*.

Definition 9.11 (Projective Representation). Given a group G , a linear representation of G over $\mathbb{C}$ of dimension n is a map $U : g \rightarrow \mathbb{C}^{n \times n}$.

9.2 Fusion Trees

We consider a fusion tree as the basis that spans $\text{Hom}(R_a \otimes R_b \dots \otimes R_m, \otimes R_{a_j} \dots)$. Since $R_a \otimes R_b$ is semisimple, we have the relationship

$$R_a \otimes R_b \cong \bigoplus_c R_c^{\oplus N_{ab}^c}$$

with embedding (make this formal)

$$g_{c,\mu}^{ab} : R_c \rightarrow R_a \otimes R_b$$

and projections

$$f_{ab}^{c,\mu} : R_a \otimes R_b \rightarrow R_c$$

$$id_{R_a \otimes R_b} = \sum_{c \in L} \sum_{\mu=1}^{N_{ab}^c} g_{c,\mu}^{ab} f_{ab}^{c,\mu}$$

Lemma 9.12.

$$\text{Hom}(R_{a_1} \otimes \dots \otimes R_{a_N}, R_b)$$

is spanned by ordinary fusion trees.

Proof. Base case: $N = 1$. Nothing to prove here.

Inductive case:

Under semisimplicity we get that

$$\text{Hom}(R_{a_1} \otimes \dots \otimes R_{a_n}, R_b) \cong \bigoplus_{c,\mu} \text{Hom}(R_c \otimes R_{a_3} \otimes \dots \otimes R_{a_n}, R_b)$$

□

9.3 Monoidal data

Definition 9.13 ($\text{Rep}(SU(2))$). Let (V, ρ_V) be a representation.

Definition 9.14 (Twists from R moves [Sim23, Equation ?]). Suppose we are in a braided monoidal category with F and R symbols. Then we compute the *twist* (a self braiding) as follows:

$$R_c^{ab} R_c^{ba} = \frac{\theta_c}{\theta_a \theta_b}.$$

(state more possibilities here.)

Example 9.15. Take $sVect$. Our simples are 1 and f . This *braiding* is in fact symmetric. Thus we have that

$$1 = \frac{\theta_1}{\theta_f^2} =$$

I think we may be able to build a nice summary with this diagram:

$$\begin{array}{ccc}
 (A \otimes B) \otimes C & \xrightarrow{\alpha_{ABC}} & A \otimes (B \otimes C) \\
 \downarrow \text{Hom}(-, D) & & \downarrow \text{Hom}(-, D) \\
 \text{Hom}((A \otimes B) \otimes C, D) & \xrightarrow[\text{associator induces isomorphisms under yoneda}]{\alpha_{A,B,C}} & \text{Hom}(A \otimes (B \otimes C), D) \\
 \downarrow \text{semisimplicity} \parallel & & \downarrow \text{semisimplicity} \parallel \\
 \bigoplus_{c \in \text{Irr}(\text{CAT})} \text{Hom}(A \otimes B, c) \otimes \text{Hom}(c \otimes C, D) & \cong & \bigoplus_{d \in \text{Irr}(\text{Cat})} \text{Hom}(A \otimes d, D) \otimes \text{Hom}(B \otimes C, d) \\
 \parallel & & \parallel \\
 \bigoplus_c V_c^{AB} \otimes V_D^{cC} & \cong & \bigoplus_c V_D^{Ac} \otimes V_c^{BC} \\
 \cap & & \cap \\
 |(A, B); c; \mu\rangle \otimes |c, C; D; \nu\rangle & \xrightarrow[\text{6j-symbols}]{F_D^{ABC}} & \sum_{f, \alpha, \beta} [F_D^{ABC}]_{(c\mu\nu)}^{(f\alpha\beta)} |A, f; D; \beta\rangle \otimes |B, C; f; \alpha\rangle
 \end{array}$$

The diagrammatic equation shows two tree diagrams representing the same operation. The left tree has three input lines labeled A, B, and C. Lines A and B are connected at a vertex labeled μ. The output of this vertex is a line labeled ν, which then connects to line C at a second vertex labeled e. The final output line is labeled D. The right tree also has three input lines labeled A, B, and C. Lines A and B are connected at a vertex labeled v1. The output of this vertex is a line labeled i, which then connects to line C at a second vertex labeled v2. The final output line is labeled D. The two trees are separated by an equals sign, and the sum over f, α, β of the 6j symbol [F_D^{ABC}]_{(eμν)}^{(fαβ)} is placed to the left of the right tree.

10 Appendix

Implementations that are available:

Name	Symmetry	Citation(Software, Publication)	Language
ITensor	Nonabelian (Fermion?)	[FWS22]	Julia
Cyten	Categorical	[UH] [Unf24]	python
TeNPy	Abelian	[Hau+24]	python
QSpace	Nonabelian	[Wei24] [Wei12a]	?
ChemTensor	$SU(2)$	[Men26]	C
UniTensor	?	[<empty citation>]	? (Python, C++)
YasTn	Abelian	[Ram+25]	Python
TensorKit.jl	Categorical	[DH25]	Julia
Cytnx	Abelian only (i think)	[<empty citation>]	C++
quimb	Fermion?	[<empty citation>]	julia
MPSKit.jl	like TensorKit?	[<empty citation>]	julia
TeMFpy	Fermion?	[HS25]	python
abeliantensor	$U(1)$ and $\mathbb{Z}_n$	[Hau26]	python
symmray	Fermion	[Gao+25] [Gra26]	?
grassmanntn	Grassmann	[Yos26]	?
SymLibrary	? (OnSite)	[McM19] [MSB20]	MatLab?
Telum.jl	NonAbelian	[Lee26]	Julia

Table 1: Overview of tensor network libraries and symmetry support. (There are some closed source things here as well, also what is the software used by singh et al.). Software is included if it has a website or documentation and an account of the symmetries it exploits. Applications or theory are listed in the Citation column.

References

- [AK23] Fatimah Rita Ahmadi and Aleks Kissinger. “The ZX-calculus as a Language for Topological Quantum Computation”. In: *Journal of Physics A: Mathematical and Theoretical* 56.41 (Oct. 13, 2023), p. 415301. DOI: [10.1088/1751-8121/acef7e](https://doi.org/10.1088/1751-8121/acef7e). arXiv: [2211.03855](https://arxiv.org/abs/2211.03855) [quant-ph]. (Visited on 11/21/2024) (cit. on p. 3).
- [Aye17] Babatunde Moses Ayeni. “Studies of Braided Non-Abelian Anyons Using Anyonic Tensor Networks”. PhD thesis. Macquarie University, Dec. 3, 2017. DOI: [10.25949/19435697.v1](https://doi.org/10.25949/19435697.v1). arXiv: [1708.06476](https://arxiv.org/abs/1708.06476) [quant-ph]. (Visited on 06/28/2026) (cit. on p. 10).
- [BG17] Matthew Buican and Andrey Gromov. “Anyonic Chains, Topological Defects, and Conformal Field Theory”. In: *Communications in Mathematical Physics* 356.3 (Dec. 1, 2017), pp. 1017–1056. DOI: [10.1007/s00220-017-2995-6](https://doi.org/10.1007/s00220-017-2995-6). arXiv: [1701.02800v2](https://arxiv.org/abs/1701.02800v2) [hep-th]. (Visited on 06/28/2026) (cit. on pp. 2, 10).
- [BKP17] Parsa Bonderson, Christina Knapp, and Kaushal Patel. “Anyonic Entanglement and Topological Entanglement Entropy”. In: *Annals of Physics* 385 (Oct. 1, 2017), pp. 399–468. DOI: [10.1016/j.aop.2017.07.018](https://doi.org/10.1016/j.aop.2017.07.018). (Visited on 07/05/2026) (cit. on p. 2).
- [BSS08] Parsa Bonderson, Kirill Shtengel, and J. K. Slingerland. “Interferometry of Non-Abelian Anyons”. In: *Annals of Physics* 323.11 (Nov. 1, 2008), pp. 2709–2755. DOI: [10.1016/j.aop.2008.01.012](https://doi.org/10.1016/j.aop.2008.01.012). (Visited on 07/26/2026) (cit. on p. 2).
- [Bul+17] Nick Bultinck et al. “Fermionic Matrix Product States and One-Dimensional Topological Phases”. In: *Physical Review B* 95.7 (Feb. 3, 2017), p. 075108. DOI: [10.1103/PhysRevB.95.075108](https://doi.org/10.1103/PhysRevB.95.075108). arXiv: [1610.07849v2](https://arxiv.org/abs/1610.07849v2) [cond-mat.str-el]. (Visited on 05/04/2026) (cit. on p. 8).
- [Bur16] Simon Burton. *A Short Guide to Anyons and Modular Functors*. Oct. 17, 2016. DOI: [10.48550/arXiv.1610.05384](https://doi.org/10.48550/arXiv.1610.05384). arXiv: [1610.05384](https://arxiv.org/abs/1610.05384) [quant-ph]. (Visited on 07/15/2026). Pre-published (cit. on p. 2).
- [Cir+17] J. I. Cirac et al. “Matrix Product Density Operators: Renormalization Fixed Points and Boundary Theories”. In: *Annals of Physics* 378 (Mar. 1, 2017), pp. 100–149. DOI: [10.1016/j.aop.2016.12.030](https://doi.org/10.1016/j.aop.2016.12.030). arXiv: [1606.00608v4](https://arxiv.org/abs/1606.00608v4) [quant-ph] (cit. on p. 8).

Symbol	Meaning
$\mathbb{C}$	Complex numbers
$\mathbb{F}$	Base field
$\mathbb{N}$	Natural numbers
$\mathfrak{g}$	Lie algebra
$\mathfrak{sl}_2$	Special linear Lie algebra of degree 2
$U_q(\mathfrak{sl}_2)$	Quantum group associated with $\mathfrak{sl}_2$
$\mathcal{C}$	Category
$\mathbf{1}$	Monoidal unit object
V^*	Dual vector space/object
$\text{Hom}(V, W)$	Space of morphisms from V to W
$\text{Rep}(G)$	Representation category of group G
$\text{Hom}_G(V, W)$	Space of intertwiners between representations of G
Vect_k	Category of k -vector spaces
$\text{Hom}_k(-, -)$	Linear morphisms between k -vector spaces

Table 2: Notation conventions.

- [DH25] Lukas Devos and Jutho Haegeman. *TensorKit.jl: A Julia Package for Large-Scale Tensor Computations, with a Hint of Category Theory*. Version 1. 2025. DOI: [10.48550/arxiv.2508.10076](https://doi.org/10.48550/arxiv.2508.10076). (Visited on 06/01/2026). Pre-published (cit. on pp. [1](#), [2](#), [7](#), [10](#), [13](#), [23](#)).
- [DN05] Christopher M. Dawson and Michael A. Nielsen. *The Solovay-Kitaev Algorithm*. Aug. 23, 2005. DOI: [10.48550/arXiv.quant-ph/0505030](https://doi.org/10.48550/arXiv.quant-ph/0505030). arXiv: [quant-ph/0505030](https://arxiv.org/abs/quant-ph/0505030). (Visited on 07/15/2026). Pre-published (cit. on pp. [2](#), [3](#)).
- [DT24] Clement Delcamp and Apoorv Tiwari. “Higher Categorical Symmetries and Gauging in Two-Dimensional Spin Systems”. In: *SciPost Physics* 16.4 (Apr. 23, 2024), p. 110. DOI: [10.21468/SciPostPhys.16.4.110](https://doi.org/10.21468/SciPostPhys.16.4.110). arXiv: [2301.01259 \[hep-th\]](https://arxiv.org/abs/2301.01259). (Visited on 06/08/2026) (cit. on p. [3](#)).
- [Eck25] Luisa Eck. “Quantum Lattice Models from Fusion Categories”. PhD thesis. University of Oxford, 2025. DOI: [10.5287/ora-zozm87o5w](https://doi.org/10.5287/ora-zozm87o5w). (Visited on 06/28/2026) (cit. on p. [10](#)).
- [Fei+07] Adrian Feiguin et al. “Interacting Anyons in Topological Quantum Liquids: The Golden Chain”. In: *Physical Review Letters* 98.16 (Apr. 20, 2007), p. 160409. DOI: [10.1103/PhysRevLett.98.160409](https://doi.org/10.1103/PhysRevLett.98.160409). (Visited on 06/28/2026) (cit. on pp. [2](#), [9](#)).
- [FNW92] M. Fannes, B. Nachtergaele, and R. F. Werner. “Finitely Correlated States on Quantum Spin Chains”. In: *Communications in Mathematical Physics* 144.3 (Mar. 1, 1992), pp. 443–490. DOI: [10.1007/BF02099178](https://doi.org/10.1007/BF02099178). (Visited on 07/15/2026) (cit. on p. [1](#)).
- [Fre98] Michael H. Freedman. “P/NP, and the Quantum Field Computer”. In: *Proceedings of the National Academy of Sciences* 95.1 (Jan. 6, 1998), pp. 98–101. DOI: [10.1073/pnas.95.1.98](https://doi.org/10.1073/pnas.95.1.98). (Visited on 07/25/2026) (cit. on p. [2](#)).
- [FWS22] Matthew Fishman, Steven R. White, and E. Miles Stoudenmire. “The ITensor Software Library for Tensor Network Calculations”. In: *SciPost Phys. Codebases* (2022), p. 4. DOI: [10.21468/SciPostPhysCodeb.4](https://doi.org/10.21468/SciPostPhysCodeb.4) (cit. on p. [23](#)).
- [Gao+25] Yang Gao et al. “Fermionic Tensor Network Contraction for Arbitrary Geometries”. In: *Physical Review Research* 7.2 (May 27, 2025), p. 023193. DOI: [10.1103/PhysRevResearch.7.023193](https://doi.org/10.1103/PhysRevResearch.7.023193). (Visited on 05/04/2026) (cit. on pp. [8](#), [23](#)).
- [Gil+13] C. Gils et al. “Anyonic Quantum Spin Chains: Spin-1 Generalizations and Topological Stability”. In: *Physical Review B* 87.23 (June 17, 2013), p. 235120. DOI: [10.1103/PhysRevB.87.235120](https://doi.org/10.1103/PhysRevB.87.235120). (Visited on 06/28/2026) (cit. on pp. [2](#), [10](#)).
- [Gra26] Johnnie Gray. *Jcmgray/Symmray*. June 21, 2026. URL: <https://github.com/jcmgray/symmray> (visited on 06/30/2026) (cit. on p. [23](#)).

- [Hau+24] Johannes Hauschild et al. “Tensor Network Python (TeNPy) Version 1”. In: *SciPost Phys. Codebases* (2024), p. 41. doi: [10.21468/SciPostPhysCodeb.41](https://doi.org/10.21468/SciPostPhysCodeb.41) (cit. on p. 23).
- [Hau26] Markus Hauru. *Mhauru/Abeliantensors*. May 16, 2026. URL: <https://github.com/mhauru/abeliantensors> (visited on 06/30/2026) (cit. on p. 23).
- [HJ12] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. 2nd ed. Cambridge: Cambridge University Press, 2012. ISBN: 978-1-139-02041-1. doi: [10.1017/CBO9781139020411](https://doi.org/10.1017/CBO9781139020411). (Visited on 06/30/2026) (cit. on p. 7).
- [HS25] Simon H. Hille and Attila Szabó. *TeMFpy: A Python Library for Converting Fermionic Mean-Field States into Tensor Networks*. 2025. arXiv: [2510.05227](https://arxiv.org/abs/2510.05227) [[cond-mat.str-el](#)] (cit. on p. 23).
- [IR19] Alberto Ibort and Miguel A. Rodriguez. *An Introduction to Groups, Groupoids and Their Representations*. Boca Raton: CRC Press, Oct. 28, 2019. 362 pp. ISBN: 978-1-315-23294-2. doi: [10.1201/b22019](https://doi.org/10.1201/b22019) (cit. on p. 20).
- [Kir+23] Nico Kirchner et al. “Numerical Simulation of Non-Abelian Anyons”. In: *Physical Review B* 107.19 (May 15, 2023), p. 195129. doi: [10.1103/PhysRevB.107.195129](https://doi.org/10.1103/PhysRevB.107.195129). arXiv: [2206.14730](https://arxiv.org/abs/2206.14730) [[cond-mat.str-el](#)]. (Visited on 07/16/2026) (cit. on p. 2).
- [Kit03] A. Yu Kitaev. “Fault-Tolerant Quantum Computation by Anyons”. In: *Annals of Physics* 303.1 (Jan. 2003), pp. 2–30. doi: [10.1016/S0003-4916\(02\)00018-0](https://doi.org/10.1016/S0003-4916(02)00018-0). arXiv: [quant-ph/9707021](https://arxiv.org/abs/quant-ph/9707021). (Visited on 07/17/2026) (cit. on p. 2).
- [Kit06] Alexei Kitaev. “Anyons in an Exactly Solved Model and Beyond”. In: *Annals of Physics* 321.1 (Jan. 2006), pp. 2–111. doi: [10.1016/j.aop.2005.10.005](https://doi.org/10.1016/j.aop.2005.10.005). arXiv: [cond-mat/0506438v3](https://arxiv.org/abs/cond-mat/0506438v3). (Visited on 07/14/2026) (cit. on pp. 2, 11).
- [Kit97] A Yu Kitaev. “Quantum Computations: Algorithms and Error Correction”. In: *Russian Mathematical Surveys* 52.6 (Dec. 31, 1997), pp. 1191–1249. doi: [10.1070/RM1997v052n06ABEH002155](https://doi.org/10.1070/RM1997v052n06ABEH002155). (Visited on 07/26/2026) (cit. on p. 3).
- [Kon+20] Liang Kong et al. “Algebraic Higher Symmetry and Categorical Symmetry: A Holographic and Entanglement View of Symmetry”. In: *Physical Review Research* 2.4 (Oct. 15, 2020), p. 043086. doi: [10.1103/PhysRevResearch.2.043086](https://doi.org/10.1103/PhysRevResearch.2.043086). (Visited on 07/05/2026) (cit. on p. 3).
- [KZ22] Liang Kong and Zhi-Hao Zhang. *An Invitation to Topological Orders and Category Theory*. May 31, 2022. doi: [10.48550/arXiv.2205.05565](https://doi.org/10.48550/arXiv.2205.05565). arXiv: [2205.05565](https://arxiv.org/abs/2205.05565) [[cond-mat.str-el](#)]. (Visited on 07/02/2026). Pre-published (cit. on pp. 1, 3, 16).
- [KZ23] Liang Kong and Hao Zheng. “Categorical Computation”. In: *Frontiers of Physics* 18.2 (Apr. 2023), p. 21302. doi: [10.1007/s11467-022-1251-5](https://doi.org/10.1007/s11467-022-1251-5). arXiv: [2102.04814](https://arxiv.org/abs/2102.04814) [[quant-ph](#)]. (Visited on 07/01/2026) (cit. on p. 2).
- [Lee26] Seung-Sup Lee. *Ssblee/Telum.jl*. June 30, 2026. URL: <https://github.com/ssblee/Telum.jl> (visited on 07/15/2026) (cit. on p. 23).
- [Liu+22] Yu-Jie Liu et al. “Methods for Simulating String-Net States and Anyons on a Digital Quantum Computer”. In: *PRX Quantum* 3.4 (Nov. 7, 2022), p. 040315. doi: [10.1103/PRXQuantum.3.040315](https://doi.org/10.1103/PRXQuantum.3.040315). (Visited on 07/16/2026) (cit. on p. 2).
- [LM77] J. M. Leinaas and J. Myrheim. “On the Theory of Identical Particles”. In: *Il Nuovo Cimento B* (1971-1996) 37.1 (Jan. 1, 1977), pp. 1–23. doi: [10.1007/BF02727953](https://doi.org/10.1007/BF02727953). (Visited on 07/19/2026) (cit. on p. 2).
- [McM19] Nathan McMahon. *Qnla/SymLibrary*. Mar. 4, 2019. URL: <https://github.com/qnla/SymLibrary> (visited on 07/05/2026) (cit. on p. 23).
- [Men26] Christian B. Mendl. *Qc-Tum/Chemtensor*. qc-tum, May 28, 2026. URL: <https://github.com/qc-tum/chemtensor> (visited on 06/30/2026) (cit. on p. 23).
- [MG02] I. P. McCulloch and M. Gulácsi. “The Non-Abelian Density Matrix Renormalization Group Algorithm”. In: *Europhysics Letters* 57.6 (Mar. 1, 2002), p. 852. doi: [10.1209/epl/i2002-00393-0](https://doi.org/10.1209/epl/i2002-00393-0). (Visited on 07/16/2026) (cit. on p. 1).

- [Moc03] Carlos Mochon. “Anyons from Nonsolvable Finite Groups Are Sufficient for Universal Quantum Computation”. In: *Physical Review A* 67.2 (2003). doi: [10.1103/PhysRevA.67.022315](#) (cit. on p. 2).
- [Moc04] Carlos Mochon. “Anyon Computers with Smaller Groups”. In: *Physical Review A* 69.3 (Mar. 11, 2004), p. 032306. doi: [10.1103/PhysRevA.69.032306](#). arXiv: [quant-ph/0306063](#). (Visited on 07/01/2026) (cit. on p. 2).
- [Mor+25] Quinten Mortier et al. “Fermionic Tensor Network Methods”. In: *SciPost Physics* 18.1 (Jan. 10, 2025), p. 012. doi: [10.21468/SciPostPhys.18.1.012](#). (Visited on 05/04/2026) (cit. on p. 8).
- [MSB20] Nathan A. McMahon, Sukhbinder Singh, and Gavin K. Brennen. “A Holographic Duality from Lifted Tensor Networks”. In: *npj Quantum Information* 6.1 (Apr. 24, 2020), p. 36. doi: [10.1038/s41534-020-0255-7](#). (Visited on 07/05/2026) (cit. on p. 23).
- [MSS24] David Jaz Myers, Hisham Sati, and Urs Schreiber. “Topological Quantum Gates in Homotopy Type Theory”. In: *Communications in Mathematical Physics* 405.7 (July 2024), p. 172. doi: [10.1007/s00220-024-05020-8](#). arXiv: [2303.02382 \[quant-ph\]](#). (Visited on 07/17/2026) (cit. on p. 2).
- [Nay+08] Chetan Nayak et al. “Non-Abelian Anyons and Topological Quantum Computation”. In: *Reviews of Modern Physics* 80.3 (Sept. 12, 2008), pp. 1083–1159. doi: [10.1103/RevModPhys.80.1083](#). arXiv: [0707.1889 \[cond-mat.str-el\]](#). (Visited on 07/17/2026) (cit. on p. 2).
- [OR95] Stellan Ostlund and Stefan Rommer. “Thermodynamic Limit of the Density Matrix Renormalization for the Spin-1 Heisenberg Chain”. In: *Physical Review Letters* 75.19 (Nov. 6, 1995), pp. 3537–3540. doi: [10.1103/PhysRevLett.75.3537](#). arXiv: [cond-mat/9503107](#). (Visited on 07/15/2026) (cit. on p. 1).
- [Pen71] Roger Penrose. “Applications of Negative Dimensional Tensors”. In: (1971). Ed. by D.J.A. Welsh, pp. 221–224. URL: <https://ncatlab.org/nlab/files/PenroseNegativeDimensionalT.pdf> (visited on 06/30/2026) (cit. on p. 1).
- [Per+08] D. Perez-Garcia et al. “String Order and Symmetries in Quantum Spin Lattices”. In: *Physical Review Letters* 100.16 (Apr. 22, 2008), p. 167202. doi: [10.1103/PhysRevLett.100.167202](#). arXiv: [0802.0447 \[cond-mat.str-el\]](#). (Visited on 06/28/2026) (cit. on p. 8).
- [Pfe+10] Robert N. C. Pfeifer et al. “Simulation of Anyons with Tensor Network Algorithms”. In: *Physical Review B* 82.11 (Sept. 29, 2010), p. 115126. doi: [10.1103/PhysRevB.82.115126](#). arXiv: [1006.3532 \[cond-mat.str-el\]](#). (Visited on 06/16/2026) (cit. on pp. 1, 2, 10).
- [Pfe12] Robert N. C. Pfeifer. *Simulation of Anyons Using Symmetric Tensor Network Algorithms*. Feb. 10, 2012. doi: [10.48550/arXiv.1202.1522](#). arXiv: [1202.1522 \[cond-mat\]](#). (Visited on 05/11/2026). Pre-published (cit. on pp. 2, 10).
- [Pfe14] Robert N. C. Pfeifer. “Measures of Entanglement in Non-Abelian Anyonic Systems”. In: *Physical Review B* 89.3 (Jan. 6, 2014), p. 035105. doi: [10.1103/PhysRevB.89.035105](#). (Visited on 07/05/2026) (cit. on p. 2).
- [Ram+25] Marek M. Rams et al. “YASTN: Yet Another Symmetric Tensor Networks; A Python Library for Abelian Symmetric Tensor Network Calculations”. In: *SciPost Phys. Codebases* (2025), p. 52. doi: [10.21468/SciPostPhysCodeb.52](#) (cit. on p. 23).
- [RT91] N. Reshetikhin and V. G. Turaev. “Invariants of 3-Manifolds via Link Polynomials and Quantum Groups”. In: *Inventiones mathematicae* 103.1 (Dec. 1, 1991), pp. 547–597. doi: [10.1007/BF01239527](#). (Visited on 07/15/2026) (cit. on p. 2).
- [Sch+20] Philipp Schmoll et al. “A Programming Guide for Tensor Networks with Global $U(2)$ Symmetry”. In: *Annals of Physics* 419 (Aug. 1, 2020), p. 168232. doi: [10.1016/j.aop.2020.168232](#). (Visited on 05/07/2026) (cit. on p. 2).

- [Sch05] Issai Schur. “Neue Begründungen der Theorie der Gruppencharaktere”. In: *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* (Januar bis Juni 1905), pp. 406–433. URL: http://archive.org/details/bub_gb_KwUoAAAAYAAJ (visited on 07/15/2026) (cit. on p. 20).
- [Sch20] Philipp Schmoll. “Tensor Network Simulations with Global SU(2) Symmetry”. Mainz: Johannes Gutenberg University, July 2020. 150 pp. URL: <https://openscience.ub.uni-mainz.de/server/api/core/bitstreams/d9d9ca4e-46c8-4a8e-a831-7611475d2504/content> (visited on 05/08/2026) (cit. on p. 2).
- [Sel11] P. Selinger. “A Survey of Graphical Languages for Monoidal Categories”. In: *New Structures for Physics*. Ed. by Bob Coecke. Berlin, Heidelberg: Springer, 2011, pp. 289–355. ISBN: 978-3-642-12821-9. DOI: [10.1007/978-3-642-12821-9_4](https://doi.org/10.1007/978-3-642-12821-9_4). arXiv: [0908.3347v1](https://arxiv.org/abs/0908.3347v1) [math.CT] (cit. on p. 3).
- [Sim+06] S. H. Simon et al. “Topological Quantum Computing with Only One Mobile Quasiparticle”. In: *Physical Review Letters* 96.7 (Feb. 23, 2006), p. 070503. DOI: [10.1103/PhysRevLett.96.070503](https://doi.org/10.1103/PhysRevLett.96.070503). arXiv: [quant-ph/0509175](https://arxiv.org/abs/quant-ph/0509175). (Visited on 07/19/2026) (cit. on p. 3).
- [Sim23] Steven H. Simon. *Topological Quantum*. Oxford University Press, Sept. 29, 2023. ISBN: 978-0-19-888672-3. DOI: [10.1093/oso/9780198886723.001.0001](https://doi.org/10.1093/oso/9780198886723.001.0001). (Visited on 06/24/2026) (cit. on pp. 2, 22).
- [Sin+14] Sukhwinder Singh et al. “Matrix Product States for Anyonic Systems and Efficient Simulation of Dynamics”. In: *Physical Review B* 89.7 (Feb. 13, 2014), p. 075112. DOI: [10.1103/PhysRevB.89.075112](https://doi.org/10.1103/PhysRevB.89.075112). (Visited on 07/08/2026) (cit. on p. 11).
- [SM98] G. Sierra and M. A. Martin-Delgado. *The Density Matrix Renormalization Group, Quantum Groups and Conformal Field Theory*. Version 3. Nov. 30, 1998. DOI: [10.48550/arXiv.cond-mat/9811170](https://doi.org/10.48550/arXiv.cond-mat/9811170). arXiv: [cond-mat/9811170](https://arxiv.org/abs/cond-mat/9811170). (Visited on 07/16/2026). Pre-published (cit. on p. 1).
- [SMB18] Sukhwinder Singh, Nathan A. McMahon, and Gavin K. Brennen. “Holographic Spin Networks from Tensor Network States”. In: *Physical Review D* 97.2 (Jan. 26, 2018), p. 026013. DOI: [10.1103/PhysRevD.97.026013](https://doi.org/10.1103/PhysRevD.97.026013). arXiv: [1702.00392](https://arxiv.org/abs/1702.00392) [cond-mat.str-el]. (Visited on 05/11/2026) (cit. on p. 1).
- [SO20] Philipp Schmoll and Roman Orus. “Benchmarking Global SU(2) Symmetry in 2d Tensor Network Algorithms”. In: *Physical Review B* 102.24 (Dec. 1, 2020), p. 241101. DOI: [10.1103/PhysRevB.102.241101](https://doi.org/10.1103/PhysRevB.102.241101). arXiv: [2005.02748](https://arxiv.org/abs/2005.02748) [cond-mat]. (Visited on 05/07/2026) (cit. on p. 2).
- [SPV10] Sukhwinder Singh, Robert N. C. Pfeifer, and Guifre Vidal. “Tensor Network Decompositions in the Presence of a Global Symmetry”. In: *Physical Review A* 82.5 (Nov. 8, 2010), p. 050301. DOI: [10.1103/PhysRevA.82.050301](https://doi.org/10.1103/PhysRevA.82.050301). arXiv: [0907.2994](https://arxiv.org/abs/0907.2994) [cond-mat.str-el]. (Visited on 05/11/2026) (cit. on pp. 2, 5).
- [SPV11] Sukhwinder Singh, Robert N. C. Pfeifer, and Guifre Vidal. “Tensor Network States and Algorithms in the Presence of a Global U(1) Symmetry”. In: *Physical Review B* 83.11 (Mar. 15, 2011), p. 115125. DOI: [10.1103/PhysRevB.83.115125](https://doi.org/10.1103/PhysRevB.83.115125). arXiv: [1008.4774v1](https://arxiv.org/abs/1008.4774v1) [cond-mat.str-el]. (Visited on 05/06/2026) (cit. on p. 2).
- [SS26] Hisham Sati and Urs Schreiber. “Entanglement of Sections: The Pushout of Entangled and Parameterized Quantum Information”. In: *Quantum Studies: Mathematics and Foundations* 13.2 (June 2026), p. 23. DOI: [10.1007/s40509-026-00397-8](https://doi.org/10.1007/s40509-026-00397-8). arXiv: [2309.07245](https://arxiv.org/abs/2309.07245) [quant-ph]. (Visited on 07/19/2026) (cit. on p. 3).
- [SV12] Sukhwinder Singh and Guifre Vidal. “Tensor Network States and Algorithms in the Presence of a Global SU(2) Symmetry”. In: *Physical Review B* 86.19 (Nov. 9, 2012), p. 195114. DOI: [10.1103/PhysRevB.86.195114](https://doi.org/10.1103/PhysRevB.86.195114). arXiv: [1208.3919v1](https://arxiv.org/abs/1208.3919v1) [cond-mat.str-el]. (Visited on 05/11/2026) (cit. on pp. 1, 2, 13, 16).

- [Tre+08] Simon Trebst et al. “A Short Introduction to Fibonacci Anyon Models”. In: *Progress of Theoretical Physics Supplement* 176 (June 1, 2008), pp. 384–407. doi: [10.1143/PTPS.176.384](https://doi.org/10.1143/PTPS.176.384). arXiv: [0902.3275v1](https://arxiv.org/abs/0902.3275v1) [[cond-mat.stat-mech](#)]. (Visited on 06/28/2026) (cit. on pp. [2](#), [10](#)).
- [UH] Jakob Unfried and Johannes Michael Hauschild. *Cyten* (cit. on p. [23](#)).
- [Unf24] Jakob Unfried. “Advancing Tensor Network Simulations -”. PhD thesis. Technische Universität München, 2024, p. 208. URL: <https://mediatum.ub.tum.de/1755273> (cit. on pp. [1](#), [2](#), [10](#), [23](#)).
- [Val21] Sachin Jayesh Valera. “Topological Quantum and Skein-Theoretic Aspects of Braided Fusion Categories”. PhD thesis. Norway: University of Bergen, 2021. ISBN: 978-82-308-5158-6 978-82-308-6990-1. URL: [11250/2770245](https://nbn-resolving.org/urn:nbn:no:bi-11250-2770245) (visited on 06/08/2026) (cit. on p. [2](#)).
- [VC04] F. Verstraete and J. I. Cirac. *Renormalization Algorithms for Quantum-Many Body Systems in Two and Higher Dimensions*. July 2, 2004. doi: [10.48550/arXiv.cond-mat/0407066](https://doi.org/10.48550/arXiv.cond-mat/0407066). arXiv: [cond-mat/0407066](https://arxiv.org/abs/cond-mat/0407066). (Visited on 07/15/2026). Pre-published (cit. on p. [1](#)).
- [Vid08] G. Vidal. “A Class of Quantum Many-Body States That Can Be Efficiently Simulated”. In: *Physical Review Letters* 101.11 (Sept. 12, 2008), p. 110501. doi: [10.1103/PhysRevLett.101.110501](https://doi.org/10.1103/PhysRevLett.101.110501). arXiv: [quant-ph/0610099](https://arxiv.org/abs/quant-ph/0610099). (Visited on 07/15/2026) (cit. on p. [1](#)).
- [VW18] Gert Vercleyen, Frank Verstraete, and Universiteit Gent. Faculteit Wetenschappen. “The Mathematical Structure of Tensor Networks”. MA thesis. Master of Science in de wiskunde, 2018. URL: https://lib.ugent.be/fulltxt/RUG01/002/479/664/RUG01-002479664_2018_0001_AC.pdf (cit. on pp. [3](#), [8](#)).
- [Wei12a] Andreas Weichselbaum. “Non-Abelian Symmetries in Tensor Networks: A Quantum Symmetry Space Approach”. In: *Annals of Physics* 327.12 (Dec. 1, 2012), pp. 2972–3047. doi: [10.1016/j.aop.2012.07.009](https://doi.org/10.1016/j.aop.2012.07.009). (Visited on 05/07/2026) (cit. on pp. [2](#), [23](#)).
- [Wei12b] Andreas Weichselbaum. “Tensor Networks and the Numerical Renormalization Group”. In: *Physical Review B* 86.24 (Dec. 20, 2012), p. 245124. doi: [10.1103/PhysRevB.86.245124](https://doi.org/10.1103/PhysRevB.86.245124). (Visited on 06/30/2026) (cit. on p. [1](#)).
- [Wei20] Andreas Weichselbaum. “X-Symbols for Non-Abelian Symmetries in Tensor Networks”. In: *Physical Review Research* 2.2 (June 23, 2020), p. 023385. doi: [10.1103/PhysRevResearch.2.023385](https://doi.org/10.1103/PhysRevResearch.2.023385). (Visited on 06/30/2026) (cit. on p. [2](#)).
- [Wei24] Andreas Weichselbaum. “QSpace - An Open-Source Tensor Library for Abelian and Non-Abelian Symmetries”. In: *SciPost Physics Codebases* (Nov. 14, 2024), p. 40. doi: [10.21468/SciPostPhysCodeb.40](https://doi.org/10.21468/SciPostPhysCodeb.40). arXiv: [2405.06632v1](https://arxiv.org/abs/2405.06632v1) [[cond-mat.str-el](#)]. (Visited on 05/08/2026) (cit. on p. [23](#)).
- [Whi92] Steven R. White. “Density Matrix Formulation for Quantum Renormalization Groups”. In: *Physical Review Letters* 69.19 (Nov. 9, 1992), pp. 2863–2866. doi: [10.1103/PhysRevLett.69.2863](https://doi.org/10.1103/PhysRevLett.69.2863). (Visited on 07/15/2026) (cit. on p. [1](#)).
- [Whi93] Steven R. White. “Density-Matrix Algorithms for Quantum Renormalization Groups”. In: *Physical Review B* 48.14 (Oct. 1, 1993), pp. 10345–10356. doi: [10.1103/PhysRevB.48.10345](https://doi.org/10.1103/PhysRevB.48.10345). (Visited on 07/15/2026) (cit. on p. [1](#)).
- [Wil90] Frank Wilczek. *Fractional Statistics and Anyon Superconductivity*. WORLD SCIENTIFIC, 1990. doi: [10.1142/0961](https://doi.org/10.1142/0961). eprint: <https://www.worldscientific.com/doi/pdf/10.1142/0961> (cit. on p. [2](#)).
- [Yos26] Atis Yosprakob. *Ayosprakob/Grassmannn*. June 7, 2026. URL: <https://github.com/ayosprakob/grassmannn> (visited on 06/30/2026) (cit. on p. [23](#)).