Unbounded central charge at fixed local dimension?

A candidate refutation of the c–d conjecture, prepared for expert review

Compiled for J. Kudler-Flam (Institute for Advanced Study)
from the repository `github.com/jkudlerflam/cd-conjecture`, folder `notes/cd_conjecture`

28 October 2026, revised 29 October 2026

Abstract

Latorre and Sierra conjecture that a one-site translation-invariant, nearest-neighbour chain on $(\mathbb{C}^d)^{\otimes L}$ whose low-energy spectrum is a collection of unitary CFTs has total central charge $c \le d-1$. We describe a construction that, for every n, gives a Hermitian nearest-neighbour bond on $\mathbb{C}^{4464} \otimes \mathbb{C}^{4464}$ whose predicted infrared theory is $n+1$ compact bosons, so no bound of the form $c \le f(d)$ would hold. A related 24-state chain is predicted to have $c = 24 > d-1$. Every finite-dimensional ingredient is proved, or certified in exact arithmetic at small n and checked in high precision beyond. The infrared conclusion rests on one standard but unproved input S, the elimination of a massive local sector at fixed coupling followed by weak-coupling RG for a multi-component Luttinger liquid with extremely separated scales. Since the first version, the elimination step, uniformly in volume, and the transfer to the physical ring in thermodynamic form have been proved at fixed n, and S now reduces to two constructive-RG estimates about the logical fermion action, neither of them closed (Section 6). An independent rebuild of the wrapper agrees with the original, and its projected $n = 1$ chain gives a calibrated $c = 2$. A logical vacancy model gives calibrated $c = 2$ and 2.9 ± 0.1 at $n = 1, 2$, and a ten-letter demonstrator flows to the Luttinger liquid of its matched logical chain, as S predicts. An eleven-letter alphabet gives refereed conditional candidates at $d = 11$ (frozen, $c = 11$) and $d = 13$ (mobile, $c = 14$). No chain with $c > d-1$ has been simulated. The work was produced with AI assistance and has not been refereed by humans. This note states what a reviewer would need to check, and in which order.

1 The conjecture and the claim

Latorre and Sierra [1] consider

$$H = \sum_{i=1}^{L} h_{i,i+1}, \qquad h = h^\dagger \text{ on } \mathbb{C}^d \otimes \mathbb{C}^d, \qquad h_{L,L+1} \equiv h_{L,1}, \tag{1}$$

with one fixed bond h on every link. If the low-energy spectrum is described by a collection of unitary CFTs with central charges c_a, possibly with different velocities, they conjecture

$$c_T = \sum_a c_a \le d - 1. \tag{2}$$

The bound is saturated by the Uimin–Lai–Sutherland chain $h = P$ (the swap), whose infrared theory is $SU(d)_1$ [2]. Blocking k sites raises the local dimension to d^k, a remark in [1] that will matter below. A weaker question is whether any bound $c \le f(d)$ exists. Known proofs cover restricted classes (quasi-free chains obey $c \le \log_2 d$, for example), and every general lattice quantity we know that bounds c grows like a length scale times $\log d$.

In this note, we describe two forms of one construction.

Claim A (unbounded c at fixed d)**.** For every $n \geq 1$ there is a Hermitian bond h_n on $\mathbb{C}^{d_0} \otimes \mathbb{C}^{d_0}$, $d_0 = 4464$, built from one fixed alphabet and one fixed list of terms with n-dependent coefficients, such that, assuming the input S of Section 5.3, the periodic chain (1) has on every ring with $L \geq (n+3)(n+4)$ an infrared theory of $n+1$ compact bosons with positive velocities, so $c_T = n+1$. Hence c_T is unbounded at $d = 4464$, and $c \leq d-1$ itself fails once $n \geq 4463$.

Claim B (a concrete instance of $c > d-1$)**.** An explicit bond h_{24} on $\mathbb{C}^{24} \otimes \mathbb{C}^{24}$, at a certified parameter point with $n = 23$, is predicted to have $c_T = 24 > d - 1 = 23$. The prediction assumes the same kind of input as Claim A for a frozen 23-letter chain, plus a matching statement for a gapless vacancy fluid and a positive dressed defect energy (Section 3.2).

Claim A is the stronger statement and the one we ask reviewers to focus on. Claim B is cheaper and more concrete, but it has more open inputs. The finite-dimensional ingredients of both are established at the working standard of theoretical physics (proofs in the repository, exact rational or interval certificates where they exist, high-precision numerics otherwise). The infrared step of Claim A has been reduced, with explicit constants, to the single input S, and S has since been reduced to two named estimates (Section 6). In our assessment both claims are credible conditional candidates and neither is an established counterexample. The AI referee reports on the two forms reach the same verdict [48, 49]. Smaller conditional candidates of the same kind, down to $d = 11$, are described in Section 7.

2 The mechanism

The idea fits in a paragraph. Sites group spontaneously into long cells. Each cell carries an exactly degenerate code of n logical fermion modes, which acts as one large effective site. Neighbouring cells interact only through the single bond at their junction, and only at second order in that bond, so the cells form an effective chain with a logical Hamiltonian we can design. We design a stable multi-component Luttinger liquid with $c = n$. Finally the cells are made mobile, so that translation is unbroken and the cell lengths contribute one more boson. Each step is described below in physical terms.

2.1 Spontaneous cells

Letters carry conserved types (marker, data, token, hole), so a configuration is a sequence of segments between markers. A walker confined to a segment of ℓ sites has a confinement energy of order $t[1 - \cos(\pi/(\ell+1))]$, which favours long segments, while a marker chemical potential favours short ones. The energy per site is then minimized at one chosen length b, with a selection margin of order b^{-4} in units of t. Wrong lengths and illegal words (pairs whose arrows do not match) are excluded by operator lower bounds, at any fixed small value of the inter-cell coupling [50, 22]. A ten-letter example forms cells of length three unaided in VUMPS [54].

2.2 Codes with idle ports

Inside a selected cell the Hamiltonian is a sum of positive terms whose common zero space is one fixed state of a "token" (a walker in its vacuum species) tensored with n free data bits. This is the code, of dimension 2^n, and it is frustration free. The two end sites of a cell, the ports, are touched by the inter-cell coupling λV only through operators that change the token species at the port from vacuum to excited. Their compression onto the code therefore vanishes, $PVP = 0$, and first order in λ is absent.

2.3 Second-order exchange

At order λ^2 the port excites the tokens of the two cells adjacent to a junction. Each excited token runs into its cell, where a number-conserving conversion of amplitude g exchanges one unit of charge with the data site j under it, flipping that data bit, and then returns. The port de-excites the pair. When the two conversions carry opposite charge, the net process moves one logical fermion from one cell to the next, with an amplitude set by return filters f_j of the excited token. A Jordan–Wigner sign carried by the neutral token makes this a fermion bilinear, and the total charge Q (data occupation plus token charge) is an exact on-site $U(1)$. At order $\lambda^2 g^2$ the effective chain is

$$H_{\text{eff}} = \sum_x \Big[\sum_{i,j=1}^{n} T_{ij}\, c^\dagger_{x,i} c_{x+1,j} + \text{h.c.} + \sum_{i,j} M_{ij}\, c^\dagger_{x,i} c_{x,j} \Big] + \text{const} + O(\lambda^2 g^4) + O(\lambda^3), \tag{3}$$

with T and M of order $\lambda^2 g^2$ and no three-cell term at order λ^2. Non-Gaussian terms first appear at order g^4. A large energy D on an "express" rail makes T a Gram matrix, $T \propto -D^{-(2n+4)}\Gamma$ with $\Gamma > 0$, so every band crosses a common chemical potential [19].

2.4 An incommensurate multi-flavour target with exact $U(1)$

The logical symbol $h(k) = Te^{ik} + T^\dagger e^{-ik} + M$ gives n bands. Write $p_{a,\pm}$ for the two Fermi points of band a and $\theta_a = (p_{a,+} + p_{a,-})/2$ for its sea centre. The target requires $2n$ simple Fermi points, distinct centres, and positive margins

$$\delta_2 = \min_{\alpha \neq \beta} |p_\alpha - p_\beta|, \qquad \delta_4 = \min_{\{\alpha,\beta\} \neq \{\gamma,\delta\}} \big| p_\alpha + p_\beta - p_\gamma - p_\delta \bmod 2\pi \big|, \tag{4}$$

where the minimum runs over unordered pairs with repetition, so it includes pair transfer between bands, exchange, and every umklapp. With exact $U(1)$, a number-conserving vertex is then either a density or current product (Gaussian and marginal), or oscillating with momentum at least δ_4, or of scaling dimension at least three. A relative change of the Gaussian stiffness of at most $1/4$ keeps those dimensions above $3\sqrt{3/5} \approx 2.32$, whatever the velocity ratios [23]. This is the standard stability of a non-resonant multi-component Luttinger liquid [9, 10]. Symmetric targets are avoided on purpose. $SU(N)_1$ and the $SO(2n)$ point sit on separatrices and run away under small anisotropy, which is the familiar fate of the $SU(3)$ point of the spin-1 chain [13].

The difficulty is the centres. A real hopping matrix pins every centre at 0, and pair transfer $2\theta_a - 2\theta_b = 0$ is then resonant. The 46-letter core of [19] solves this with two independent excited buses, with stub hoppings w and wv and port phases differing by $\pi/2$, which give

$$T = -J\big[\Gamma(w) + i\eta\,\Gamma(wv)\big], \qquad J = 2\lambda^2 g^2\, \frac{q^{2n+2}}{S^2}\, D^{-(2n+4)}, \qquad S = \sum_{\ell=0}^{n+1} q^{2\ell}, \tag{5}$$

where $q = (wu)^2$ is the bias of the vacuum walker, $J_0 = J/\lambda^2$ is the coupling-independent part, and

$$\Gamma_{ij}(w) = \int_0^\infty f_i(t) f_j(t)\, \mathrm{d}t, \qquad f_j(t) = \int_0^t \big[e^{-s\mathcal{A}}\big]_{0j} \big[e^{-(t-s)\mathcal{A}}\big]_{j0}\, \mathrm{d}s, \qquad \mathcal{A} = 4I - w\,\mathrm{adj}(P_n), \tag{6}$$

with P_n the path on the n data sites. The relative factor i is a gauge-invariant loop phase. With $D_w = \mathrm{diag}(w^{2j})$, $C(w) = D_w^{-1}\Gamma(w)D_w^{-1}$ and $B(v) = D_v C(wv) D_v$, a positive congruence turns the zero-energy problem into $h(k) = -2C(w)\cos k + 2\eta B(v)\sin k + \mu E_u$. Its centres at $\mu = 0$ are $\theta_j = -\arctan(\eta r_j)$, with $r_j = \kappa_j v^{4j}(1 + o(1))$ the generalized eigenvalues of $B(v)$ against $C(w)$. A

weak on-site readout, which counts a data bit only where the biased walker sits, adds a geometric field μE_u, $E_u = \mathrm{diag}(u^{4j})$, which moves the two Fermi points of band j in opposite directions with slopes $s_j \propto u^{4j}$. Centres and slopes are both lacunary, with consecutive ratios below $1/32$. A nonzero signed combination of at most four of them is then dominated by its first nonzero term, and this proves, for every n, that a small field leaves

$$\delta_4 \;\geq\; \min\big(|\theta_{n-1}|/4,\; \mu s_{n-1}/4\big) > 0. \tag{7}$$

The parameters (η, u, v, μ) come from a dyadic search that accepts a tuple only after exact Sturm isolation of all $2n$ roots and rational comparison of all pair sums. Its termination is proved analytically, though no depth bound is given [19, 21, 47].

2.5 Mobile cells

With frozen cells the chain breaks translation b-fold on rings with $b \mid L$ and carries immobile defects on other rings. It is then outside the hypothesis of [1] on most ring lengths, and Latorre and Sierra's blocking remark applies [46]. There are two mobile completions.

Form W (wrapper). A separate selector track lets cells take two lengths, $n+3$ and $n+4$. Typed holes and a length filter keep $n+2$ occupied core sites in both, a helix of compensators makes the port coupling independent of the two lengths, and a marker move moves a hole over a cell boundary while acting as the identity on the payload [16, 17, 22]. The sequence of long and short cells is a hard-core gas, a $c = 1$ "length fluid", so the predicted total is $c = n + 1$. The payload rides on the length fluid as spin rides on charge in the Ogata–Shiba picture [3].

Form V (vacancy). Add one vacancy state $|\varnothing\rangle$ that swaps freely with any letter, and let a weak copy of the frozen bond act only on adjacent occupied sites. In squeezed coordinates the vacancies are a free hard-core gas (one charge boson) and the occupied sites carry the frozen chain. The b translates of a frozen pattern become charge boundary twists, that is, a finer charge-winding lattice, instead of a b-fold degeneracy [50, 20]. This completion was proposed in the campaign of a collaborator (GitHub: salpace) [50], and at $n = 23$ it gives $c = 23 + 1 = 24$ with $d = 24$.

3 The explicit ingredients

3.1 Form W at $d_0 = 4464$

The site space is

$$\mathbb{C}^{4464} = \mathbb{C}^4_{\mathrm{sel}} \otimes \Big[\big(\mathbb{C}^3_{\mathrm{tag}} \otimes \mathbb{C}^2_{\mathrm{filter}} \otimes \mathbb{C}^{46}_{\mathrm{core}}\big) \oplus \mathbb{C}^3_{\mathrm{hole}}\Big] \otimes \mathbb{C}^2_{\mathrm{helix\,A}} \otimes \mathbb{C}^2_{\mathrm{helix\,B}}, \tag{8}$$

so $d_0 = 4 \cdot (3 \cdot 2 \cdot 46 + 3) \cdot 4 = 4464$, and the bond is

$$h_n = h_{\mathrm{sel}} + P_{\mathrm{int}} \otimes h_{\mathrm{wrap}} + h_{\mathrm{helix}} + h_{\mathrm{move}} + \lambda\, h_{\mathrm{port}} + h_{\mathrm{field}}, \tag{9}$$

with every term written out in [52, §2]. The 46 core letters are ten shared data, vacuum and marker letters plus eighteen excited letters for each of the two buses. Every core move enters as a positive edge matrix, and each vacuum edge is replaced by $(q|j\rangle - |j+1\rangle)(q\langle j| - \langle j+1|)$, so the core is positive and frustration free with code

$$G|b\rangle = S^{-1/2} \sum_{\ell=0}^{n+1} q^{\ell}\, |\ell, v; b\rangle, \qquad b \in \{0,1\}^n, \qquad \gamma_{\mathrm{core}} \geq \min\big(K,\, 1-|g|,\, 1+q^2-2q\cos\tfrac{\pi}{n+2}\big). \tag{10}$$

The exact coefficient of λ^2 between two cells is [19]

$$K_2 = -\frac{q^{2n+2}}{S^2} \sum_{r=1,2} b_r^2 \, B_r^\dagger \big(H_r \otimes I + I \otimes H_r\big)^{-1} B_r, \qquad b_1 = 1, \; b_2 = \sqrt{\eta}, \tag{11}$$

with H_r the excited-cell Hamiltonian of bus r and B_r its pair-source map. Its Gaussian part at order g^2 is (5). The literal 2116×2116 core bond is in `codex/work_C39/literal.py`. Certified points at $n = 1, 2, 3$ ($w = 1/4$, $v = 1/1024$, $u = 1/64$, $\eta = 1/100$, $\mu = 10^{-(8n+10)}$) have $\delta_4 > 5.11 \times 10^{-16}$, 1.23×10^{-28}, 2.36×10^{-41}. They transfer to finite D and g with $\log_{10} D \approx 102, 121, 155$ and $g = 2^{-1138}, 2^{-1738}, 2^{-2761}$ [21].

3.2 Form V at $d = 24$

With h_{23} the explicit 23-letter frozen bond (data letters, a token on a data or express rail, five marker letters, a port λV and on-site terms [45]) and $n_{\mathrm{occ}} = 1 - |\varnothing\rangle\langle\varnothing|$, the bond is

$$h_{24} = -t \sum_{a=1}^{23} \big(|\varnothing a\rangle\langle a\varnothing| + \text{h.c.} \big) + J \, (\iota \otimes \iota) \, h_{23} \, (\iota \otimes \iota)^\dagger - \frac{\mu}{2} \big(n_{\mathrm{occ}} \otimes 1 + 1 \otimes n_{\mathrm{occ}} \big). \tag{12}$$

Occupied number, internal charge and marker number are conserved. In squeezed coordinates, and for any internal bond, the vacancy hopping becomes $-t(T_{\mathrm{bulk}} \otimes 1 + B \otimes R + \text{h.c.})$, where R cyclically shifts the internal word when a particle crosses the cut, and the internal coupling becomes $\sum_j A_j \otimes h_{23}(j, j+1)$, with A_j the indicator that occupied sites j and $j+1$ are adjacent. On a charge sea of density ρ the first-order coupling is

$$P_{\mathrm{sea}} \, V \, P_{\mathrm{sea}} = \alpha(\rho) \, H_{23}^{\mathrm{squeezed}}, \qquad \alpha(\rho) = \rho - \frac{\sin^2(\pi\rho)}{\pi^2 \rho}, \qquad \alpha(\tfrac{1}{2}) = \tfrac{1}{2} - \tfrac{2}{\pi^2}. \tag{13}$$

The vacancy problem is the $q = 0$ phase model [12], and for $2/5 < \rho < 3/5$ every uniform vertex carrying a nonzero charge harmonic has dimension above $20/9$, so only the payload's own resonances remain, excluded by its certified margins [50, 20]. The internal point at $n = 23$ (cell length 24, $g = 2^{-365}$ [18]) has an integer interval certificate giving 46 simple Fermi points with $\delta_2 > 0.0140193$ and $\delta_4 > 3.1558 \times 10^{-7}$. The cell-length selector has margin $\gamma_{24} = 7.567 \times 10^{-5}$, and all wrong lengths and illegal words are excluded for $0 < \lambda \le 2^{-20}$ [50].

4 Order of limits and scales

The order is as follows. (i) Fix n. (ii) Choose every coupling, each after the quantities it must beat. For Form W this is the rational witness (η, u, v, μ) with its margins and a length density avoiding the finitely many mixed resonances, then D, then g, then the wrapper ratios, then the marker hopping and readout, and the port coupling λ last [23]. For Form V it is the internal point, then λ, then t, then a density in $(2/5, 3/5)$, then J/t, then μ. None of these is sent to zero and none depends on L. (iii) Take $L \to \infty$ over all ring lengths above a fixed threshold. (iv) Only then compare different n. Nothing is uniform in n.

Two interchanges are illegal. A finite-cluster or finite-ring expansion in λ is not uniform in L, since the retained code band has width growing with L and the length or charge fluid is gapless, so no global Schrieffer–Wolff argument reaches the thermodynamic limit. And the limits that equalize the two cell lengths of Form W are never taken. The mismatch is bounded at fixed small wrapper ratios chosen before λ.

Ingredient	Where	Status
46-letter core, code, K_2, all-n band lemma	`codex/work_C39/` (`PROOFS.md`, `literal.py`, `certify.py`)	proved, certified $n \le 3$, AI referee [47]
Finite D, g budgets for every n	`codex/work_C42/` (`PROOF.md`, `budgets.py`)	proved
Wrapper on the 46-letter core, $d_0 = 4464$	`codex/work_C44/`, built on `codex/work_C8/`, `codex/work_C32/`	proved, AI referee [48]
Rule-table rebuild of C44, calibrated c at $n = 1$	`contrib/verify_C44/`	numerical
Local Schrieffer–Wolff, threshold, schedule	`codex/work_C45/` (`PROOF.md` §3–8)	proved, conditional on S
Massive matching, ensemble transfer, c from a Gaussian limit, no-go, adversarial ledger	`codex/work_C48/`, `work_C58/`, `work_C51/`, `work_C52/`, `work_C54/`	proved, with stated conditions
Lemma N and lemma TP	`codex/work_C55/`, `work_C60/`, `work_C61/`; `work_C57/`, `work_C59/`, `work_C62/`; progress in `work_C64/` to `work_C79/`	open, partial results
23-letter frozen bond, $n = 23$ certificate	`contrib/explicit_candidate/`, `codex/work_C33/`, `codex/work_C37/`	certified, rerun twice
Margins of the 23-letter chain, $n \le 23$	`contrib/margins_audit/`	high-precision numerics
Vacancy lift, squeezing, sector bounds, harmonics	`contrib/counterexample_campaign/`	finite parts proved, AI referee [49]
Assembled argument, both forms	`contrib/routeA_final/README.md`	synthesis
Mechanism checks at small d	`contrib/routeA_demonstrator/`, `contrib/vacancy_mobile_numerics/`, `contrib/mobile_cells/`	numerical
Vacancy count at $n = 2$, S on the microscopic demonstrator	`contrib/vacancy_n2_dmrg/`, `contrib/demonstrator_luttinger/`	numerical
Frozen $d = 11$, mobile $d = 13$	`codex/work_C84/`, `work_C85/`, `work_C88/`, referees in `contrib/referee_C84_C85/`, `contrib/referee_C88/`; companion note `contrib/small_d_note/`	certified, AI referee, infrared assumed
$d = 2$ surveys	`contrib/d2_grid/`, `contrib/d2_staggered/`	numerical

Table 1: Where each ingredient lives in the repository. Paths are relative to `notes/cd_conjecture/`.

The order within step (ii) also matters, and one ordering is destructive. The length fluid carries an attractive density interaction V, and a hard-core gas with hopping t_ℓ phase-separates once $V < -2t_\ell$ (standard, and proved for this fluid in [30]). With the rebuilt bond at small wrapper ratio, $s/\delta \to 0$, V scales as ρ^4 in the wrapper parameter ρ, with $V/|t| = -49, -6.3, -0.32$ at $\rho = 0.16, 0.1, 0.05$ (the scheduled point) in units of the payload hopping (numerical [39]). At C44's finite diagnostic ratios a fast term gives $V/|t|$ between -3×10^4 and -3×10^5 instead, which would phase-separate the length fluid and remove the length boson unless the length hopping were correspondingly large. The schedule takes s/δ small before λ and equalizes the two lengths with g first, which excludes this (proved within the schedule [30]). A reviewer should treat the C44 diagnostic ratios as checks of formulas, not as a point in the phase of Claim A.

The central charge should be read from the conformal towers or from the entanglement of an interval of the infinite chain,

$$S(\ell) = \frac{n+1}{3} \log \ell + O(1), \tag{14}$$

which does not depend on the velocities. The Casimir coefficient is velocity weighted, and at irrational length density the zero-mode sectors drift with L (as for an XXZ chain in a generic field), so $L(E_0 - eL)$ has no limit [48].

The scales are extreme. Carrying out the Form W schedule with the constants of S set to one gives, in $\log_{10}$, $D \approx 114$, $J_0 \approx -1472$, $\lambda \approx -38263$ and a payload energy unit $W \approx -65472$ at $n = 1$, with $W \approx -230507$ and -639244 at $n = 2, 3$. The conformal regime begins beyond about 10^{35}, 10^{130} and 10^{303} cells [48]. For Form V at $n = 23$ the referee estimates J/t below 10^{-510} and payload speeds below $10^{-1000}t$ [49]. Existence at fixed n does not need moderate numbers, since the conjecture concerns each fixed h and no step bounds a small scale from below by a larger one. But nothing at these parameters will ever be checked by simulation.

Two consistency checks follow. At fixed d the normalized bonds $h_n/\|h_n\|$ accumulate, so a limit point must be gapless and non-conformal or have a diverging crossover scale, and here the crossover energy vanishes faster than any power of $\|h_n\|$. Second, a proved conditional bound in the repository, $c \le 1 + (6/\pi)K \log d$ when c conformal levels appear on rings of length at most K, forces any counterexample to have a conformal crossover length of at least $\pi(c-1)/(6 \log d)$ sites. The crossover lengths above grow much faster.

5 What is proved, what is computed, and what is assumed

5.1 Proved

For Claim A the following hold for every n, with proofs in the cited folders. The core is positive and frustration free with the exact 2^n code (10), exact $U(1)$ and fermion parity [19]. The second-order bond (11) has no first-order or three-cell part and its order-g^2 part is the Gaussian chain (5). The lacunary pencil gives $2n$ simple Fermi points, distinct centres and $\delta_2, \delta_4 > 0$, and explicit D and g preserve all of this together with a bound on the complete non-Gaussian remainder [19, 21]. The wrapper keeps positivity, the code, the gap and the $U(1)$, its occupation filter and selector secant bound exclude every illegal, marker-free and wrong-length sector, and on every ring with $L \ge (n+3)(n+4)$ the unperturbed ground space is the set of selected tilings with their codes [22]. The two lengths' bonds agree up to an explicit error that can be made smaller than any prescribed fraction of J_0 [22].

The infrared reduction [23] writes the chain in ordered cell coordinates as $H = A + \lambda V + \lambda^2 F$ with $A_j \ge \Delta(1 - P_j)$ and $\dim P_j = 2^{n+1}$, and builds a range-three generator $S_{\mathrm{loc}} = \lambda S_1 + \lambda^2 S_2$ such that

$$e^{S_{\mathrm{loc}}} H e^{-S_{\mathrm{loc}}} = D + R, \qquad PDP = \lambda^2 \big[PFP - PVQA^{-1}QVP \big], \qquad \|R\|_a \le C_{\mathrm{SW}} \lambda^3, \tag{15}$$

with D locally block diagonal, $C_{\mathrm{SW}} = 2^{16}(v_0 G/\Delta + v_0^3/\Delta^2)$, $G = f_0 + 48 v_0^2/\Delta$, valid for $\lambda \le \min(v_0/G, \Delta/(192 v_0))$, and the norm $\|\cdot\|_a$ independent of the number of cells. A completely positive extension argument gives the ground-relative threshold

$$D \;\ge\; E_P\, I + \frac{\Delta}{2} \sum_j (1 - P_j), \tag{16}$$

which allows moving defects and does not require a gap inside the extensive code band. The schedule then ends with

$$\lambda \;\le\; \frac{\epsilon_n\, J_0/s}{16\, A_{\mathrm{dec},n}\, C_{\mathrm{SW}}}, \qquad W = \lambda^2 J_0/s, \tag{17}$$

where ϵ_n is the tolerance built from the margins and $A_{\mathrm{dec},n}$ is a constant of S. Results on S itself, starting with the partial results of [24], are collected in Section 6.

For Claim B the proved parts are the bond and its conservation laws, the squeezing identity and (13), global length selection and illegal-word exclusion at fixed nonzero λ, the free-charge momentum threshold at half filling, and the harmonic selection rule within the bosonization dictionary [50, 20].

5.2 Computed

Exact rational certificates exist for the band lemma at $n = 1, 2, 3$, and a referee reproduced the margins at the certified tuple through $n = 8$ in 500 to 1300 digits [47]. The literal core bond and the complete second-order bond at $n \leq 3$ agree with the formulas, and an independent reimplementation reproduces the bond to 7×10^{-17} [47]. For the wrapper, the $n = 1$ legal cell (dimensions 1728 and 4320) has the two-dimensional code at both lengths, and projected marker-graph spectra match the joint length–payload momentum formula to 2.4×10^{-14}. The referee of C44 and C45 rebuilt the charge-gap bounds, the selector bounds for b up to 4470, the equalization inequality, the local Schrieffer–Wolff identity (to 7×10^{-16}, with $\|R\|/\lambda^3$ converging and linear in the number of cells) and the Gaussian metric bound, from the prose and without importing the authors' code [48].

That referee asked for two further checks, now done by a separate Claude agent that imported nothing from C32, C39, C44 or C45 [39]. First, the core and wrapper were rebuilt from the rule text alone. The bare bond agrees with C44 to 8×10^{-17}, the second-order wrapped bond (C44 Eq. 11) to 5×10^{-11} at $n = 1$ and 3.4×10^{-9} at $n = 2$, the first-order port compression vanishes identically, and the marker-move overlap (C44 Eq. 9) is $A_b^2 \beta_v \cos(1/2b)^{2b}$ times the identity to 10^{-16}, all at $n = 1, 2, 3$. C44's traceless length–payload numbers at $n = 2$ turn out to be below the roundoff of its own eigensolves. Second, the projected $n = 1$ chain (a length bit and a payload qubit per cell, the rebuilt bond at $s/\delta \to 0$) at the scheduled point $\rho = 0.05$, calibrated against an exact $c = 2$ reference, gives

$$c_{\mathrm{FES}} = 1.97 \text{ (2.00 two-scale)}, \qquad c_{\mathrm{interval}} = 2.03 \ (L = 32),\ 2.00 \ (L = 48), \tag{18}$$

from finite-entanglement scaling and open-chain interval entanglement. Couplings multiplied by 300 still give 1.98 to 2.03, and a locked control gives $c = 1$. These numerical results concern the projected model, so they test part (b) of S at $n = 1$, not part (a), the 4464-state chain or large n.

For Claim B, the 576-by-576 bond was rebuilt independently entry by entry, the $n = 23$ certificate was regenerated bit for bit twice, and a margins audit finds positive δ_4 at every $n \leq 23$ in 115- to 313-digit arithmetic [49, 53].

The mechanism has been seen at small d. A ten-letter frozen bond forms cells unaided and its microscopic chain gives $c = 1.00$ to 1.02 by finite-entanglement scaling at three couplings, with ring spectra matching the effective logical chain to better than 6×10^{-3} relative and $z = 1$ [54, 15]. Calibration chains at the same correlation lengths read 0.90 to 0.92, so that estimate carries a systematic uncertainty near 0.1. A $d = 6$ mobile toy has a unique ground state on rings $L = 8$ to 18 and $c = 2$ from three estimators [55]. The vacancy lift of the ten-letter bond ($d = 11$) has a unique ground state on rings $L = 5$ to 11 up to a payload doublet, and a calibrated charge boson with $c = 0.998$, but its payload is 360 times slower and its half of $c = 2$ is not resolved [56]. No chain with $c > d - 1$ has been simulated. None of the small-d checks in this paragraph uses the two-bus core, and the projected $n = 1$ chain of (18) is the only computation of c that does.

The vacancy count [numerical]. The payloads of the real lifts are far too slow to resolve. A logical vacancy model keeps the mechanism, each site being a vacancy or a cell of n Jordan–Wigner flavours that hops through vacancies by the same lift, with the small non-Gaussian part dropped. At $n = 1$ two calibrated estimators give 1.92 to 2.04, against 1.05 for a gapped-payload control [56]. At $n = 2$, a DMRG conserving cell and flavour numbers, with interval entanglement calibrated

against the exact three-free-chain reference, gives 2.86 to 2.91 on open chains of $L = 40$ to 100 and 2.8 to 3.2 by finite-entanglement scaling, while the control with one flavour gapped gives 1.85 to 2.0 [58]. So

$$c(n = 1) = 2.0, \qquad c(n = 2) = 2.9 \pm 0.1 \quad \text{against a } c = 2 \text{ control}, \tag{19}$$

and c rises as $n + 1$. The step from the logical model back to the microscopic chain is first-order spin–charge separation [heuristic].

Input S on a full microscopic chain [numerical]. In the ten-letter demonstrator with the collaborator's coupling tuning, the second-order logical chain is XXZ at anisotropy $g = 27/182$, $27/364$ or $27/728$, and the exact two-cell bond shifts it to $g_x(\lambda)$. DMRG on the unchanged microscopic chain (up to 48 cells, $0.0125 \le \lambda \le 0.15$) gives $c = 0.98$ to 1.01, equal to the logical chain within 0.002, Bethe-ansatz velocities within 1.5 percent, and

$$K_{\text{micro}}(\lambda) = K_{\text{XXZ}}\big(g_x(\lambda)\big) + O(\lambda^2) = K_{\text{XXZ}}(g) + \lambda \, \frac{\mathrm{d}K_{\text{XXZ}}}{\mathrm{d}g} \, \frac{\mathrm{d}g_x}{\mathrm{d}\lambda} + O(\lambda^2), \tag{20}$$

with fitted slopes 0.16, 0.35, 0.74 against predicted 0.18, 0.35, 0.72, and agreement with $K_{\text{XXZ}}(g_x(\lambda))$ to 2×10^{-4} for $\lambda \le 0.05$ [59]. This is S in a family where the answer is known: the full chain flows to the Luttinger liquid of its matched logical chain, with the $O(\lambda)$ correction a local shift of the anisotropy. It has one channel and no velocity hierarchy.

5.3 The load-bearing assumption

Everything above is finite-dimensional or a fixed-order local statement. The passage to an infrared CFT rests on one input, stated in [23, §5] and restated by the referee [48].

Assumption S. Let $D + R$ be as in (15), with D locally block diagonal and the threshold (16). Suppose the retained block is W times a number-conserving quadratic reference with finitely many simple Fermi points and positive margins (4), plus a relative correction ϵ in a weighted local coefficient norm, and R is symmetric, quasi-local and $\|R\|_a \le \epsilon W$. Suppose also that the local operator lattice is that of the compact bosons with identity transport, with no extra fractional vertex. Then (a) integrating out the auxiliary sector, with the retained fields left unintegrated, changes the connected low-frequency retained vertices by

$$\|\delta S_R\|_{\text{LE}} \le A_{\text{dec}} \|R\|_a, \qquad \|\delta S_R(\omega) - \delta S_R(0)\|_{\text{LE}} \le A_{\text{dec}} \|R\|_a \, |\omega|/\Delta, \tag{21}$$

with A_{dec} finite and uniform as $W/\Delta \to 0$, and (b) the resulting multi-component compact-boson theory obeys weak-coupling RG, with a marginal majorant $x(\ell) \le X_0 + A_\beta \int_0^\ell x^2$ and irrelevant couplings decaying as a positive power of the energy, set by $\sigma = 3\sqrt{3/5} - 2$. All constants depend only on the normalized reference, not on Δ, the core scale or the wrapper ratios.

Under S, choosing ϵ_n so that $A_\beta X_0 \, \ell_* < 1/2$ over the whole interval $\ell_* = \log(e_{\text{top}}/e_{\text{det}})$ between the top bandwidth and the lowest detuning scale gives n payload bosons and one length boson. The last clause of S matters. If the constants depended on Δ or on the wrapper ratios, the choice of ϵ_n in (17) would be circular [48].

In physics terms, S is how the Heisenberg chain is obtained from the large-U Hubbard chain. One writes $(4t^2/U) \sum (\mathbf{S}_i \cdot \mathbf{S}_j - \frac{1}{4})$, treats higher orders as small local symmetric corrections, and invokes the stability of $SU(2)_1$ [3]. Part (a) is the first step and part (b) the second. The application here is unusual in three ways. The protection comes from momentum margins, not a symmetry

($\delta_4 \sim 10^{-16}, 10^{-28}, 10^{-41}$ at $n = 1, 2, 3$). Many small parameters are chosen in sequence. And the velocities span many decades, with n separate crossover scales.

Literature status. We know of no theorem that covers S [62]. Hamiltonian simulation needs couplings that grow with system size [7, 8], local Schrieffer–Wolff and finite-order effective Hamiltonians give only an extensive spectral error [4, 5], and prethermal bounds hold for finite times [6]. Constructive RG controls correlations of multi-channel Luttinger liquids with arbitrary velocities and pairwise non-resonance [9, 10, 11], which is close to part (b), but it has not been applied to a model obtained by integrating out a massive sector, and it does not compute c. On the physics side the step is confirmed in large-U Hubbard and related models with up to four flavours and two velocity scales. The documented failures involve marginal flows or symmetry-allowed relevant operators missed at low order [13, 14], which the margins are designed to exclude. Since the raw Grassmann expansion of the eliminated action can diverge [24], a rigorous route needs a resummed retarded action, which C48 has since supplied (Section 6).

5.4 Additional inputs for Form V

Claim B needs S for the frozen 23-letter chain after blocking 24-site cells (input P), and two further statements [49, 20, 50]. M (matching) says that, with the charge field retained, the remaining induced correction obeys $\|R\|_{\mathrm{LE}}/J \to 0$ as $J/t \to 0$, for example $\|R\|_{\mathrm{LE}} \le C(J^2/t)(1 + |\log J/t|^p)$. The dressed cost of a wrong-length cell must stay positive at finite J. Its bare cost is of order J and the dressing per defect is argued to be $O(J^2/t)$ from a finite local susceptibility, with tagged-defect numerics to $L = 20$ and 264 exact diagonalizations without an undercut. The collaborator's own adversarial audit [51] withdrew an earlier "verified" verdict because a uniform finite-J dressed-defect comparison is not established. Uniqueness of the ground state on every ring is also open for these complex bonds, though [1] does not require it.

6 Status of input S

After the reduction to S, sixteen gpt-6-astra runs (C47 to C62) attacked it in four rounds, and the coordinating agent reran every check script. No human has read these proofs. The upshot is that part (a) of S and the passage to the physical ring are proved at fixed n, while part (b) is reduced to two estimates about the constructive RG of the logical fermion action. Later runs (C64 to C79) made progress on both, summarized at the end of Section 6.2.

6.1 What is now proved

Resummed massive matching (C48) [proved, given the finite inputs of C42, C44, C45]. Each cell's whole excited Hamiltonian is kept as an auxiliary operator and only its finite logical blocks are expanded. A coarse-time polymer expansion then gives exponentially decaying retarded kernels of all degrees, uniformly in the number of cells and for $\beta \ge 2T_n/\Delta$ (including $\beta \to \infty$), and the induced change of the retained action is linear in the C45 remainder,

$$\|\delta S_R\|_{\mathrm{LE}} \le A_{\mathrm{dec},n}\, C_{\mathrm{SW}}\, \lambda^3, \qquad A_{\mathrm{dec},n} = 2^{22} e^{\mathcal{L}} \big[2 + 16(16T_n/\mathcal{L})^2\big], \tag{22}$$

with $\mathcal{L}$, T_n explicit in n and the output spatial exponent, plus a matching bound on the frequency derivative [25, PROOF Eqs. 20, 26, 29]. At fixed n the constant does not depend on Δ, the wrapper ratios or the logical bandwidth, which is the uniformity clause of S, so a fixed positive λ makes the relative error small independently of the ring size. With no expansion in u/Δ, this removes the

obstruction of [24]. The operator-valued route was first proposed in the collaborator's matching notes [40].

Physical-ensemble transfer (C58) [proved, given C48 and the finite construction]. For the original wrapped chain, at fixed n and small fixed couplings, the rooted pressure and the pressure of the exact matched thermal logical action obey

$$\left|p_{\mathrm{orig}}(T,s) - p_{\mathrm{matched}}(T,s)\right| \le C\,T\,e^{-\delta/T}, \qquad \delta > 0, \tag{23}$$

uniformly on an interval of the length source s, with $L \to \infty$ before $T \to 0$. The same holds at fixed physical length, both with marker number summed and in canonical sectors at an interior density, with the cyclic identification and periodic boundary condition included. Illegal cells are handled by a positive completion with an invariant mismatch flag (building on the collaborator's completion of C44's helix pins [41]), fixed-marker sectors by a new isometry [34, PROOF §§3, 8, 9]. Conditional on the two estimates of Section 6.2, the physical chain then has

$$f(T) = e_{\mathrm{phys}} - \frac{\pi}{12}\sum_{a=1}^{n+1}\Big(\frac{1}{u_{a,R}} + \frac{1}{u_{a,L}}\Big)T^2 + o(T^2) \tag{24}$$

with dressed physical speeds. No $o(1/L)$ spectral theorem is obtained. The free energy is the route.

c from a complete Gaussian limit (C51) [proved, given a complete Gaussian scaling limit]. For $n+1$ compact bosons with any positive quadratic Hamiltonian, including density–current mixing and arbitrarily separated velocities, $c_L = c_R = n+1$, and the drifting zero-mode sectors at irrational density do not change this. A complete limit means normalized chiral current algebras, the compact charged sectors and exhaustion of the infrared theory by these fields, which power laws of selected correlations do not establish. Since (24) weighs inverse speeds, it fixes c only with the dressed speeds [27].

Exact decoupling is impossible (C52) [proved, no-go]. If a nearest-neighbour chain has an exactly invariant product of cell codes with distinct endpoints, adjacent logical bonds commute, and the three-cell one-particle matrix element of their commutator is T^2. So the logical hopping cannot be invertible, while C39's satisfies $\sigma_{\min}(T_0) \ge W\lambda_{\min}(\Gamma(w)) > 0$ for every n. S cannot be avoided by such a redesign and has to be established perturbatively [28].

In-family demonstrator inside Benfatto–Mastropietro (C53) [proved, imports [42]]. The nearest-neighbour ten-letter demonstrator has a second-order logical bond equal to XXZ with anisotropy 27/182, which does not shrink with λ, at half filling, so no published theorem applies [29]. With four-site interactions at the same $d = 10$, C53 makes the frozen logical chain exactly XXZ with tunable anisotropy and field, every non-code sector at least 2/5 higher, and free energies equal up to $T\log[1 + 15e^{-2/(5T)}]$ uniformly in volume. Theorem 1.1 of [42] then applies off half filling at small anisotropy [29]. This checks cell selection and the frozen sector rigorously, not the port mechanism or the interacting T^2 coefficient.

Adversarial ledger (C54) [proved within the operator lattice of S]. No mechanism breaking S inside the schedule was found. The margins detune every non-Gaussian operator of dimension at most two (the list is exhaustive), and all others have dimension at least $3\sqrt{3/5}$. An amplitude audit shows that the conversion and length interactions are suppressed relative to W by g and the wrapper equalization, not by λ. Weaker substitutes for S (an $O(\lambda^3)$ pressure error, a prethermal remainder, an extensive eigenvalue error) provably cannot fix c. The one destructive mechanism found is the phase separation of Section 4, outside the schedule [30].

Partial results [proved, partial]. These include a quasi-local dressing valid on an exponentially large but finite window of sizes and finite-scale integration uniform in the velocity hierarchy [26],

all-scale control of the part of the flow linear in the initial action [31], a conditional multichannel counterterm map [32], and the first-order thermal coefficient $T^2/(12t^2)$ of interacting spinless fermions [33].

6.2 What remains

Both remaining estimates concern the logical fermion action, not the microscopic chain, and each is stated for single-channel XXZ. The construction needs them for the matched retarded action with $2n+2$ chiral speeds. C56 finds that the multichannel comparison needs the same tree estimate as Lemma N, and C57 that its thermal bootstrap extends to finitely many channels in matrix norms, with constants depending on the minimal speed and the detunings. At the physics level both are the expected behaviour of a weakly interacting non-resonant Luttinger liquid [heuristic]. As theorems they are open [assumed].

Lemma N, corrected form [31, PROOF §9], [36], [37, PROOF §2]. It is a normalized renormalized-tree majorant for arbitrary-degree retarded initial endpoints, with no cumulative loss of field radius or kernel moments, a summable short-memory gain on trees reaching an initial irrelevant endpoint, and source marks that preserve the estimates. C60 proves the degree hierarchy $\|K_{2m}\| \le C^{2m}\epsilon^{\max(1,m-1)}$ for the matched action, compatible with the scale-zero kernels of [42], and finds a free counterexample to the original source budget. At zero interaction, for $P_x = f_x f_{x+1}$ at $p_F = \pi/3$ and $x = 3m$,

$$\langle P_x P_0^\dagger \rangle - \frac{\sin^2 p_F}{\pi^2 x^2} = \frac{3}{4\pi^2 x^2 (x^2-1)}, \tag{25}$$

which no remainder proportional to the interaction can absorb. C61 therefore splits the source difference as

$$D_O(u) = D_O(0) + \big[D_O(u) - D_O(0)\big], \tag{26}$$

with only the bracket carrying an interaction prefactor, and reduces C60's scale-allocation inequality to a cut-coverage condition in the reduced scale tree. C79 has since proved that inequality for general trees (below). Given Lemma N, the scaling dimensions K and $1/K$ transfer [31, PROOF §11].

Lemma TP [35, PROOF §7], [38, PROOF §7]. Let X_h^T, Y_h^T be the effective data of the lattice and of a Luttinger reference with the dressed v, K at scale q_h and temperature T. TP asks for

$$\|X_h^0 - Y_h^0\| \le C\, q_h^\theta, \qquad \big\|(X_h^T - Y_h^T) - (X_h^0 - Y_h^0)\big\| \le C\, q_h^\theta\, (T/q_h)^p, \tag{27}$$

with $\theta > 0$, $p > 2$ and compatible vacuum derivative and Hessian bounds, uniformly in depth and regulators. Under TP, $f(T) = e_0 - (\pi/6v)T^2 + o(T^2)$ for XXZ at fixed nonzero anisotropy [35]. C62 builds a trace-defined, smoothed reference whose scalar is the oscillator trace, with coefficient $-\pi/(6v)$, and shows that $p = 3$ suffices given a joint marked-tree majorant with one short-memory factor. That majorant is the missing step. No counterexample to the thermal formula was found. The analogous bound is proved for non-solvable Ising models ($c = 1/2$) [43], and a 2024 talk announces the thermal correction for weakly interacting chains, with no manuscript located [44].

Progress on Lemma N (C64, C70, C72, C74, C79) [proved, partial]. After partial steps (C64 to C74, including all quartic pair ladders), C79 treats general trees. A moment born at scale u and routed to an edge at scale $t \ge u$ carries an extra $2^{-(t-u)}$, so an edge with load b has a surplus

$$q(b) = \big\lfloor (b-1)^2/4 \big\rfloor, \qquad b \ge 1, \tag{28}$$

half of which pays $(b!)^2$ and half the routing choices, for a total $C^{\sum_i p_i+n}$ independent of depth, with several quartic children per vertex and retarded endpoints of any degree. This proves C60's scale-allocation inequality, the source-free localization estimate [57]. The comparison and source parts of Lemma N remain open [assumed].

Progress on lemma TP (C65, C71, C73, C78) [proved, partial]. With C62's sharp windows the thermal density bubble has a $|p|$ cusp at every finite cutoff, so TP cannot hold uniformly in that regulator, and C65 supplies a smooth one with the same Gibbs limit. C71 shows that a majorant-only route fails, so cancellations must be kept before norms. C73 integrates one chirality to all orders on a cutoff-independent disk, with $\|\partial_A^j(F^T - \mathrm{Per}_\beta F^0)\| \le C|g|^2 T^P$, $P > 1$, and C78 does the same for the bubble-ring subseries of both chiralities, finding

$$f^{(2)}(T) - f^{(2)}(0) = -\frac{g^2 T^2}{48\pi A^3} + o(T^2), \tag{29}$$

consistent with the sound dispersion of C62's reference. The complete two-chirality integration and the infrared comparison remain open [assumed] [57].

In sum, Lemma N with its multichannel comparison would give the complete Gaussian limit with dressed speeds, TP the thermal coefficient, C58 would carry it to the physical ring as (24), and C51 would identify $c = n + 1$. The collaborator's ledger lists the same two obligations [40].

7 Smaller local dimension

A second line of runs (C75 to C95) asked how small d can be made with the same mechanism. Full details are in a companion note being written in parallel, `contrib/small_d_note/`. The two smallest candidates, conditional on the inputs below, are

$$d = 11: \quad c_T = 11 > d - 1 = 10 \ \text{(frozen)}, \qquad d = 13: \quad c_T = 14 > d - 1 = 12 \ \text{(mobile)}. \tag{30}$$

Frozen, $d = 11$ (C84) [proved finite construction, certified, AI referee]. The eleven letters are $L_{0,1}$, $R_{0,1}$, $D_{A,B}$, $X_{A,B}$, E, $M_{A,B}$ (bond in `codex/work_C84/bond11.py`). Tokens carry no bit, the displaced bit is shifted to the next site, and a vacuum domain wall replaces the vacuum token. A certified member at $n = 11$ (cell length 13) has 22 simple Fermi points and pair-sum separation above 9.12×10^{-7} over all 253 pair sums, with all finite-coupling errors in budget. As a frozen chain it carries the caveat of Section 2.5.

Mobile, $d = 13$ (C88) [proved finite construction, certified, AI referee]. Two vacancy colours V_0, V_1 carry the grammar state of the preceding material letter, so every illegal squeezed pair forces a local mismatch across any number of vacancies. The boundary pin of the eleven-letter grammar is a superselection label, which removes the third colour of an earlier $d = 14$ lift (C85), and in the pin-free sector C85's certified $n = 13$ member applies unchanged (26 simple Fermi points, pair-sum separation above 3.25×10^{-6}). The predicted infrared is 13 payload bosons plus the vacancy charge boson. Claude referees rebuilt both bonds, reproduced both certificates bit for bit and found no fatal or serious new gap [60].

Inputs [assumed]. Both inherit the infrared inputs of the larger chains, S for the frozen chain and, at $d = 13$, vacancy matching and a positive dressed defect cost. C90 transfers C48, C58 and C51 to them under stated hypotheses [proved]. Open are the legal charge comparison that the $d = 24$ chain also needs (C94 refutes a sufficient finite-volume form, C95 proves parts of the thermodynamic one) and the finite-J mobile representation for C48 (C91, C93).

Lower bounds. Eleven letters are minimal in the single-token product class (C86, C87) [proved within stated classes], and simpler mechanisms at $3 \le d \le 12$ give at best $c = d - 1$ (C76) [proved or numerical, case by case].

The case $d = 2$ [numerical]. A calibrated survey sampled about 103,000 points across the four residual symmetry strata of the qubit chain and decided all 173 flagged points by finite-entanglement scaling calibrated on Ising, XX, cluster-Ising (1.500) and two XX chains (2.01). None has $c > 1$. A charge-resolved DMRG on the staggered branch, open chains to $L = 256$ at 22 points, finds a single boson tower in every charge sector, no coexisting Ising sector, and at most 1.05 ± 0.05 [61]. A value slightly above 1 could escape the screens.

8 What a reviewer should check, in order

1. Scope. Does an infrared theory of $n + 1$ compact bosons with different velocities, unbroken translation and L-dependent zero-mode sectors count as "a collection of unitary CFTs" in (2)? Latorre and Sierra allow different velocities [1], and the XXZ chain in a generic field has the same zero-mode drift. If the answer is no, the construction does not bear on the conjecture as stated and nothing below matters.

2. Assumption S (Section 5.3). Is it true here, with margins near 10^{-41} at $n = 3$, many sequential small parameters and velocities spread over many decades? Is there a mechanism, such as a relevant operator generated at high order, a marginal flow, or an orthogonality effect from the extensive code band, that defeats it? Are its constants independent of Δ and the wrapper ratios? This is the decisive question. For part (a) a reviewer can now check C48 and C58 (Section 6), and for part (b) the two open estimates, Lemma N in the corrected form of C61 and lemma TP, together with their multichannel extensions.

3. The band lemma for the 46-letter core [19, PROOFS §2–5] and its finite-parameter transfer [21, PROOF]. The lacunarity argument and the pair-sum classification can be checked by hand. `certify.py` reruns the exact certificates in seconds.

4. The local reduction [23, PROOF §3–4], Eqs. (15) and (16), which are short, and the schedule [23, PROOF §8] for circularity.

5. The wrapper [22, PROOFS]. The exclusion bounds were rebuilt by the referee. The transport overlap and the source formula (C44 Eqs. 9 and 11) have now been rebuilt from the rule text and agree at $n = 1, 2, 3$ [39], but C44's traceless length–payload numbers at $n = 2$ are roundoff and should be disregarded.

6. For Claim B, the squeezing identity, the sector bounds, the $n = 23$ certificate, the harmonic selection rule, and inputs M and the dressed defect cost [50, 51].

7. For the smaller candidates, the finite constructions C84 and C88 and the legal charge comparison (Section 7 and `contrib/small_d_note/`).

The scripts that reproduce every finite claim are listed in `contrib/expert_note/REPRODUCE.md`. The two AI referees asked, before acceptance, for a calibrated $c = n + 1$ from two estimators on a small member of the same family at couplings where the payload is resolved, for the rebuild in item 5, and for a human reading of C39 to C45 [48, 49]. The first two are now done at the level stated in Section 5.2, the first only for the projected $n = 1$ chain and for the logical vacancy model at $n = 1, 2$. The human reading is still outstanding, and it now extends to C48, C51, C52 and C58.

9 Provenance and contributions

This note and the work it summarizes were produced with AI assistance, and none of it has been refereed by a human. J. Kudler-Flam directed the project, set the working standard, and coordinated the work, with Claude Opus 5.5 (Anthropic) as coordinating agent, Claude agents (Opus 5.5, and Sonnet 5.5 for the literature check) as authors of the referee, red-team, literature and numerical folders, and closed-book runs of gpt-6-astra (OpenAI, through the Codex CLI) for most derivations. The runs are labelled C1 to C97, and C32, C33, C37, C39, C41, C42, C44 to C62 and C64 to C95 are the ones cited here. The rebuild of C44 and the $n = 1$ central charge [39], the vacancy and demonstrator numerics [56, 58, 59] and the $d = 2$ surveys [61] are the work of separate Claude Opus 5.5 agents, partly run on the IAS Typhon cluster. The tuning of the demonstrator used in [59] is the collaborator's. A collaborator (GitHub: salpace) ran a parallel campaign with GPT-6-family Codex workers. It proposed the one-vacancy completion, wrote the $d = 24$ synthesis and its sector bounds, regenerated the $n = 23$ certificate, ran an adversarial audit that revised its own $d = 24$ verdict from verified to conditional, proposed the matching route that C48 completes, and keeps a ledger of the obligations on S [50, 51, 40, 41]. The 23-letter internal bond and its certificate came from the coordinating side. The referees and the red team were separate agents, but all were AI models, and the C39 to C45 chain comes from one model family. This note was drafted by Claude Opus 5.5 from the repository on 28 October 2026 and revised the same day to include C47 to C62 and the rebuild of C44, and again on 29 October 2026 to include C64 to C95, the smaller candidates and the numerics of REPORT §§9.34 to 9.58.

References

[1] J. I. Latorre and G. Sierra, "The c–d conjecture," J. Stat. Mech. (2024) 113103, arXiv:2403.17242.

[2] B. Sutherland, "Model for a multicomponent quantum system," Phys. Rev. B 12 (1975) 3795. G. V. Uimin, JETP Lett. 12 (1970) 225. C. K. Lai, J. Math. Phys. 15 (1974) 1675.

[3] M. Ogata and H. Shiba, "Bethe-ansatz wave function, momentum distribution, and spin correlation in the one-dimensional strongly correlated Hubbard model," Phys. Rev. B 41 (1990) 2326.

[4] S. Bravyi, D. P. DiVincenzo and D. Loss, "Schrieffer–Wolff transformation for quantum many-body systems," Ann. Phys. 326 (2011) 2793, arXiv:1105.0675.

[5] N. Datta, R. Fernández and J. Fröhlich, "Effective Hamiltonians and phase diagrams for tight-binding models," J. Stat. Phys. 96 (1999) 545, arXiv:math-ph/9809007.

[6] C. Yin and A. Lucas, "Prethermalization and the local robustness of gapped systems," arXiv:2209.11242.

[7] T. Cubitt, A. Montanaro and S. Piddock, "Universal quantum Hamiltonians," Proc. Natl. Acad. Sci. 115 (2018) 9497, arXiv:1701.05182.

[8] T. Kohler, S. Piddock, J. Bausch and T. Cubitt, "Translationally-invariant universal quantum Hamiltonians in 1D," arXiv:2003.13753.

[9] G. Benfatto, P. Falco and V. Mastropietro, "Universality of one-dimensional Fermi systems, I. Response functions and critical exponents," Commun. Math. Phys. 330 (2014) 153, arXiv:1303.3681.

[10] V. Mastropietro and M. Porta, "Multi-channel Luttinger liquids at the edge of quantum Hall systems," Commun. Math. Phys. 395 (2022), arXiv:2108.13708.

[11] M. Porta, G. Scola and H. P. Singh, "Large scale dynamical response of interacting $1d$ Fermi systems," arXiv:2509.08665.

[12] M. Bortz and S. Sergeev, "The q-deformed Bose gas: integrability and thermodynamics," arXiv:cond-mat/0603093.

[13] C. Itoi and M.-H. Kato, "An extended massless phase and the Haldane phase in a spin-1 isotropic antiferromagnetic chain," Phys. Rev. B 55 (1997) 8295, arXiv:cond-mat/9605105.

[14] M. Oshikawa and I. Affleck, "Theory of the field-induced gap in $S = 1/2$ antiferromagnetic chains," Phys. Rev. Lett. 79 (1997) 2883, arXiv:cond-mat/9706085.

[15] F. Pollmann, S. Mukerjee, A. Turner and J. E. Moore, "Theory of finite-entanglement scaling at one-dimensional quantum critical points," Phys. Rev. Lett. 102 (2009) 255701, arXiv:0812.2903.

[16] gpt-6-astra run C8, mobile selector, arrow wrapper, length filter and helix equalizer, `codex/work_C8/` (unrefereed).

[17] gpt-6-astra run C32, the original mobile core at $d_0 = 2736$, `codex/work_C32/` (unrefereed).

[18] gpt-6-astra run C33, finite-g interval certificate of the 23-letter chain at $n = 23$, `codex/work_C33/`.

[19] gpt-6-astra run C39, two folded buses and a geometric readout, `codex/work_C39/`.
[20] gpt-6-astra run C41, vacancy completion of the 23-letter core, `codex/work_C41/`.
[21] gpt-6-astra run C42, all-n schedule with explicit finite-D and finite-g budgets, `codex/work_C42/`.
[22] gpt-6-astra run C44, audit of the mobile wrapper on the 46-letter core, `codex/work_C44/`.
[23] gpt-6-astra run C45, local reduction and the infrared budget, `codex/work_C45/`.
[24] gpt-6-astra run C46, attempted rigorous infrared step, `codex/work_C46/`.
[25] gpt-6-astra run C48, resummed cluster expansion for the massive sector, `codex/work_C48/`.
[26] gpt-6-astra runs C49 and C50, quasi-local dressing and constructive multichannel RG, `codex/work_C49/`, `codex/work_C50/`.
[27] gpt-6-astra run C51, from a Gaussian scaling limit to c, `codex/work_C51/`.
[28] gpt-6-astra run C52, exact-decoupling no-go, `codex/work_C52/`.
[29] gpt-6-astra runs C47 and C53, the demonstrator as XXZ and inside the Benfatto–Mastropietro theorem, `codex/work_C47/`, `codex/work_C53/`.
[30] gpt-6-astra run C54, adversarial search and amplitude ledger for S, `codex/work_C54/`.
[31] gpt-6-astra run C55, single-channel retarded RG and Lemma N, `codex/work_C55/`.
[32] gpt-6-astra run C56, multichannel comparison and counterterm map, `codex/work_C56/`.
[33] gpt-6-astra run C57, the T^2 coefficient and a source audit, `codex/work_C57/`.
[34] gpt-6-astra run C58, transfer to the physical ring, `codex/work_C58/`.
[35] gpt-6-astra run C59, estimate P for XXZ and lemma TP, `codex/work_C59/`.
[36] gpt-6-astra run C60, assembling Lemma N and the free pair-source counterexample, `codex/work_C60/`.
[37] gpt-6-astra run C61, corrected Lemma N and the cut-coverage reduction, `codex/work_C61/`.
[38] gpt-6-astra run C62, a trace-defined reference for lemma TP, `codex/work_C62/`.
[39] Rule-table rebuild of C44 and the $n = 1$ central charge, `contrib/verify_C44/README.md`.
[40] A collaborator (GitHub: salpace) and GPT-6-family Codex workers, operator-valued matching notes and obligations ledger, `contrib/direct_attack_20261028/infrared/` and `contrib/direct_attack_20261028/CURRENT_OBLIGATIONS.md`.
[41] A collaborator (GitHub: salpace) and GPT-6-family Codex workers, thermal-ensemble completion of the wrapper, `contrib/thermal_ensemble_20261028/README.md`.
[42] G. Benfatto and V. Mastropietro, "Universality relations in non-solvable quantum spin chains," J. Stat. Phys. 138 (2010) 1084, arXiv:0907.2837.
[43] A. Giuliani and V. Mastropietro, "Universal finite size corrections and the central charge in non-solvable Ising models," Commun. Math. Phys. 324 (2013) 179, arXiv:1211.3030.
[44] B. Renzi, with A. Giuliani, V. Mastropietro and V. A. Santini, short talk B12, 127th Statistical Mechanics Conference, Rutgers University, 17 December 2024 (abstract only).
[45] The explicit 23-letter bond, `contrib/explicit_candidate/README.md`.
[46] Red-team report on the 23-letter chain, `contrib/redteam_routeA_cells/README.md`.
[47] Referee report on C39, `contrib/referee_C39/README.md`.
[48] Referee report on C44 and C45, `contrib/referee_C44_C45/README.md`.
[49] Referee report on the $d = 24$ chain, `contrib/referee_d24/README.md`.
[50] A collaborator (GitHub: salpace) and GPT-6-family Codex workers, the $d = 24$ campaign, `contrib/counterexample_campaign/` (synthesis in `COUNTEREXAMPLE_D24.md`).
[51] A collaborator (GitHub: salpace) and GPT-6-family Codex workers, adversarial audit and matching attack, `contrib/counterexample_audit_20261028/` and `contrib/direct_attack_20261028/`.
[52] Assembled argument for both mobile forms, `contrib/routeA_final/README.md`.
[53] Margins audit of the 23-letter chain, `contrib/margins_audit/README.md`.
[54] Ten-letter demonstrator, `contrib/routeA_demonstrator/README.md`.
[55] Mobile-cell toy model at $d = 6$, `contrib/mobile_cells/README.md`.
[56] Vacancy lift numerics, `contrib/vacancy_mobile_numerics/README.md`.
[57] gpt-6-astra runs C64 to C79, `codex/work_C64/` to `work_C79/`.
[58] The $n = 2$ vacancy model, `contrib/vacancy_n2_dmrg/`.
[59] S on the microscopic demonstrator, `contrib/demonstrator_luttinger/`.
[60] Referee reports on C84, C85 and C88, `contrib/referee_C84_C85/`, `contrib/referee_C88/`.
[61] The $d = 2$ surveys, `contrib/d2_grid/`, `contrib/d2_staggered/`.
[62] Literature check of the integrate-out-then-bosonize step, `contrib/lit_ir_step/README.md`.