

Attentional modulation of functional connectivity in the face processing network of the brain

Find more info here:



UNIVERSITÀ
DI TRENTO

Jan-Luca Schröder^{1,2} & Daniel Baldauf¹

¹Center for Mind/Brain Sciences (CIMEC), University of Trento, Italy

²Institute of Cognitive Science, Universität Osnabrück, Osnabrück, Germany


Introduction

- The face processing network, including regions like the FFA [3], OFA [5] and a region in the STS [4], is most important for our social interactions
- The prefrontal cortex was shown to play a crucial role guiding selective attention, especially the IFJ (non-spatial attention, object and feature encoding) and the FEF (spatial attention) [6, 7]
- Research question:**
Is there a modulation of **spectral activity** and **functional connectivity patterns** in a **face detection task** based on what the **covert spatial attention** is set on?

Methods

- 10 participants saw composite face stimuli in the MEG and had to attend either to the **eyes**, the **mouth** or the facial **identity** [1]
- We applied the HCP-MMP 1.0 atlas [2] onto the individual anatomy of subjects
- The Regions of interest were:
IFJa/p, FEF, FFC, VVC, V4, TPOJ2, LIPd, MT, PeEC
- Analysis in **theta** (5 – 8 Hz), **alpha** (8 – 12 Hz), **beta** (15 – 25 Hz), **gamma** (30 – 100 Hz)
- Analysis: ‘Attend IN – Attend OUT’ contrast**
(Mean of non-attended conditions subtracted from attended condition)


Paradigm



Note. Examples of the database of stimuli and the respective trial sequence used in the study.


Figure adapted from [1]

Regions of Interest




- Regions: IFJa/p, FEF, FFC, VVC, V4, TPOJ2, LIPd, MT, PeEC
- Performed anatomic likelihood estimation (ALE) for FFA, OFA, STS

Results – spectral activity



Results – functional connectivity



Conclusion

- Attention modulates spectral activity and functional connectivity
- FEF and IFJa → crucial role in selective attention during face processing
- Difference between holistic and single facial feature processing

Discussion

Attend to mouth

- The spectral inhibition and functional connectivity in FEF's theta and beta band suggest lesser effect of spatial attention
- The spectral excitation of MT in theta suggests perception of motion or activity of OFA
- The functional connectivity between IFJa and TPOJ2 in alpha suggests increase feature attention and involvement of STS

Attend to identity

- The spectral excitation of the MT in alpha suggests perception of motion
- The functional connectivity of the FEF suggests an effect of spatial attention
- The spectral excitation and functional connectivity of the PeEC in alpha suggest processing of identity

Attend to eyes

- The functional connectivity between IFJa and TPOJ2 suggests an effect of feature attention and an involvement of the STS

References

- [1] de Vries, E., & Baldauf, D. (2019). Attentional Weighting in the Face Processing Network: A Magnetic Response Image-guided Magnetoencephalography Study Using Multiple Cyclic Entrainments. *Journal of Cognitive Neuroscience*, 31 (10), 1573–1588.
- [2] Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A Multi-Modal Parcellation of Human Cerebral Cortex. *Nature*, 536(7615), 171–178.
- [3] Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The Fusiform Face Area: A Module in Human Extrastriate Cortex Specialized for Face Perception.
- [4] Haxby, J. V., Hoffman, E. A., & Gobbini, M. (2000). The Distributed Human Neural System for Face Perception. *Trends in Cognitive Sciences*, 4 (6), 223–233.
- [5] Puce, A., Allison, T., Asgari, M., Gore, J. C., & McCarthy, G. (1996). Differential Sensitivity of Human Visual Cortex to Faces, Letterstrings, and Textures: A Functional Magnetic Resonance Imaging Study. *The Journal of Neuroscience*, 16(16), 5205–5215.
- [6] Bedini, M., Olivetti, E., Avesani, P., & Baldauf, D. (2023). Accurate Localization and Coactivation Profiles of the Frontal Eye Field and Inferior Frontal Junction: An ALE and MACM fMRI Meta-Analysis. *Brain Structure and Function*, 228(3–4), 997–1017.
- [7] Soyuhos, O., & Baldauf, D. (2023). Functional Connectivity Fingerprints of the Frontal Eye Field and Inferior Frontal Junction Suggest Spatial versus Nonspatial Processing in the Prefrontal Cortex. *European Journal of Neuroscience*, 57(7), 1114–1140.