

PSET 2

(ASSUMPTION OF VARIOUS PROPERTIES OF \exp , \ln FUNCTIONS) Lin. Alg.

Jonathan Lam
Prof. Minkowski
MA 326
Lin. Alg.
9/19/19

1. Let $V = \mathbb{R}^+ = (0, +\infty)$, $F = \mathbb{R}$. 27/30

DEF: Given $x, y \in V$, $x \cdot y := xy$ (real multiplication)

DEF: Given $a \in F$, $x \in V$, $a \cdot x := x^a = \exp(a \ln(x))$ (real exponentiation)

Show that $(V, +_v, \cdot_v)$ is a v.s. over \mathbb{R} .

~~1.3.23 | 5~~

~~1.4.13 | 5~~

10

CLAIM: \cdot_v is binary over V .

PROOF: $\forall x, y \in V$, $x, y \in \mathbb{R}$, and $xy \in \mathbb{R}$ (since \cdot is binary over \mathbb{R}). Also, since $x, y > 0$, $xy > 0$ (two positive reals multiply to a positive real), so $xy \in \mathbb{R}^+$.

Then \cdot_v maps $V \times V \rightarrow V$ and thus is binary over V .

CLAIM: \cdot_v is external binary on V with field F .

PROOF: $\forall a \in F$, $x \in V$. $a \cdot_v x = \exp(a \ln(x))$.

$(\ln(x))$ exists and $\in \mathbb{R}$ $\forall x \in \mathbb{R}^+ = V$, so $a \ln(x) \in \mathbb{R}$

(since \cdot is binary over \mathbb{R}). $\forall c \in \mathbb{R}$, $\exp(c) \in \mathbb{R}^+$

(property of \exp), thus $\exp(a \ln(x)) \in \mathbb{R}^+ = V$, i.e.,

\cdot_v maps $F \times V \rightarrow V$ and thus is external binary on V with field F .

VS1) CLAIM: $\forall x, y \in V$, $x +_v y = y +_v x$.

PROOF: $\forall x, y \in V$,

$$x +_v y = xy = yx = y +_v x$$

commutativity of \cdot over \mathbb{R}

VS2) CLAIM: $\forall x, y, z \in V$, $x +_v (y +_v z) = (x +_v y) +_v z$

PROOF: $\forall x, y, z \in V$,

$$x +_v (y +_v z) = x +_v (yz) = x(yz) = (xy)z = (xy) +_v z = (x +_v y) +_v z$$

associativity of \cdot over \mathbb{R} .

VS3) CLAIM: $\exists 0_V \in V$ s.t. $\forall x \in V$, $0_V +_v x = x$

PROOF: Let $0_V = 1_{(\mathbb{A})}$. Then, $\forall x \in V$,

$$0_V +_v x = 1x = x$$

(F3) on \mathbb{R}

VS4) CLAIM: $\forall x \in V, \exists y \in V$ s.t. $x +_V y = 0_V$.

PROOF: Let $x \in V = \mathbb{R}^+$. Then $\exists y = \frac{1}{x}$. Since $x \neq 0 \in \mathbb{R}$, $y \in \mathbb{R}$, and since $1, x > 0$, $y = \frac{1}{x} > 0 \in \mathbb{R}^+ = V$, and:
 $x +_V y = x\left(\frac{1}{x}\right) = 1_{(\mathbb{R})} = 0_V$

VS5) CLAIM: $\forall x \in V, 1 \cdot_V x = x$

PROOF: $\forall x \in V$:

$$\begin{aligned} 1 \cdot_V x &= \exp(1 \cdot \ln x) \\ &= \exp(\ln x) && (\text{F3 of } \mathbb{R}) \\ &= x && (\exp = \ln^{-1}) \end{aligned}$$

VS6) CLAIM: $\forall a, b \in F, \forall x \in V, (ab) \cdot_V x = a \cdot_V (b \cdot_V x)$

PROOF: $\forall a, b \in F, \forall x \in V$,

$$\begin{aligned} (ab) \cdot_V x &= \exp(ab \ln(x)) = \exp(a(b \ln x)) \\ &= \exp(a \ln(\exp(b \ln x))) && (\exp = \ln^{-1}) \\ &= \exp(a \ln(b \cdot_V x)) \\ &= a \cdot_V (b \cdot_V x) \end{aligned}$$

VS7) CLAIM: $\forall a \in F, x, y \in V, a \cdot_V (x +_V y) = a \cdot_V x +_V a \cdot_V y$.

PROOF: $\forall a \in F, x, y \in V$,

$$\begin{aligned} a \cdot_V (x +_V y) &= a \cdot_V (xy) = \exp(a \ln(xy)) \\ &= \exp(a(\ln x + \ln y)) && (\text{prop. of } \ln) \\ &= \exp(a \ln x + a \ln y) \\ &= \exp(a \ln x) \cdot \exp(a \ln y) && (\text{prop. of } \exp) \\ &= (a \cdot_V x) \cdot (a \cdot_V y) = a \cdot_V x +_V a \cdot_V y. \end{aligned}$$

VS8) CLAIM: $\forall a, b \in F, x \in V, (a+b) \cdot_V x = a \cdot_V x +_V b \cdot_V x$.

PROOF: $\forall a, b \in F, x \in V$,

$$\begin{aligned} (a+b) \cdot_V x &= \exp((a+b) \ln(x)) = \exp(a \ln x + b \ln x) \\ &= \exp(a \ln(x)) \cdot \exp(b \ln(x)) && (\text{prop. of } \exp) \\ &= (a \cdot_V x) \cdot (b \cdot_V x) = a \cdot_V x +_V b \cdot_V x \end{aligned}$$

Since $+_V$ is binary on V , \cdot_V is external binary on V with field F , and V satisfies (VS1-8), $(V, +_V, \cdot_V)$ is a vector space.

PSET 2, cont'd

Jonathan Low
Peer-Matched
MA 326
Lin. Alg.
9/19/19

2. DEF: $C(\mathbb{R}) := \{f: \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ is continuous on } \mathbb{R}\}$

DEF: $C^n(\mathbb{R}) := \{f: \mathbb{R} \rightarrow \mathbb{R} \mid f^{(n)} \text{ is continuous on } \mathbb{R}\}$

a. Show that $C(\mathbb{R})$ is a v.s. over \mathbb{R} .

CLAIM: $C(\mathbb{R}) \subseteq F(\mathbb{R}, \mathbb{R})$

The following proofs will prove that $C(\mathbb{R}) \subseteq F(\mathbb{R}, \mathbb{R})$ using (THM 1.3), the necessary conditions for a subspace.

CLAIM: $C(\mathbb{R}) \subseteq F(\mathbb{R}, \mathbb{R})$

PROOF: $F(\mathbb{R}, \mathbb{R})$ is the set of all functions $f: \mathbb{R} \rightarrow \mathbb{R}$ over the domain \mathbb{R} . $C(\mathbb{R})$ is a set of some functions $f: \mathbb{R} \rightarrow \mathbb{R}$ that exist (and are continuous) over the domain \mathbb{R} , thus $C(\mathbb{R}) \subseteq F(\mathbb{R}, \mathbb{R})$.

DEF: $z := z(x) = 0 \quad \forall x \in \mathbb{R}$

(Note that this is the additive identity for $F(\mathbb{R}, \mathbb{R})$, since $\forall f \in F(\mathbb{R}, \mathbb{R})$, $f + z = f(x) + z(x) = f(x) + 0 = f(x) = f$).

CLAIM: $z \in C(\mathbb{R})$.

PROOF: $z(a) = \lim_{x \rightarrow a} z(x) = 0 \quad \forall a \in \mathbb{R}$, so z is cts. over \mathbb{R} and $z \in C(\mathbb{R})$

CLAIM: $\forall f, g \in C(\mathbb{R})$, $f + g \in C(\mathbb{R})$.

PROOF: $\forall a \in \mathbb{R}$,

$$\begin{aligned} f(a) &= \lim_{x \rightarrow a} f(x), \quad g(a) = \lim_{x \rightarrow a} g(x), \quad (\text{DEF continuity}) \\ f + g &= f(a) + g(a) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \\ &= \lim_{x \rightarrow a} (f(x) + g(x)) \quad (\text{linearity of lim}) \end{aligned}$$

Therefore $f + g$ is continuous on \mathbb{R} , so $f + g \in C(\mathbb{R})$.

CLAIM: $\forall a \in \mathbb{R}$, $\forall f \in C(\mathbb{R})$, $af \in C(\mathbb{R})$

PROOF: $\forall b \in \mathbb{R}$, $f(b) = \lim_{x \rightarrow b} f(x)$, (DEF continuity)

$$af = af(b) = a \lim_{x \rightarrow b} f(x) = \lim_{x \rightarrow b} af(x) \quad (\text{linearity of lim})$$

Therefore af is continuous on \mathbb{R} , so $af \in C(\mathbb{R})$

By THM 1.3, $C(\mathbb{R})$ SUBSP. $\mathcal{F}(\mathbb{R})$. Thus $C(\mathbb{R})$ is a v.s.

b. Show that $(^{(n)})(\mathbb{R})$ SUBSP. $C(\mathbb{R}) \quad \forall n \in \mathbb{Z}^+$

CLAIM: $(^{(n)})(\mathbb{R}) \subseteq C(\mathbb{R}), \forall n \in \mathbb{Z}^+$

PROOF: $\forall f \in (^{(n)})(\mathbb{R}), \forall n \in \mathbb{Z}^+, f$ is n times differentiable ($n > 0$).

Any differentiable function is continuous over its domain,
thus $f \in C(\mathbb{R})$ and $(^{(n)})(\mathbb{R}) \subseteq C(\mathbb{R})$.

CLAIM: $z \in (^{(n)})(\mathbb{R}), \forall n \in \mathbb{Z}^+ \quad (\text{using same (DEF } z\text{)})$

PROOF: $\forall n \in \mathbb{Z}^+, z^{(n)} = z^{(n)}(x) = 0 = z \in C(\mathbb{R}) \quad \forall x \in \mathbb{R}$.

CLAIM: $\forall f, g \in (^{(n)})(\mathbb{R}), \forall n \in \mathbb{Z}^+, f + g \in (^{(n)})(\mathbb{R})$

PROOF: $\forall f, g \in (^{(n)})(\mathbb{R}), \forall n \in \mathbb{Z}^+, \forall a \in \mathbb{R},$

$$f^{(n)}(a) = \lim_{x \rightarrow a} f^{(n)}(x), \quad (\text{DEF continuity})$$

$$g^{(n)}(a) = \lim_{x \rightarrow a} g^{(n)}(x), \quad (\text{DEF continuity})$$

$$\frac{d}{dx^n}(f+g) = \frac{d}{dx^n}(f(a) + g(a)) \quad (\text{linearity of } \frac{d}{dx})$$

$$= f^{(n)}(a) + g^{(n)}(a) = \lim_{x \rightarrow a} f^{(n)}(x) + \lim_{x \rightarrow a} g^{(n)}(x)$$

$$= \lim_{x \rightarrow a} (f^{(n)}(x) + g^{(n)}(x)) \quad (\text{linearity of lim})$$

$$= \lim_{x \rightarrow a} (f^{(n)} + g^{(n)}) = \lim_{x \rightarrow a} \left(\frac{d}{dx^n} (f+g) \right) \quad (\text{linearity of } \frac{d}{dx})$$

Thus $f+g$ has a continuous n th-derivative, so $f+g \in (^{(n)})(\mathbb{R})$.

CLAIM: $\forall f \in (^{(n)})(\mathbb{R}), \forall a \in \mathbb{R}, \forall n \in \mathbb{Z}^+, af \in (^{(n)})(\mathbb{R})$.

PROOF: $\forall f \in (^{(n)})(\mathbb{R}), \forall a, b \in \mathbb{R},$

$$f^{(n)}(b) = \lim_{x \rightarrow b} f^{(n)}(x), \quad (\text{DEF continuity})$$

$$\frac{d}{dx^n}(af) = a \frac{d}{dx^n} f = af^{(n)}(b)$$

$$= a \lim_{x \rightarrow b} f^{(n)}(x) = \lim_{x \rightarrow b} (af^{(n)}(x))$$

$$= \lim_{x \rightarrow b} \left(\frac{d}{dx^n} (af) \right)$$

Thus af has a continuous n th-derivative, so $af \in (^{(n)})(\mathbb{R})$

2b, (contd.) By Thm 1.3, $C^{(i)}(\mathbb{R})$ subsp. $C(\mathbb{R}) \ \forall n \in \mathbb{Z}^+$.

2c. Show $j > i \Rightarrow C^{(j)}(\mathbb{R})$ subsp. $C^{(i)}(\mathbb{R})$, $j, i \in \mathbb{Z}^+$.

CLAIM: $C^{(j)}(\mathbb{R}) \subseteq C^{(i)}(\mathbb{R})$

PROOF: $\forall f \in C^{(j)}(\mathbb{R})$, $f^{(j)}$ is continuous. In general,
for $\forall k \in \mathbb{Z}^+$, if $f^{(k+1)}$ is continuous, $f^{(k)}$ is continuous.

By induction, $f^{(j)}, f^{(j-1)}, \dots, f^{(1)}$ are continuous.

Since $i < j$, $f^{(i)}$ is continuous. Thus $f \in C^{(i)}(\mathbb{R})$,
so $C^{(j)}(\mathbb{R}) \subseteq C^{(i)}(\mathbb{R})$.

Since $C^{(n)}(\mathbb{R})$ is a v.s. (proved in (2b)), then $C^{(i)}(\mathbb{R})$,
 $C^{(j)}(\mathbb{R})$ are v.s. over \mathbb{R} . Since $C^{(j)}(\mathbb{R}) \subseteq C^{(i)}(\mathbb{R})$,
 $C^{(j)}(\mathbb{R})$ subsp. $C^{(i)}(\mathbb{R})$.

SET 2 #3

Jonathan Lam
Prof. Mutchler
MA 326
Lin. Alg.
9/19/19

3. Let $V = \mathcal{F}(\mathbb{R}, \mathbb{R})$,

$$E = \{f \in V \mid f(-x) = f(x) \quad \forall x \in \mathbb{R}\}$$

$$O = \{f \in V \mid f(-x) = -f(x) \quad \forall x \in \mathbb{R}\}$$

a. Show E and O are proper subspaces of V .

CLAIM: E, O are proper subsets of V ($E, O \subseteq V, E, O \neq V$).

PROOF: $\forall f \in E, \forall g \in O$, by (DEF E) and (DEF O),
 $f \in V, g \in V$, thus $E, O \subseteq V$.

$\forall h \neq 0 \in E, i \neq 0 \in O$. Then, $h \notin O$ ($h(-x) = h(x) \neq -h(x)$)
and $i \notin E$ ($i(-x) = -i(x) \neq i(x)$). Since $h \in V, i \in V$,
 $E, O \neq V$.

CLAIM: $z \in E, O$. (using same (DEF z) from problem 2)

(Note: z is the 0 vector of V .)

PROOF: $\forall x, z = z(x) = 0 = z(-x)$, so $z \in E$,
 $z = z(x) = 0 = (-1) \cdot 0 = -z(x)$, so $z \in O$.

CLAIM: $\forall f, g \in E, f + g \in E$,

$\forall h, i \in O, h + i \in O$

PROOF: $\forall f, g \in E, \forall x \in \mathbb{R}$,

$$f+g = (f+g)(x) = f(x) + g(x) = f(-x) + g(-x) = (f+g)(-x)$$

$$\therefore f+g \in E.$$

$\forall h, i \in O, h + i \in O$.

$$(h+i)(-x) = h(-x) + i(-x) = -h(x) - i(x)$$

$$= -1(h(x) + i(x)) = -(h+i)(x)$$

$$\therefore h+i \in O.$$

CLAIM: $\forall a \in \mathbb{R}, \forall f \in E, af \in E,$
 $\forall b \in \mathbb{R}, \forall g \in O, bg \in O.$

PROOF: $\forall a \in \mathbb{R}, \forall f \in E,$

$$af = (af)(x) = af(x) = af(-x) = (af)(-x)$$

$$\therefore af \in E.$$

$\forall b \in \mathbb{R}, \forall g \in O,$

$$(bg)(x) = bg(-x) = b(-g(x))$$
$$= -bg(x) = -(bg)(x)$$

$$\therefore bg \in O.$$

By (THM 1.3), E, O subsp. V . Since E, O are proper subsets of V , E, O proper subsp. V .

b. Argue that $\text{span}(E) = E$, $\text{span}(O) = O$.

LEM 1: $\forall V$ v.s., $\text{span}(V) = V$.

PROOF: $V \subseteq V$, so $\text{span}(V) \subseteq V$ (THM 1.5)

Also, if $x \in V$, x can be written

as a linear combination of vectors in V ,

i.e., $1 \cdot x = x$. Thus $V \subseteq \text{span}(V)$.

Since $\text{span}(V) \subseteq V$; $V \subseteq \text{span}(V)$, $V = \text{span}(V)$.

CLAIM: $\text{span}(E) = E$, $\text{span}(O) = O$.

PROOF: E and O are both v.s. (proved in part a).

(LEM 1) states that the span of any v.s. is itself,
so this result is immediate.

PSET 2, P3, cont'd.

Jonathan Lam
Prof. Mutchler
MA 326
Lin. Alg.
9/19/14

3c) Show that $E \cup O$ generates V .

CLAIM: $V \subseteq \text{span}(E \cup O)$

PROOF: $\forall f \in V$, let $f_e = f_e(x) = \frac{f(x) + f(-x)}{2}$,

$$f_o = f_o(x) = \frac{f(x) - f(-x)}{2}.$$

$$\text{Since } f_e(-x) = \frac{f(-x) + f(x)}{2} = \frac{f(x) + f(-x)}{2} = f_e(x),$$

and $f_e: \mathbb{R} \rightarrow \mathbb{R}$, $f_e \in E$.

$$\text{Since } f_o(-x) = \frac{f(-x) - f(x)}{2} = -\left(\frac{f(x) - f(-x)}{2}\right) = -f_o(x),$$

and $f_o: \mathbb{R} \rightarrow \mathbb{R}$, $f_o \in O$.

Then $\forall g \in V$, g can be expressed as the linear combination of g_e, g_o :

$$1 \cdot g_e + 1 \cdot g_o = g_e(x) + g_o(x) = \frac{g(x) + g(-x)}{2} + \frac{g(x) - g(-x)}{2} \\ = \left(\frac{g(x)}{2} + \frac{g(x)}{2}\right) + \left(\frac{g(x) - g(-x)}{2}\right) = g(x) = g.$$

Since $g_e \in E$, $g_o \in O$, $g_e, g_o \in E \cup O$, and
 $V \subseteq \text{span}(E \cup O)$.

CLAIM: $E \cup O$ generates V .

PROOF: Since $E, O \subseteq V$, $E \cup O \subseteq V$. Then, by

(Thm 1.5), $\text{span}(E \cup O) \subseteq V$. Since $V \subseteq \text{span}(E \cup O)$,

$V = \text{span}(E \cup O)$ $\therefore E \cup O$ generates V .

1.2 # 8, 9, 21

8.9, 5 TEC
7/17,

CLAIM:

8. \forall v.s. V , show $(a+b)(x+y) = ax + ay + bx + by$.
 $\forall x, y \in F$, $\forall a, b \in F$.

PROOF: $(a+b)(x+y)$

$$= (a+b)x + (a+b)y \quad (\text{VS 7})$$

$$= ax + bx + ay + by \quad (\text{VS 8})$$

$$= ax + ay + bx + by \quad (\text{VS 1})$$

9. Prove corollaries 1, 2 of (THM 1.1), (THM 1.2 c).

(THM 1.1)
(cor. 1)

CLAIM: The vector 0 described in (VS3) is unique.

PROOF: Let $0_1, 0_2$ satisfy additive identity property described in (VS3). Then, $\forall x \in V$,

$$0_1 + x = x = 0_2 + x$$

By (THM 1.2), $0_1 = 0_2$ \therefore the additive identity for a v.s. is unique.

(THM 1.1)
(cor. 2)

CLAIM: The vector y described in (VS4) is unique.

PROOF: Let y_1, y_2 satisfy additive inverse property described in (VS4) if $x \in V$. Then,

$$x + y_1 = 0 = x + y_2$$

By (THM 1.1), $y_1 = y_2$ \therefore the additive inverse $\forall x \in V$ is unique.

(THM 1.2 c) CLAIM: $a0 = 0$ $\forall a \in F$ in any v.s. over F .

PROOF: $\forall a \in F$,

$$a0 + a0 = \underbrace{a(0+0)}_{\text{by (VS7)}} = \underbrace{a0}_{(\text{VS3})} = \underbrace{a0 + 0}_{(\text{VS3})}$$

By (THM 1.1), $a0 = 0$.

21. Let V, W be v.s. over field F . Let $Z = \{(v, w) : v \in V, w \in W\}$.

Prove that Z v.s. over F w/ the operations:

$$(v_1, w_1) + (v_2, w_2) := (v_1 + v_2, w_1 + w_2),$$

$$c(v_1, w_1) := (cv_1, cw_1)$$

CLAIM: + binary over Z .

PROOF: $\forall z_1 = (v_1, w_1), z_2 = (v_2, w_2) \in Z$. Then $v_1, v_2 \in V$, $w_1, w_2 \in W$, and by binary-ness of vector addition, $v_1 + v_2 \in V$, $w_1 + w_2 \in W$. $\therefore z_1 + z_2 \in Z$, so + maps $Z \times Z \rightarrow Z$ and is thus binary.

CLAIM: - is external binary over Z w/ field F .

PROOF: $\forall c \in F, \forall z = (v, w)$. Since v, w are v.s. over field F , scalar multiplication by F is external binary, i.e., $cv \in V$, $cw \in W$. $\therefore cz \in Z$, so - maps $F \times Z \rightarrow Z$, and is therefore external binary over F .

VS1) CLAIM: $\forall z_1, z_2 \in Z, z_1 + z_2 = z_2 + z_1$.

PROOF: $\forall z_1 = (v_1, w_1), z_2 = (v_2, w_2) \in Z$,

$$z_1 + z_2 = (v_1 + v_2, w_1 + w_2) = \underbrace{(v_2 + v_1, w_2 + w_1)}_{(VS1 \text{ for } W, V)} = z_2 + z_1$$

VS2) CLAIM: $\forall z_1, z_2, z_3 \in Z, z_1 + (z_2 + z_3) = (z_1 + z_2) + z_3$.

PROOF: $\forall z_1 = (v_1, w_1), z_2 = (v_2, w_2), z_3 = (v_3, w_3)$,

$$\begin{aligned} z_1 + (z_2 + z_3) &= (v_1, w_1) + (v_2 + v_3, w_1 + w_2 + w_3) \\ &= (v_1 + (v_2 + v_3), w_1 + (w_2 + w_3)) \\ &= ((v_1 + v_2) + v_3, (w_1 + w_2) + w_3) \quad (VS2 \text{ for } V, W) \\ &= (v_1 + v_2, w_1 + w_2) + (v_3, w_3) \\ &= (z_1 + z_2) + z_3 \end{aligned}$$

VS3) CLAIM: $\forall z \in Z, z + 0 = z$.

(Proven directly by $(v, w) + (0, 0) = (v, w)$ additivity from v, w)

VS4) CLAIM: $\forall z \in Z, z + (-z) = 0$.

(Proven directly by $(v, w) + (-v, -w) = (v - v, w - w) = (0, 0)$ for v, w)

1.2 # 21, cont'd.

VS3) CLAIM: $\exists \mathbf{0}_z \in \mathbb{Z}$ s.t. $\forall z \in \mathbb{Z}, \mathbf{0}_z + z = z$.

PROOF: \exists additive identities $\mathbf{0}_v \in V, \mathbf{0}_w \in W$ (VS3 for v, w).

Let $\mathbf{0}_z = (\mathbf{0}_v, \mathbf{0}_w)$. Then, $\forall z = (v, w) \in \mathbb{Z}$,

$$\begin{aligned} \mathbf{0}_z + z &= (\mathbf{0}_v + v, \mathbf{0}_w + w) \\ &= (v, w) = z. \end{aligned} \quad (\text{VS3 for } v, w)$$

VS4) CLAIM: $\forall z_1 \in \mathbb{Z}, \exists z_2 \in \mathbb{Z}$ s.t. $z_1 + z_2 = \mathbf{0}_z$.

PROOF: $\forall z_1 = (v, w) \in \mathbb{Z}$,

v has an additive inverse ($-v$), $(\text{VS4 for } v)$

w has an additive inverse ($-w$), $(\text{VS4 for } w)$

Let $z_2 = (-v, -w)$, then:

$$\begin{aligned} z_1 + z_2 &= (v + (-v), w + (-w)) \\ &= (\mathbf{0}_v, \mathbf{0}_w) = \mathbf{0}_z. \end{aligned} \quad (\text{VS4 for } v, w)$$

VS5) CLAIM: $\forall z \in V, 1z = z$.

PROOF: $\forall z = (v, w) \in V, 1z = (1v, 1w) = (\underbrace{v, w}) = v$

$(\text{VS5 for } v, w)$

VS6) CLAIM: $\forall a, b \in F, \forall z \in \mathbb{Z}, (a \cdot b)z = a \cdot (b \cdot z)$

PROOF: $\forall a, b \in F, \forall z = (v, w) \in \mathbb{Z}$,

$$(a \cdot b)z = ((a \cdot b)v, (a \cdot b)w) = (a \cdot (bv), a \cdot (bw)) = a(bv, bw) = a(bz) \quad (\text{VS6 for } v, w)$$

VS7) CLAIM: $\forall a \in F, \forall z_1, z_2 \in \mathbb{Z}, a(z_1 + z_2) = az_1 + az_2$

PROOF: $\forall a \in F, \forall z_1 = (v_1, w_1), z_2 = (v_2, w_2) \in \mathbb{Z}$,

$$a(z_1 + z_2) = a(v_1 + v_2, w_1 + w_2) = (a(v_1 + v_2), a(w_1 + w_2))$$

$$= (av_1 + av_2, aw_1 + aw_2) = (av_1, aw_1) + (av_2, aw_2) = az_1 + az_2$$

\curvearrowright (VS7 for v, w)

VS8) CLAIM: $\forall a, b \in F, \forall z \in \mathbb{Z}, (a + b)z = az + bz$.

PROOF: $\forall a, b \in F, \forall z = (v, w) \in \mathbb{Z}$,

$$(a + b)z = ((a + b)v, (a + b)w) = (av + bv, aw + bw) \quad (\text{VS8 for } v, w)$$

$$= (av, aw) + (bv, bw) = az + bz.$$

Since $+$ binary on \mathbb{Z} , \cdot external binary on \mathbb{Z} , and (VS1-8)

Satisfied, \mathbb{Z} is a U.S over F with the addition and multiplication defined here.

PSET 2

1.3 #10, 22, 23, 25.

Jonathan Loom
Prof. Mirtchev
MA 326
Lin. Alg.
9/19/19

10. Prove that $W_1 = \{(\alpha_1, \alpha_2, \dots, \alpha_n) \in F^n : \alpha_1 + \alpha_2 + \dots + \alpha_n = 0\}$
 is a subspace, but $W_2 = \{(\alpha_1, \alpha_2, \dots, \alpha_n) \in F^n : \alpha_1 + \alpha_2 + \dots + \alpha_n = 1\}$
 is not.

a. Prove W_1 is a subspace:

Note: the zero vector for F^n is known to be $(\underbrace{0, 0, 0, \dots, 0}_n)$.
 Denote zero vector for F^n 0_n ($n \in \mathbb{Z}^+$).

CLAIM: $0_n \in W_1$.

PROOF: In 0_n , $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$, so $\alpha_1 + \alpha_2 + \dots + \alpha_n = 0$.
 Thus $0_n \in W_1$.

CLAIM: $\forall x = (\alpha_1, \alpha_2, \dots, \alpha_n), \forall y = (b_1, b_2, \dots, b_n) \in W_1$,
 then $x+y \in W_1$.

PROOF: $x+y = (\alpha_1+b_1, \alpha_2+b_2, \dots, \alpha_n+b_n)$. Summing its terms:
 $(\alpha_1+b_1) + (\alpha_2+b_2) + \dots + (\alpha_n+b_n)$
 $= (\alpha_1 + \alpha_2 + \dots + \alpha_n) + (b_1 + b_2 + \dots + b_n)$ (F1, F2 for F)
 $= 0 + 0 = 0$.

Thus $x+y \in W_1$.

CLAIM: $\forall c \in F, \forall x = (\alpha_1, \alpha_2, \dots, \alpha_n) \in W_1, cx \in W_1$.

PROOF: $cx = (c\alpha_1, c\alpha_2, \dots, c\alpha_n)$. Summing the terms:
 $c\alpha_1 + c\alpha_2 + \dots + c\alpha_n$

$$= c(\alpha_1 + \alpha_2 + \dots + \alpha_n) = c(0) = 0$$

Thus $cx \in W_1$.

By (THM 1.3), W_1 satisfies the necessary conditions to be subsp. F^n .

b) CLAIM: W_2 not subsp. F^n .

PROOF: For 0_n , $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$. Thus, $\alpha_1 + \alpha_2 + \dots + \alpha_n = 0 \neq 1$,
 so $0_n \notin W_2$. By (THM 1.3), W_2 is not subsp. F^n
 because it doesn't contain the zero vector of F^n .

22. Let F_1, F_2 be fields.

DEF: $E := \{ f: F_1 \rightarrow F_2 : f(-t) = f(t) \}$.

DEF: $O := \{ O: F_1 \rightarrow F_2 : f(-t) = -f(t) \}$.

DEF: an element of E is called even, and an element of O is called odd.

(a) Prove E, O subsp. $\tilde{F}(F_1, F_2)$.

CLAIM: $E, O \subseteq \tilde{F}(F_1, F_2)$

This is immediate from the definitions of $\tilde{F}(F_1, F_2)$, and E, O .

DEF: Let $z_2 := z_2(x) = 0_2$, $x_1 \in F_1$, $0_2 \in F_2$ and is the zero element in F_2 . This is the zero vector of $\tilde{F}(F_1, F_2)$ (since addition of any element of $\tilde{F}(F_1, F_2)$ with z_2 results in an addition with the additive identity in F_2).

CLAIM: $z_2 \in E, O$.

PROOF: $\forall a \in F_1$, $z_2(a) = 0 = z_2(a)$, so $z_2 \in E$.

$\forall a \in F_1$, $z_2(-a) = 0 = (-1)(0) = -z_2(a)$, so $z_2 \in O$.

CLAIM: $\forall f, g \in E$, $f + g \in E$.

$\forall u, v \in O$, $u + v \in O$.

PROOF: $\forall f, g \in E$, $\forall a \in F_1$,

$$(f+g)(a) = f(-a) + g(-a) = f(a) + g(a) = (f+g)(a),$$

so $f+g \in E$.

$$\begin{aligned} (u+v)(-a) &= u(-a) + v(-a) = -u(a) - v(a) = -1(u(a) + v(a)) \\ &= -(u+v)(a), \text{ so } u+v \in O. \end{aligned}$$

CLAIM: $\forall c \in F_2$, $\forall f \in E$, $\forall g \in O$, $cf \in E$, $cg \in O$.

PROOF: $\forall a \in F_1$, $(cf)(a) = cf(a) = cf(a) = (cf)(a)$, so $cf \in E$.

$$(cg)(-a) = cg(-a) = c(-g(a)) = -(cg)(a), \text{ so } cg \in O.$$

By Thm 1.3, the necessary conditions we met to prove E, O subsp. $\tilde{F}(F_1, F_2)$.

23. Let w_1, w_2 subsp. v.s. V.

a) Prove that $w_1 + w_2$ subsp. V and contains both w_1 and w_2 .

5 Let $w = w_1 + w_2$ (i.e., $w = \{x + y : x \in w_1, y \in w_2\}$)

CLAIM: $w \in V$?

You mean $w \subseteq V$.

PROOF: $\forall w = x + y \in V, x \in w_1 \subseteq V, y \in w_2 \subseteq V$.

Since + is binary over vector addition $x + y \in V$.

$\therefore w \in V$

CLAIM: $0_V \in w$.

PROOF: All subspaces contain the zero vector of their superspace.

Thus $0_V \in w_1, w_2$, and $\exists x \in w = 0_V + 0_V = 0_V$.

CLAIM: $\forall x, y \in w, x + y \in w$.

PROOF: If $x = x_1 + x_2, y = y_1 + y_2 \in w, x_1, y_1 \in w_1$,

$x_2, y_2 \in w_2$. Then $x + y = x_1 + x_2 + y_1 + y_2$

$= (x_1 + y_1) + (x_2 + y_2)$. By binarity of + over v.s.,

$x_1 + y_1 \in w_1, x_2 + y_2 \in w_2$, thus $x + y \in w$

CLAIM: $\forall a \in F, \forall x \in w, ax \in w$.

PROOF: If $a \in F, \forall x = x_1 + x_2 \in w, x_1 \in w_1, x_2 \in w_2$,

$$ax = a(x_1 + x_2) = ax_1 + ax_2$$

(VS for V)

Scalar multiplication is external binary over a v.s.,

so $ax_1 \in w_1, ax_2 \in w_2$, so $ax \in w$.

By (Thm 1.3), w satisfies the conditions to be a subsp. of V.

CLAIM: W contains both w_1, w_2 .

PROOF: Since $0_v \in W_2$, $\forall x \in W$, $\exists w \in W = x + 0_v = x$

Similarly, since $0_v \in W_1$, $\forall y \in W_2$,

$$\exists w = 0_v + y = y. \text{ Thus } w_1, w_2 \subseteq W.$$

b. Prove that any subspace Z of V that contains both w_1 and w_2 must also contain W .

CLAIM: Sub Z contains $w_1, w_2 \Rightarrow Z$ contains W .

PROOF: $\forall x \in W_1$, $\forall y \in W_2$, $x + y \in Z$ because

+ is binary over vector addition. Therefore,

$$\text{the set of } \{x + y : x \in w_1, y \in w_2\} = W \subseteq Z.$$

25. Let W_1 denote set of all polynomials in $P(F)$ s.t. in the representation $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$,

we have $a_i = 0$ when i even. Also define W_2 set

of all polynomials in $P(F)$ s.t. in the representation:

$$g(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0, \text{ we have } b_i = 0 \text{ whenever } i \text{ is odd.}$$

Prove $P(F) = W_1 \oplus W_2$.

To show that $P(F)$ is the direct sum of W_1, W_2 ,

need to show that: (by (DEF direct sum)):

$$1) W_1, W_2 \text{ subsp } P(F)$$

$$2) W_1 \cap W_2 = \{0_p\}$$

$$3) W_1 + W_2 = V.$$

25.1) DEF: Let $0_p := 0 + 0x + 0x^2 + \dots$ be the zero vector of $P(F)$.

CLAIM: $0_p \in W_1, W_2$.

PROOF: 0_p is expressible as a polynomial with all coefficients $a_i = 0$.

Then $a_i = 0$ when i even, so $0_p \in W_1$, and $a_i = 0$ when i odd, so $0_p \in W_2$.

SET 2

1.3 # 25, cont'd.

Jonathan Lam
Prof. M. Mitchell
MA326
Lin. Alg.
7/19/19

CLAIM: $\forall w, x \in W_1, w+x \in W_1$,

$\forall y, z \in W_2, y+z \in W_2$

PROOF: If $w = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 \in W_1$,

$x = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0 \in W_1$,

$w+x = (a_j + b_j) x^j + (a_{j+1} + b_{j+1}) x^{j+1} + \dots + (a_0 + b_0)$,

$\forall i \text{ even}, (a_i + b_i) = 0 + 0 = 0, \therefore w+x \in W_1$.

$\forall y = c_n x^n + c_{n-1} x^{n-1} + \dots + c_1 x + c_0 \in W_2$,

$\forall z = d_m x^m + d_{m-1} x^{m-1} + \dots + d_1 x + d_0 \in W_2$,

$y+z = (c_j + d_j) x^j + (c_{j-1} + d_{j-1}) x^{j+1} + \dots + (c_0 + d_0)$,

$\forall i \text{ odd}, (c_i + d_i) = 0 + 0 = 0, \therefore y+z \in W_2$.

CLAIM: $\forall c \in F, \forall x \in W_1, \forall y \in W_2, cx \in W_1, cy \in W_2$.

PROOF: If $x = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 \in W_1$,

then $cx = (ca_n) x^n + (ca_{n-1}) x^{n-1} + \dots + (ca_1) x + (ca_0)$.

$\forall i \text{ even}, ca_i = c(0) = 0, \therefore cx \in W_1$.

$\forall y = b_m x^m + b_{m-1} x^{m-1} + \dots + b_1 x + b_0 \in W_2$,

then $cy = (cb_m) x^m + (cb_{m-1}) x^{m-1} + \dots + (cb_1) x + (cb_0)$.

$\forall i \text{ odd}, cb_i = c(0) = 0, \therefore cy \in W_2$.

By (Thm 1.3), W_1, W_2 subsp P(F).

25. 2) CLAIM: $W_1 \cap W_2 = \{0\}$

PROOF: Since W_1, W_2 subsp P(F), $0_p \in W_1, W_2$.

Next, choose $x = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 \neq 0_p \in P(F)$.

Then, $\forall i$ s.t. $a_i \neq 0$, i must be odd or even.

If i even, $x \notin W_2$. If i odd, $x \notin W_1$.

Thus, $\forall x \neq 0_p, x \notin W_1 \cap W_2$.

$\therefore W_1 \cap W_2 = \{0_p\}$.

25.3) Show $w_1 + w_2 = V$.

CLAIM: $w_1 + w_2 \subseteq P(F)$

PROOF: $w_1 + w_2 = \{x + y : x \in w_1, y \in w_2\}$. Since $w_1, w_2 \subseteq V$,
 $\forall x \in w_1 \subseteq P(F)$ & $y \in w_2 \subseteq P(F)$, thus,
 $w_1 + w_2 \subseteq P(F)$.

CLAIM: $P(F) \subseteq w_1 + w_2$

PROOF: $\forall x \in P(F)$, x can be represented as (for some odd n)

$$\begin{aligned} x &= a_n x^n + b_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_3 x^3 + b_2 x^2 + a_1 x + b_0 \\ &= (a_n + 0)x^n + (0 + b_{n-1})x^{n-1} + (a_{n-2} + 0)x^{n-2} + \dots \\ &\quad + (a_3 + 0)x^3 + (0 + b_2)x^2 + (a_1 + 0)x + (0 + b_0) \\ &= (a_n x^n + 0 x^{n-1} + a_{n-2} x^{n-2} + \dots + a_3 x^3 + 0 x^2 + a_1 x + 0) \\ &\quad + (0 x^n + b_{n-1} x^{n-1} + 0 x^{n-2} + \dots + 0 x^3 + b_2 x^2 + 0 x + b_0). \end{aligned}$$

Thus x can be expressed as $y + z$, $y \in w_1, z \in w_2$,
so $P(F) \subseteq w_1 + w_2$.

Since $w_1 + w_2 \subseteq P(F)$, $P(F) \subseteq w_1 + w_2$, so $w_1 + w_2 = P(F)$.

Since the three conditions for a direct sum are satisfied,

$$w_1 \oplus w_2 = P(F).$$

PSET 2

L4 #2ae, 4ab, 73, 17.

Solve the linear systems using Gaussian elimination.

Jonathan Lam
Prof. Mutchler
MA320,
Lin. Alg.
9/19/19

$$2x_1 - 2x_2 - 3x_3 = -2$$

$$3x_1 - 3x_2 - 2x_3 + 5x_4 = 7$$

$$x_1 - x_2 - 2x_3 - x_4 = -3$$

$$x_1 - x_2 - 2x_3 - x_4 = -3$$

$$0 + 0 + x_3 + 2x_4 = 4$$

$$0 + 0 + 4x_3 + 8x_4 = 16$$

$$x_1 - x_2 - 2x_3 - x_4 = -3$$

$$0 + 0 + x_3 + 2x_4 = 4$$

$$0 + 0 + \cancel{x_3} + 2x_4 = 4 \Rightarrow 0 + 0 + 0 + 0 = 0.$$

$$\text{Let } x_3 = t. \text{ Then } t + 2x_4 = 4 \Rightarrow x_4 = 2 - \frac{t}{2}.$$

$$\text{Let } x_1 = s. \text{ Then } 2s - 2x_2 - 3t = -2 \Rightarrow x_2 = 1 + s - \frac{3}{2}t.$$

$$\text{Then a solution is } (x_1, x_2, x_3, x_4) = (s, 1 + s - \frac{3}{2}t, t, 2 - \frac{t}{2}).$$

$$\forall t, s \in \mathbb{R}.$$

$$x_1 + 2x_2 - 4x_3 - x_4 + x_5 = 7$$

$$-x_1 + 10x_3 - 3x_4 - 4x_5 = -16$$

$$2x_1 + 5x_2 - 5x_3 - 4x_4 - x_5 = 2$$

$$4x_1 + 11x_2 - 7x_3 - 10x_4 - 2x_5 = 7.$$

$$x_1 + 2x_2 - 4x_3 - x_4 + x_5 = 7$$

$$2x_2 - 6x_3 - 4x_4 - 3x_5 = -9$$

$$x_2 + 3x_3 - 2x_4 - 3x_5 = -12$$

$$3x_2 + 9x_3 - 6x_4 - 6x_5 = -21$$

$$x_1 + 2x_2 - 4x_3 - x_4 + x_5 = 7$$

$$x_2 + 3x_3 - 2x_4 - 3x_5 = -12$$

$$6x_3 - 4x_4 - 3x_5 = -9$$

$$3x_5 = 15 \Rightarrow x_5 = 5$$

Let $x_3 = t$. Then $6t - 4x_4 - 3(5) = -9$

$$\Rightarrow x_4 = \frac{6t + 9 - 15}{4} = \frac{3}{2}t - \frac{3}{2}$$

$$x_2 + 3t - 2\left(\frac{3}{2}t - \frac{3}{2}\right) - 3(5) = -12$$

$$x_2 = -12 + 18 - 3t + 8t = 3t = 0$$

$$x_1 + 2t - 4t - \left(\frac{3}{2}t - \frac{3}{2}\right) + 5 = 7$$

$$\begin{aligned}x_1 &= 7 + 4t + \frac{3}{2}t - \frac{3}{2} - 5 \\&= \frac{1}{2}t + \frac{11}{2}\end{aligned}$$

$$\text{Thus a solution is } (x_1, x_2, x_3, x_4, x_5) = \left(\frac{1}{2}t + \frac{11}{2}, 0, \frac{3}{2}t - \frac{3}{2}, t, 5\right)$$

$\forall t \in \mathbb{R}$.

4. Determine if first polynomial can be expressed as a lin. comb. of other two.

$$(1) a. x^3 - 3x + 5 = a(x^3 + 2x^2 - x + 1) + b(x^3 + 3x^2 - 1)$$

$$\begin{cases} a+b = 1 \\ 2a+3b = 0 \\ -a = -3 \Rightarrow a = 3, a+b = 1 \Rightarrow b = 1-3 = -2 \\ a-b = 5. \end{cases}$$

Check for consistency:

$$3+(-2)=1, 2(3)+3(-2)=0, -3=-3, 3-(-2)=5. \checkmark$$

Yes. (solution: $(a, b) = (1, -2)$)

$$(2) 4x^3 + 2x^2 - 6 = a(x^3 - 2x^2 + 4x + 1) + b(3x^3 - 6x^2 + x + 4)$$

$$\begin{cases} a+3b=4 \\ -2a-6b=2 \quad \text{multply by -2} \\ 4a+b=0 \end{cases}$$

$$a+4b=-6 \quad \text{subtract: } b=-10,$$

$$4a+(-10)=0 \Rightarrow a=\frac{5}{2}.$$

$$a+4b=\frac{5}{2}+4(-10)=\frac{5}{2}-40=\frac{-75}{2} \neq -6.$$

This system is not consistent, thus the first polynomial cannot be expressed as a linear. comb. of the other two.

PSET 2

1.4 # 13, 17.

Jonathan Lam
Prof. Mutchler
MA 326
Lin. Alg.
9/19/19

13.

(CLAIM) Show that if $S_1, S_2 \subseteq \text{v.s. } V$ s.t. $S_1 \subseteq S_2$,

then $\text{span}(S_1) \subseteq \text{span}(S_2)$. In particular, if $S_1 \subseteq S_2$ and $\text{span}(S_1) = V$, deduce that $\text{span}(S_2) = V$.

CLAIM: If $S_1, S_2 \subseteq \text{v.s. } V$, $S_1 \subseteq S_2$, then $\text{span}(S_1) \subseteq \text{span}(S_2)$

PROOF: For any linear combination $x \in \text{span}(S_1)$,

$$x = a_1x_1 + a_2x_2 + \dots + a_nx_n, \text{ then } a_1, a_2, \dots, a_n \in F,$$

$x_1, x_2, \dots, x_n \in S_1 \subseteq S_2$. Since all $x_i \in S_2$, this

is also a linear combination in S_2 , so $\text{span}(S_1) \subseteq \text{span}(S_2)$.

CLAIM: If $S_1 \subseteq S_2$, $\text{span}(S_1) = V$, then $\text{span}(S_2) = V$.

PROOF: By the above proof, $\text{span}(S_1) = V \subseteq \text{span}(S_2)$

By (THM 1.5), $\text{span}(S_2) \subseteq V$. Therefore, $\text{span}(S_2) = V$.

17

Let w subsp v.s. V . Under what conditions are there only a finite # of distinct subsets S of w s.t. S generates w .

CLAIM: If w finite, then there are only a finite # of distinct generating subsets.

PROOF: If w finite, then there are only a finite # of all subsets (finite power set), so there can only be a finite # of generating subsets.

LEM 2: If w is an infinite v.s., $\forall x \in w$, let $S = w \setminus \{x\}$.

Then, S is a generating set of w .

PROOF: Choose an $x \in w$.

(Case 1: $x = 0$) $\forall y \in S$, $0 \cdot y = 0$, so $x \in \text{span}(S)$.

(Case 2: $x \neq 0$) $\forall y \in S$, $\exists (x-y) \in S$, since $x, y \in w$,

$x-y \in w$, and $x-y \neq x \in S$. Then $x \in \text{span}(S)$.

Thus $x \in \text{span}(S)$. Also, $\forall z \in S$, $z = 1 \cdot z \Rightarrow z \in \text{span}(S)$.

Thus $w = S \cup \{x\} \subseteq \text{span}(S)$. By (THM 1.5) $\text{span}(S) \subseteq w$, thus $\text{span}(S) = w$ and S generates w .

9/19/19

CLAIM: If W infinite, there are infinitely many generating subsets of W .

PROOF: By (LEM 2), any subset $S = W \setminus \{x\}$, $\forall x \in W$ is a generating subset of W . Since W is infinite, there are an infinite # of generating subsets.

Since W finite $\Rightarrow W$ has finitely many generating subsets, and W infinite $\Rightarrow W$ has infinitely many generating subsets, this implies the tautology.

W finite $\Leftrightarrow W$ has a finite # of generating subsets.