
HG002 maternal 9p (train): telomere-adjacent block 15q-like, then 9p-like
HG002 carrier
ONT R10 ultra-long, 259 kb: conclusive
HG002 carrier
ONT R10 ultra-long, 258 kb: conclusive
HG002 carrier
ONT R10 duplex, 79 kb: conclusive
HG002 carrier
PacBio HiFi Revio, 35 kb: conclusive
native control NA18570
ONT R9.4.1 (1KG), 36 kb: non-informative
0
10
20
30
40
position on read from the telomere boundary (kb)
HG002 maternal 17p (train): telomere-adjacent block 11p-like, then 17p-like
HG002 carrier
ONT R10 ultra-long, 349 kb: conclusive
HG002 carrier
ONT R10 ultra-long, 261 kb: conclusive
HG002 carrier
ONT R10 duplex, 53 kb: conclusive
native control HG01786
ONT R9.4.1 (1KG), 90 kb: contradicts
0
10
20
30
40
position on read from the telomere boundary (kb)
HG002 paternal 8p (train): telomere-adjacent block 5q-like, then 11p-like
HG002 carrier
ONT R10 ultra-long, 401 kb: conclusive
HG002 carrier
ONT R10 ultra-long, 343 kb: conclusive
native control NA18940
ONT R9.4.1 (1KG), 31 kb: non-informative
0
20
40
60
80
100
position on read from the telomere boundary (kb)
donor (D) allele, SNV site
proximal (P) allele, SNV site
D-private 200 bp bin
P-private 200 bp bin
neither allele
read switch, 95% interval
catalogue switch
assembly window, same unit model (bar at top)