

a
Carriers of 18 of 34 forms are less diverse than native donor alleles
carrier diversity, π (SNVs/kb)
native donor-arm diversity, π (SNVs/kb)
equal diversity
16p>9q
8p>1p
3q>11p
6p>11p
20q>3q
1p>11p
consistent with single origin (13)
clustered, low diversity, no carrier-fixed SNVs (3)
clustered, carriers as diverse as natives (7)
not supported (11)
b
Carriers sit on one tight lineage; donor lineage survives in natives (20q>3q, 3q>11p) or not (6p>11p)
carriers
natives
other recipients
distance to carrier consensus (SNVs/kb, +0.02)
20q>3q
3q>11p
6p>11p
1p>11p
c
Median-based ages of 17 of 23 clustered forms fall in 0.2-1.5 Myr; mean-based ages are higher
8p>1p
9q>15q
16q>15q
19q>16q
19q>10q
20p>7p
9q>1p
6p>1p
11p>3q
20q>3q
3q>11p
6p>12p
6p>11p
4q>4p
9q>11p
20p>16q
Xq>15q
6p>6q
8p>6q
16p>16q
1p>11p
9q>20p (0; at 5)
19q>21q (0; at 5)
dots: median-based age
mean-based (ρ) age
clock age, kyr (μ = 1.25e-8 /bp/gen, 29 y/gen)
d
No recurrent human junction is present in a great-ape T2T genome
probe not found across switch
found, but native probes align equally (shared duplicon)
recurrent human junctions tested (n=70)
chimpanzee
bonobo
gorilla
Sumatran orangutan
Bornean orangutan