
a
Common forms are present in all superpopulations
but many differ in frequency
20q>3q * 0.72 0.36 0.18 0.58 0.34 0.21
8p>1p 0.40 0.46 0.39 0.43 0.46 -0.01
2q>16q * 0.20 0.39 0.52 0.72 0.42 0.13
3q>11p 0.46 0.56 0.37 0.71 0.48 0.03
6p>11p * 0.19 0.41 0.39 0.71 0.52 0.13
1p>6q 0.44 0.64 0.41 0.45 0.54 0.01
19p>11p * 0.46 0.23 0.08 0.36 0.24 0.11
19p>15q * 0.25 0.17 0.51 0.18 0.25 0.08
5q>6q * 0.53 0.11 0.16 0.12 0.11 0.22
4p>15p 0.39 0.40 0.32 0.35 0.44 -0.01
20p>7p * 0.17 0.21 0.37 0.16 0.30 0.03
4q>15p * 0.07 0.61 0.50 0.42 0.44 0.22
10p>18p 0.27 0.21 0.22 0.26 0.35 0.00
20p>16q * 0.33 0.12 0.19 0.20 0.16 0.03
17p>11p 0.31 0.42 0.47 0.29 0.14 0.03
11p>1p * 0.14 0.16 0.09 0.37 0.20 0.05
8p>6q 0.16 0.22 0.10 0.25 0.15 0.01
9q>11p 0.17 0.19 0.21 0.15 0.16 -0.01
5q>1p * 0.30 0.03 0.12 0.04 0.14 0.10
9p>3q 0.17 0.26 0.18 0.05 0.14 0.02
9q>20p 0.17 0.06 0.24 0.17 0.24 0.02
Xq>16p * 0.40 0.06 0.35 0.32 0.44 0.08
17q>16q 0.23 0.22 0.12 0.32 0.29 0.01
Xp>Yp 0.27 0.35 0.21 0.58 0.28 0.04
4q>10q * 0.36 0.03 0.10 0.29 0.11 0.12
19q>10q * 0.24 0.09 0.06 0.04 0.12 0.05
18p>10p 0.29 0.14 0.07 0.17 0.27 0.04
9p>11p 0.13 0.07 0.17 0.12 0.24 0.01
6q>16q 0.09 0.18 0.14 0.18 0.20 -0.00
12p>16p 0.07 0.16 0.16 0.06 0.05 0.02
AFR (138)
AMR (86)
EAS (102)
EUR (59)
SAS (70)
F_{ST} (5 superpopulations)
0.0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8
carrier frequency (haplotypes)
* superpopulation difference q < 0.05 (chemistry-stratified permutation)
b
Most large junction clusters span the
out-of-Africa split (below random expectation)
observed
random labelling
fraction of clusters in both continental
African and EUR/EAS/SAS samples
0.0
0.2
0.4
0.6
0.8
1.0
0.08
0.38
0.52
0.87
2 (n=66)
3-4 (n=55)
5-9 (n=31)
≥10 (n=39)
samples sharing the junction cluster