

- LRR per bin:  $\log_2$  of the bin's median depth over the member's autosomal median (GC- and panel-corrected where applicable)
- step fit of the track (exact total-variation denoising)
- BAF: alt-allele read fraction at the member's heterozygous sites (subsampled)
- phased fraction pooled over a window of sites (depth weighted  $\times 1/2$  with equal homologue copy,  $1/2 \pm d$  along an event with the sign giving the parent (child: below  $1/2$  the paternal homologue is in excess, above  $1/2$  the maternal; parent: below  $1/2$  the event lies on the untransmitted homologue, above  $1/2$  on the transmitted one); below where the window deviates in two or more members (paralogy, excluded from file and scan)
- child's maternal fraction at sites where the father is homozygous and the mother heterozygous; departs from the main track where the child carries two different maternal homologues (maternal meiotic trisomy or heterodisomy) and rejoins it after a crossover
- the same at sites where the mother is homozygous and the father heterozygous; departs where the child carries two different paternal homologues
- child's maternal copy number: the LRR step fit's copy number ( $2 \times 2^{\text{LRR}}$ ) times the maternal fraction's step fit
- child's paternal copy number, likewise
- called gain (line at the segment's mean LRR)
- called loss
- called copy-neutral event: loss of heterozygosity or uniparental heterodisomy (drawn at LRR 0)
- reference lines: LRR 0; BAF  $1/2$ ,  $1/3$  and  $2/3$ ; the centromere