

-  LRR per bin: $\log_2$ of the bin's trimmed-mean depth over the member's autosomal median (GC- and panel-corrected where applicable)
-  step fit of the track (exact total-variation denoising)
-  BAF: alt-allele read fraction at the member's heterozygous sites (subsampled)
-  informative site whose alt allele is maternal (father 0/0, mother 1/1)
-  informative site whose alt allele is paternal (father 1/1, mother 0/0)
-  phased fraction pooled over a window of sites (depth-weighted, $1/2$ with equal homologue copy, $1/2 \pm d$ along an event with the sign giving the parent (child: below $1/2$ the paternal homologue is in excess, above $1/2$ the maternal; parent: below $1/2$ the event lies on the untransmitted homologue, above $1/2$ on the transmitted one); hollow where the window deviates in two or more members (paralogy, excluded from fits and scan)
-  the same at sites where the mother is homozygous and the father heterozygous; departs from the main track where the child carries two different maternal homologues (maternal meiotic trisomy or heterodisomy) and rejoins it after a crossover
-  carries two different paternal homologues
-  parent's transmitted-allele fraction at sites where the child is heterozygous and the other parent homozygous; departs from the main track where the child carries two different homologues of this parent
-  child's maternal copy number: the LRR step fit's copy number ($2 \times 2^{\text{LRR}}$) times the maternal fraction's step fit
-  child's paternal copy number, likewise
-  parent's copy number of the transmitted homologue, likewise
-  parent's copy number of the untransmitted homologue
-  called gain (line at the segment's mean LRR)
-  called loss
-  called copy-neutral event: loss of heterozygosity or uniparental heterodisomy (drawn at LRR 0)
-  event supplied with --events (bracket above the track)
-  reference lines: LRR 0; BAF $1/2$, $1/3$ and $2/3$; the centromere