
LRR per bin: log2 of the bin's trimmed-mean depth over the member's autosomal median (GC- and panel-corrected where applicable)
step fit of the track (exact total-variation denoising)
BAF: alt-allele read fraction at the member's heterozygous sites (subsampled)
phased fraction pooled over a window of sites (depth-weighted): 1/2 with equal homologue copy, 1/2 +/- d along an event with the sign giving the parent (child: below 1/2 the paternal homologue is in excess, above 1/2 the maternal; parent: below 1/2 the event lies on the untransmitted homologue, above 1/2 on the transmitted one); hollow where the window deviates in two or more members (paralogy, excluded from fits and scan)
phased fraction at one site: maternal allele (child) or transmitted allele (parent), reference bias removed
child's maternal fraction at sites where the father is homozygous and the mother heterozygous; departs from the main track where the child carries two different maternal homologues (maternal meiotic trisomy or heterodisomy) and rejoins it after a crossover
the same at sites where the mother is homozygous and the father heterozygous; departs where the child carries two different paternal homologues
child's maternal copy number: the LRR step fit's copy number (2 x 2^LRR) times the maternal fraction's step fit
child's paternal copy number, likewise
called gain (line at the segment's mean LRR)
called loss
called copy-neutral event: loss of heterozygosity or uniparental heterodisomy (drawn at LRR 0)
reference lines: LRR 0; BAF 1/2, 1/3 and 2/3; the centromere