

- LRR per bin: log2 of the bin's trimmed-mean depth over the member's autosomal median (GC- and panel-corrected where applicable)
- step fit of the track (exact total-variation denoising)
- BAF: alt-allele read fraction at the member's heterozygous sites (subsampled)
- phased fraction pooled over a window of sites (depth-weighted), 1/2 with equal homologue copy, 1/2 + d along an event with the sign giving the parent (child: below 1/2 the paternal homologue is in excess, above 1/2 the maternal; parent: below 1/2 the event lies on the untransmitted homologue, above 1/2 on the transmitted one); below where the window deviates in two or more members (paralogy, excluded from file and scan)
- child's maternal fraction at sites where the father is homozygous and the mother heterozygous; departs from the main track where the child carries two different maternal homologues (maternal meiotic trisomy or heterodisomy) and rejoins it after a crossover
- the same at sites where the mother is homozygous and the father heterozygous; departs where the child carries two different paternal homologues
- child's maternal copy number: the LRR step fit's copy number ($2 \times 2^{\text{LRR}}$) times the maternal fraction's step fit
- child's paternal copy number, likewise
- called gain (line at the segment's mean LRR)
- called loss
- called copy-neutral event: loss of heterozygosity or uniparental heterodisomy (drawn at LRR 0)
- reference lines: LRR 0; BAF 1/2, 1/3 and 2/3; the centromere