

# Weight-adjusted Bernstein-Bezier Discontinuous Galerkin methods

Jesse Chan

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Bernstein-Bezier bases</b>	<b>1</b>
<b>3</b>	<b>Fast <math>L^2</math> projections with Bernstein polynomials</b>	<b>1</b>
3.1	Collapsed-coordinate quadratures . . . . .	1
3.2	Polynomial multiplication . . . . .	2
3.3	Inversion of modally diagonal matrices . . . . .	2
<b>4</b>	<b>Numerical experiments</b>	<b>3</b>
4.1	Expansion kernels . . . . .	3
4.1.1	Polynomial multiplication . . . . .	3
4.1.2	Collapsed-coordinate quadrature . . . . .	3
4.2	Mass matrix inversion kernel . . . . .	3
4.3	Monolithic kernel . . . . .	3
<b>5</b>	<b>Application to Weight-adjusted Discontinuous Galerkin (WADG) methods</b>	<b>3</b>

## Abstract

Alternative formula for mass inversion. GPU-accelerated versions of Ainsworth and Kirby Duffy transforms. Comparison to polynomial multiplication-based approach.

## 1 Introduction

[1]

## 2 Bernstein-Bezier bases

## 3 Fast $L^2$ projections with Bernstein polynomials

Involves three steps - evaluation in an enriched representation (either at quadrature points or in a higher degree polynomial basis), scaling by the weight, and projection down to polynomials of degree  $N$ .

### 3.1 Collapsed-coordinate quadratures

Fast evaluation at Duffy points [2, 3, 4].

Component-wise scaling by weight evaluated at collapsed-coordinate quadrature points.

### 3.2 Polynomial multiplication

For non-constant coefficients, DG requires being able to deal with polynomial multiplication and projection onto lower-dimensional subspaces. Multiplying polynomials together may be done using a discrete convolution and polynomial multiplication (J. Sanchez-Reyes 2003). The projection operator may be derived by noting that degree elevation operators are diagonal when transformed to a modal basis.

Rescaling by binomial coefficients results in the unscaled Bernstein basis. Polynomial multiplication is then equivalent to discrete convolution of the scaled binomial coefficients.

Quadrature-free strategy for nonlinear volume terms: polynomial multiplication + projection.

1. Polynomial multiplication of two BB basis functions representable as coefficient scaling,  $N_p$  scalar multiplications and storage of  $N_p$  coeffs, and another coefficient scaling.
2. To reduce local memory costs, process coeffs for  $fg$  over one or more  $(d - 1)$  dimensional layers.
3. Store ids and load a triangular number of loads.

### 3.3 Inversion of modally diagonal matrices

The inverse of any modally diagonal matrix  $\mathbf{D}_N^{-1}$  can be represented in the form

$$\mathbf{D}_N^{-1} = \sum_{j=0}^N c_j \mathbf{E}_{N-j}^N (\mathbf{E}_{N-j}^M)^T.$$

This property was shown for the polynomial projection matrix by Waldron in [5, 6]. We give an alternative proof of this below where  $\mathbf{M}$  is any modally diagonal matrix, including rectangular matrices.

The constants  $c_j$  may be computed through the solution of an  $(N + 1) \times (N + 1)$  matrix system, using the fact that upon transformation to a modal basis,  $\mathbf{E}_{N-j}^N$  is a diagonal matrix of ones and zeros, while  $\mathbf{E}_{N-j}^M$  is a diagonal matrix with entries

$$\frac{\lambda_i^{N-j}}{\lambda_i^M}, \quad i = 0, \dots, N.$$

This may be factored into an application of  $\mathbf{E}_N^M$ , then an application of

$$\begin{aligned} \sum_{j=0}^N c_j \mathbf{E}_{N-j}^N (\mathbf{E}_{N-j}^N)^T &= c_0 \mathbf{I} + c_1 \mathbf{E}_{N-1}^N (\mathbf{E}_{N-1}^N)^T + c_2 \mathbf{E}_{N-1}^N \mathbf{E}_{N-2}^{N-1} (\mathbf{E}_{N-2}^{N-1})^T (\mathbf{E}_{N-1}^N)^T + \dots \\ &= c_0 \mathbf{I} + c_1 \mathbf{E}_{N-1}^N \left( \mathbf{I} + \frac{c_2}{c_1} \mathbf{E}_{N-2}^{N-1} (\mathbf{I} + \dots) (\mathbf{E}_{N-2}^{N-1})^T \right) (\mathbf{E}_{N-1}^N)^T. \end{aligned}$$

This may be applied in two sweeps of length  $N$ , using in-place updates to memory. Unfortunately, for shared-memory parallelization, this will require synchronizations between each application of each matrix.

The cost of applying  $(\mathbf{E}_N^M)^T$  is the application of  $(M - N)$  sparse degree elevation operations, each of which is  $O(M^d)$  cost. Assuming  $M \approx N$  (it is reasonable to match the order of the data with the order of approximation), this gives  $O(N^{d+1})$  cost.

When applying the projection operator, since each operation is  $O(N^3)$  and we apply  $O(N)$  total operations, we have an  $O(N^{d+1})$  overall cost.

This can also be used to apply the inverse mass matrix since it is diagonal under the transformation  $T$ .

## 4 Numerical experiments

### 4.1 Expansion kernels

#### 4.1.1 Polynomial multiplication

#### 4.1.2 Collapsed-coordinate quadrature

### 4.2 Mass matrix inversion kernel

## 5 Application to Weight-adjusted Discontinuous Galerkin (WADG) methods

Numerical experiment: time to solution for high orders using both NDG-WADG and BB-WADG.

[7, 8]

## References

- [1] Jesse Chan and T Warburton. GPU-accelerated Bernstein-Bezier discontinuous Galerkin methods for wave problems. *arXiv preprint arXiv:1512.06025*, 2015.
- [2] Mark Ainsworth, Miangaly Gaelle Andriamaro, and Oleg Davydov. Bernstein-Bézier finite elements of arbitrary order and optimal assembly procedures. *SIAM Journal on Scientific Computing*, 33(6):3087–3109, 2011.
- [3] Robert C Kirby. Fast simplicial finite element algorithms using Bernstein polynomials. *Numerische Mathematik*, 117(4):631–652, 2011.
- [4] Robert C Kirby and Kieu Tri Thinh. Fast simplicial quadrature-based finite element operators using Bernstein polynomials. *Numerische Mathematik*, 121(2):261–279, 2012.
- [5] Shayne Waldron. On the Bernstein-Bézier form of Jacobi polynomials on a simplex. *Journal of Approximation Theory*, 140(1):86–99, 2006.
- [6] Shayne Waldron. Computing orthogonal polynomials on a triangle by degree raising. *Numerical Algorithms*, 42(2):171–179, 2006.
- [7] Jesse Chan, Russell J Hewett, and T Warburton. Weight-adjusted discontinuous Galerkin methods: wave propagation in heterogeneous media. *arXiv preprint arXiv:1608.01944*, 2016.
- [8] Jesse Chan, Russell J Hewett, and T Warburton. Weight-adjusted discontinuous Galerkin methods: curvilinear meshes. *arXiv preprint arXiv:1608.03836*, 2016.