

SKEW-SYMMETRIC ENTROPY STABLE DISCONTINUOUS GALERKIN FORMULATIONS

JESSE CHAN

Abstract. Entropy stable high order methods for nonlinear conservation laws satisfy an inherent discrete entropy inequality. The construction of such schemes has relied on the use of carefully chosen collocation points [1, 2, 3, 4] or volume and surface quadrature rules [5, 6] to produce operators which satisfy a summation-by-parts (SBP) property. In this work, we show how to construct skew-symmetric schemes which are entropy stable even for volume and surface quadratures under which an SBP property does not hold. These skew-symmetric formulations avoid the use of a “strong” SBP property, and require only that operators exactly differentiate constants and satisfy a discrete form of the fundamental theorem of calculus. We conclude by discussing mortar-based implementations of skew-symmetric formulations for entropy stable schemes on hybrid meshes.

1. Introduction. High order methods for the simulation of time-dependent compressible flow have the potential to achieve higher levels of accuracy at lower costs compared to current low order schemes [7]. In addition to superior accuracy, the low numerical dispersion and dissipation of high order methods [8] enables the accurate propagation of waves over long distances and time scales. The same properties also make high order methods attractive for unsteady phenomena such as vorticular and turbulent flows, which are sensitive to numerical dissipation [9, 7].

However, when applied to nonlinear conservation laws, high order methods can experience artificial growth and blow-up near under-resolved features such as shocks, turbulence, or boundary layers. In practice, the application of high order methods to practical problems requires shock capturing and stabilization techniques (such as artificial viscosity) or solution regularization (such as filtering or limiting) to prevent solution blow-up. The resulting schemes for nonlinear conservation laws walk a fine line between stability, robustness, and accuracy. Aggressive stabilization or regularization can result in the loss of high order accuracy, while too little can result in instability [7]. Moreover, it can be difficult to determine robust expressions for stabilization parameters, as parameters which work for one simulation can fail when applied to a different physical regime or discretization setting.

These issues have motivated the introduction of high order *entropy stable* discretizations, which satisfy a semi-discrete entropy inequality while maintaining high order accuracy in smooth regions. Proofs of continuous entropy inequalities rely on the chain rule, which does not hold discretely due to effects such as quadrature error. By using an approach referred to as “flux differencing”, entropy stable schemes account for the loss of the chain rule while maintaining a semi-discrete analogue of the continuous entropy inequality. These schemes were first introduced as high order collocation methods on tensor product elements in [2, 3, 10, 11], and were extended to simplicial elements in [12, 13, 14, 5, 6]. Entropy stable methods have also been extended to a variety of other discretization settings, including staggered grids [15], Gauss-Legendre collocation [4], and non-conforming meshes [16].

Entropy stable discretizations are built upon flux differencing and a summation-by-parts (SBP) property, which serves as a discrete analogue of integration by parts for quadrature-based discretization matrices. However, the SBP property does not hold for some under-integrated quadrature rules, which naturally arise in certain discretization settings. For example, on hybrid meshes consisting of both quadrilateral and triangular elements, it is convenient to utilize the same quadrature rule on shared faces between different element types. On degree N tensor product elements, a pop-

ular choice of quadrature is an $(N + 1)$ -point Gauss-Legendre-Lobatto (GLL) rule. When both volume and surface integrals are approximated using $(N + 1)$ point GLL quadrature rules, the SBP property holds, despite the fact that GLL quadrature is inexact for the integrands which appear in finite element formulations [2]. However, while GLL quadrature induces an SBP property on quadrilateral elements, but does not always result in an SBP property if used for triangular elements [5].

This work proposes an alternative formulation which utilizes a skew-symmetric form of flux differencing. Under such a formulation, the proof of entropy stability requires only a weaker “variational” form of the SBP property, which holds under a more general class of quadrature rules compared to the decoupled SBP property introduced in [5, 6]. We show that the skew-symmetric formulation is entropy stable, locally conservative, and free-stream preserving on curved elements, and confirm theoretical results with numerical experiments on triangular, quadrilateral, and hybrid triangular-quadrilateral meshes.

It should be noted that a similar approach to entropy stable discretizations was introduced within a finite difference framework [13, 14] using multidimensional differencing operators which are constructed to satisfy accuracy conditions and an SBP property [23]. These operators do not correspond to any specific basis or approximation space, but can be shown to exist for nodal points corresponding to sufficiently accurate choices of volume and surface quadrature. The formulations presented in this work require similar accuracy conditions, but differ from those of [13, 14] in that they build upon an explicit approximation space. In contrast with traditional SBP coupling terms [12], the use of decoupled SBP operators also accomodates general choices of volume and surface quadrature (e.g. volume quadratures without boundary nodes and over-integrated rules) while ensuring compact coupling conditions between neighboring elements.

The structure of the paper is as follows: Section 2 describes the continuous entropy inequality which we aim to replicate discretely. Section 3 and Section 4 introduce polynomial approximation spaces and quadrature-based SBP operators on simplicial and tensor product elements. Section 5 then introduces the variational SBP property and an entropy conservative skew-symmetric formulation. Section 6 extends the skew-symmetric formulation to curved elements, and Section 8 presents numerical experiments which verify the theoretical assumptions, stability, and accuracy of the proposed formulations.

2. Entropy stability for systems of nonlinear conservation laws. We begin by reviewing the dissipation of entropy for a d -dimensional system of nonlinear conservation laws on a domain Ω

$$(1) \quad \frac{\partial \mathbf{u}_h}{\partial t} + \sum_{j=1}^d \frac{\partial \mathbf{f}_j(\mathbf{u})}{\partial x_j} = 0, \quad \mathbf{u} \in \mathbb{R}^n, \quad \mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n,$$

where \mathbf{u} are the conservative variables and $\mathbf{f}(\mathbf{u})$ is a vector-valued nonlinear flux function. We are interested in nonlinear conservation laws for which a convex entropy function $U(\mathbf{u})$ exists. For such systems, the *entropy variables* are an invertible mapping $\mathbf{v}(\mathbf{u}) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined as the derivative of the entropy function with respect to the conservative variables

$$(2) \quad \mathbf{v}(\mathbf{u}) = \frac{\partial U}{\partial \mathbf{u}}.$$

Several widely used equations in fluid modeling (Burgers, shallow water, compressible Euler and Navier-Stokes equations) yield convex entropy functions $U(\mathbf{u})$ [18, 13]. Let $\partial\Omega$ be the boundary of Ω with outward unit normal \mathbf{n} . By multiplying the equation (1) with $\mathbf{v}(\mathbf{u})^T$, the solutions \mathbf{u} of (1) can be shown to satisfy an entropy inequality

$$(3) \quad \int_{\Omega} \frac{\partial U(\mathbf{u})}{\partial t} dx + \int_{\partial\Omega} \sum_{j=1}^d (\mathbf{v}(\mathbf{u})^T \mathbf{f}_j(\mathbf{u}) - \psi_j(\mathbf{v}(\mathbf{u}))) n_j dx \leq 0,$$

where $\mathbf{n} = (n_1, \dots, n_d)$ denotes the outward unit normal, and $\psi_j(\mathbf{u})$ is some function referred to as the entropy potential.

The proof of (3) requires the use of the chain rule [19, 20, 21]. The instability-in-practice of high order schemes for (1) can be attributed in part to the fact that the discrete form of the equations do not satisfy the chain rule, and thus do not satisfy (3). As a result, discretizations of (1) do not typically possess an underlying statement of stability. For low order schemes, this can be offset in practice by the inherent numerical dissipation. However, because high order discretizations possess low numerical dissipation, the lack of an underlying discrete stability has been conflated with the idea that high order methods are inherently less stable than low order methods.

3. Polynomial approximation spaces. In this work, we consider either simplicial reference elements (triangles and tetrahedra) or tensor product reference elements (quadrilaterals and hexahedra). We define an approximation space using degree N polynomials on the reference element; however, the natural polynomial approximation space differs depending on the element type [22].

On a d -dimensional reference simplex, the natural polynomial space are total degree N polynomials

$$P^N(\widehat{D}) = \left\{ \widehat{x}_1^{i_1} \dots \widehat{x}_d^{i_d}, \quad \widehat{\mathbf{x}} \in \widehat{D}, \quad 0 \leq \sum_{k=1}^d i_k \leq N \right\}.$$

In contrast, the natural polynomial space on a d -dimensional tensor product element is the space of maximum degree N polynomials

$$Q^N(\widehat{D}) = \left\{ \widehat{x}_1^{i_1} \dots \widehat{x}_d^{i_d}, \quad \widehat{\mathbf{x}} \in \widehat{D}, \quad 0 \leq i_k \leq N, \quad k = 1, \dots, d \right\}.$$

We denote the natural approximation space on a given reference element \widehat{D} by V^N . Furthermore, we denote the dimension of V^N as $N_p = \dim(V^N(\widehat{D}))$.

The proofs presented in this work will also use anisotropic tensor product polynomial spaces, where the maximum polynomial degree varies depending on the coordinate direction. We denote such spaces by Q^{N_1, \dots, N_d} , where N_k are non-negative integers and

$$Q^{N_1, N_2, \dots, N_d}(\widehat{D}) = \left\{ \widehat{x}_1^{i_1} \dots \widehat{x}_d^{i_d}, \quad \widehat{\mathbf{x}} \in \widehat{D}, \quad 0 \leq i_k \leq N_k, \quad k = 1, \dots, d \right\}.$$

For example, the isotropic tensor product space Q^N is the same as $Q^{N, \dots, N}$.

We also define trace spaces for each reference element. Let \widehat{f} be a face of the reference element \widehat{D} . The trace space $V^N(\widehat{f})$ is defined as the restrictions of functions in $V^N(\widehat{D})$ to \widehat{f} , and denote the dimension of the trace space as $\dim(V^N(\widehat{f})) = N_p^f$.

$$V^N(\widehat{f}) = \left\{ u|_{\widehat{f}}, \quad u \in V^N(\widehat{D}), \quad \widehat{f} \in \partial\widehat{D} \right\}.$$

For example, on a d -dimensional simplex, $V^N(\partial\hat{D})$ consists of total degree N polynomials on simplices of dimension $(d-1)$. On a d -dimensional tensor product element, $V^N(\partial\hat{D})$ consists of maximum degree N polynomials on a tensor product element of dimension $(d-1)$.

4. Quadrature-based matrices and decoupled SBP operators. Let $\hat{D} \subset \mathbb{R}^d$ denote a reference element with surface $\partial\hat{D}$. The high order schemes in [5, 6] begin by approximating the solution in a degree N polynomial basis $\{\phi_j(\mathbf{x})\}_{j=1}^{N_p}$ on \hat{D} . These schemes also assume volume and surface quadrature rules (\mathbf{x}_i, w_i) , (\mathbf{x}_i^f, w_i^f) on \hat{D} . We will specify the accuracy of each quadrature rule later, and discuss how quadrature accuracy implies specific summation-by-parts properties.

Let $\mathbf{V}_q, \mathbf{V}_f$ denote interpolation matrices, and let \mathbf{D}^i be the differentiation matrix with respect to the i th coordinate such that

$$(4) \quad (\mathbf{V}_q)_{ij} = \phi_j(\mathbf{x}_i), \quad (\mathbf{V}_f)_{ij} = \phi_j(\mathbf{x}_i^f), \quad \frac{\partial \phi_j(\mathbf{x})}{\partial x_i} = \sum_{k=1}^{N_p} (\mathbf{D}_{jk}^i) \phi_k(\mathbf{x}).$$

The interpolation matrices $\mathbf{V}_q, \mathbf{V}_f$ map basis coefficients to evaluations at volume and surface quadrature points respectively, while the differentiation matrix \mathbf{D}_i maps basis coefficients of a function to the basis coefficients of its derivative with respect to x_k . The interpolation matrices are used to assemble the mass matrix \mathbf{M} , the quadrature-based projection matrix \mathbf{P}_q , and lifting matrix \mathbf{L}_f

$$(5) \quad \mathbf{M} = \mathbf{V}_q^T \mathbf{W} \mathbf{V}_q, \quad \mathbf{P}_q = \mathbf{M}^{-1} \mathbf{V}_q^T \mathbf{W}, \quad \mathbf{L}_f = \mathbf{M}^{-1} \mathbf{V}_f^T \mathbf{W}_f,$$

where \mathbf{W}, \mathbf{W}_f are diagonal matrices of volume and surface quadrature weights, respectively. The matrix \mathbf{P}_q is a quadrature-based discretization of the L^2 projection operator Π_N onto degree N polynomials.

Interpolation, differentiation, and L^2 projection matrices can be combined to construct finite difference operators. For example, the matrix $\mathbf{D}_q^i = \mathbf{V}_q \mathbf{D}^i \mathbf{P}_q$ maps function values at quadrature points to approximate values of the derivative at quadrature points. By choosing specific quadrature rules, \mathbf{D}_q^i recovers high order summation-by-parts finite difference operators in [1, 24, 25] and certain operators in [23]. However, to address difficulties in designing efficient entropy stable interface terms for nonlinear conservation laws, the PI introduced in [5] a new “decoupled” summation by parts matrix which builds interface terms directly into the approximation of the derivative.

Let $\hat{\mathbf{n}}$ denote the scaled outward normal vector $\hat{\mathbf{n}} = \{\hat{n}_1 \hat{J}_f, \dots, \hat{n}_d \hat{J}_f\}$, where \hat{J}_f is the determinant of the Jacobian of the mapping of a face of $\partial\hat{D}$ to a reference face. Let $\hat{\mathbf{n}}_i$ denote the vector containing values of the i th component $\hat{n}_i \hat{J}_f$ at all surface quadrature points, and let $\mathbf{Q}^i = \mathbf{W} \mathbf{D}_q^i$. The “decoupled” summation by parts operator \mathbf{D}_N^i is defined as the block matrix involving both volume and surface quadratures

$$(6) \quad \mathbf{D}_N^i = \mathbf{W}_N^{-1} \mathbf{Q}_N^i, \quad \mathbf{W}_N = \begin{pmatrix} \mathbf{W} & \\ & \mathbf{W}_f \end{pmatrix},$$

$$\mathbf{E} = \mathbf{V}_f \mathbf{P}_q, \quad \mathbf{B}^i = \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i), \quad \mathbf{Q}_N^i = \begin{bmatrix} \mathbf{Q}^i - \frac{1}{2} \mathbf{E}^T \mathbf{B}^i \mathbf{E} & \frac{1}{2} \mathbf{E}^T \mathbf{B}^i \\ -\frac{1}{2} \mathbf{B}^i \mathbf{E} & \frac{1}{2} \mathbf{B}^i \end{bmatrix}.$$

Here, \mathbf{B}^i is a boundary matrix, and \mathbf{E} denotes the extrapolation matrix which maps values at volume quadrature points to values at surface quadrature points using quadrature-based L^2 projection and polynomial interpolation.

It was shown in [5, 4] that, when combined with projection and lifting matrices, \mathbf{D}_N^i produces a high order polynomial approximation of $f \frac{\partial g}{\partial x_i}$. Let f, g be differentiable functions, and let $\mathbf{f}_i = f(\mathbf{x}_i)$, $\mathbf{g}_i = g(\mathbf{x}_i)$ denote values of f, g at volume and surface quadrature points. Then,

$$(7) \quad f \frac{\partial g}{\partial x_i} \approx \begin{bmatrix} \mathbf{P}_q & \mathbf{L}_f \end{bmatrix} \text{diag}(\mathbf{f}) \mathbf{D}_N^i \mathbf{g}.$$

For sufficiently accurate quadrature rules, the matrix \mathbf{Q}_N^i also satisfies a decoupled summation-by-parts (SBP) property, which is used to prove semi-discrete entropy stability for nonlinear conservation laws.

LEMMA 1. *Assume that the volume and surface quadrature rules are sufficiently accurate such that the quantities*

$$\int_{\hat{D}} \frac{\partial u}{\partial x_i} v, \quad \int_{\partial \hat{D}} u v n_i$$

are integrated exactly for all $u, v \in V^N(\hat{D})$ and $i = 1, \dots, d$. Then, the decoupled SBP operator \mathbf{D}_N^i (6) satisfies a summation by parts property:

$$(8) \quad \mathbf{Q}_N^i + (\mathbf{Q}_N^i)^T = \mathbf{B}_N^i, \quad \mathbf{B}_N^i = \begin{pmatrix} \mathbf{0} & \\ & \mathbf{B}^i \end{pmatrix}.$$

Proof. The proof is a straightforward extension of Theorem 1 in [5] to polynomial approximation spaces on non-simplicial elements. \square

The assumptions of Lemma 1 are satisfied for sufficiently accurate volume and surface quadratures. For example, on simplicial elements, (8) holds if the volume quadrature is exact for polynomial integrands of total degree $(2N-1)$, and the surface integral is exact for degree $2N$ polynomials on each face. Tensor product elements require stricter conditions: (8) holds if both the volume and surface quadratures are exact for polynomial integrands of degree $2N$ in each coordinate, due to the fact that derivatives of $u \in Q^N$ are degree $(N-1)$ polynomials with respect to one coordinate and degree N with respect to others.

REMARK. *It is worth noting that the assumptions of Lemma 1 are sufficient but not necessary conditions. For example, suppose \hat{D} is a quadrilateral element, and that the volume quadrature is a tensor product of $(N+1)$ point one-dimensional Gauss-Legendre-Lobatto (GLL) rule. If the surface quadrature also taken to be an $(N+1)$ -point GLL rule over each face, then \mathbf{Q}^i satisfies a traditional SBP property, which can be used to prove the decoupled SBP property for \mathbf{Q}_N^i .*

5. Skew-symmetric entropy conservative formulations on a single element. While the SBP property has been used to derive entropy stable schemes, the strong SBP property (8) is difficult to enforce in certain discretization settings, such as hybrid and non-conforming meshes. This difficulty is a result of the choices of volume and surface quadrature which naturally arise in these settings. We first illustrate how specific pairings of volume and surface quadratures can result in the loss of the SBP property (8). We then propose a skew-symmetric formulation which is entropy conservative without explicitly requiring operators which satisfy the SBP property.




Fig. 1: Volume and surface quadrature pairs which do not satisfy the assumptions of Lemma 1, and thus do not possess the decoupled SBP property (8). Volume quadrature nodes are drawn as circles, while surface quadrature nodes are drawn as squares.

5.1. Loss of the SBP property. In this section, we give examples of specific pairings of volume and surface quadratures under which the decoupled SBP property does not hold (see Figure 1). We consider two dimensional reference elements \hat{D} with spatial coordinates x, y .

Quadrilateral elements (Figure 1a). We first consider a quadrilateral element \hat{D} with an $(N+1)$ point tensor product GLL volume quadrature and $(N+1)$ point Gauss quadrature on each face. Let $u, v \in Q^N$ denote two arbitrary degree N polynomials. The assumptions of Lemma 1 are that the volume quadrature exactly integrates $\int_{\hat{D}} \frac{\partial u}{\partial x} v$ and that the surface quadrature exactly integrates $\int_{\partial \hat{D}} uv n_i$ on \hat{D} . Because the $(N+1)$ -point Gauss rule is exact for polynomials of degree $2N+1$ and the product $uv \in P^{2N}$ on each face, the surface quadrature satisfies the assumptions of Lemma 1. However, the 1D GLL rule is only exact for polynomials of degree $(2N-1)$. The derivative $\frac{\partial u}{\partial x}$ is a polynomial of degree $(N-1)$ in x , but is degree N in y . Thus, $\frac{\partial u}{\partial x} v$ is a polynomial of degree $(2N-1)$ in x but degree $2N$ in y , and is not integrated exactly by the volume quadrature.

Triangular elements (Figure 1b). We next consider a triangular element \hat{D} , where the volume quadrature is exact for degree $2N$ polynomials [26] and an $(N+1)$ -point GLL quadrature on each face. Let $u, v \in P^N$ denote two arbitrary degree N polynomials. The derivative $\frac{\partial u}{\partial x} \in P^{(N-1)}$, and $\frac{\partial u}{\partial x} v \in P^{(2N-1)}$, so the volume quadrature satisfies the assumptions of Lemma 1. However, because the surface quadrature is exact only degree $(2N-1)$ polynomials and the trace of $uv \in P^{2N}$, the surface quadrature does not satisfy the assumptions of Lemma 1.

These specific pairings of volume and surface quadratures appear naturally for hybrid meshes consisting of DG-SEM quadrilateral elements (using GLL volume quadrature) and triangular elements, as shown in Figure 2. In Figure 2a, the surface quadrature is a $(N+1)$ point GLL rule, and results in a loss of the SBP property on the triangle. In Figure 2b, the surface quadrature is a $(N+1)$ point Gauss-Legendre rule, and results in a loss of the SBP property on the quadrilateral element. The goal of this work is to construct high order accurate discretizations which preserve entropy conservation for situations in which the decoupled SBP property (8) does not hold.




Fig. 2: Examples of interface couplings which do not result in a decoupled SBP property (8). Volume quadrature nodes are drawn as circles, while surface quadrature nodes are drawn as squares.

5.2. A variational SBP property. The property (8) (which we will refer to as the “strong” SBP property) relates the polynomial exactness of specific quadrature rules to algebraic properties of quadrature-based matrices. We will relax accuracy conditions on these quadrature rules by utilizing a weaker “variational” version of the SBP property variational, which relies on the exactness of volume and surface quadrature rules only for specific polynomials.

On tensor product elements, we restrict ourselves to isotropic volume quadrature rules which are constructed from tensor products of one-dimensional quadrature formulas. For the remainder of this work, the degree of the multi-dimensional quadrature rule on tensor product elements will refer to the degree of exactness of the one-dimensional rule. For example, we refer to the quadrature rule constructed through a tensor product of one-dimensional $(N + 1)$ -point GLL quadrature rules as a degree $(2N - 1)$ quadrature rule. This choice of quadrature is sufficient to guarantee that the mass matrix is positive definite [27].

Throughout the remainder of this work, we make the following assumptions:

ASSUMPTION 1. Let $v \in V^N$ denote a fixed polynomial. We assume that:

1. the mass matrix \mathbf{M} is positive definite under the volume quadrature rule,
2. the volume quadrature rule is exact for integrals of the form

$$\int_{\hat{D}} \frac{\partial u}{\partial \hat{x}_j} v \text{ for all } u \in V^N(\hat{D}), j = 1, \dots, d.$$

3. the surface quadrature rule is exact for integrals of the form

$$\int_{\partial \hat{D}} u v n_j \text{ for all } u \in V^N(\hat{D}), j = 1, \dots, d, \text{ and } f \in \partial \hat{D}.$$

The conditions of Assumption 1 are relatively standard within the SBP literature [23, 5, 14], though they have not previously depended on the specific choice of polynomial $v(\mathbf{x})$. The strong SBP property (8) can be derived if Assumption 1 is satisfied for all $v \in V^N$. In contrast, we will derive a weaker SBP property which assumes only that Assumption 1 holds for a few specific polynomials v . In Sections 5.3 and 6.1, specific polynomials v will be motivated by proofs of entropy stability, and we will present examples of volume and surface quadrature rules on simplicial and tensor product elements which satisfy Assumption 1 for these choices of v .

The conditions of Assumption 1 imply the following variational SBP property:

LEMMA 2. Suppose Assumption 1 holds, and let \mathbf{v}_q denote the vector of values of $v(\mathbf{x})$ of at volume quadrature points. Let $u(\mathbf{x})$ be an L^2 integrable function, and let

279 \mathbf{u}_q denote the values of $u(\mathbf{x})$ at quadrature points. Then,

$$280 \quad \mathbf{v}_q^T \mathbf{Q}_i \mathbf{u}_q = \mathbf{v}_q^T (\mathbf{E}^T \mathbf{B}^i \mathbf{E} - \mathbf{Q}_i^T) \mathbf{u}_q.$$

281 *Proof.* Recall that $\mathbf{Q}_i = \mathbf{W} \mathbf{V}_q \mathbf{D}^i \mathbf{P}_q$. The quadrature-based L^2 projection $\Pi_N u$
 282 is a polynomial of degree N , and is computed discretely as $\mathbf{P}_q \mathbf{u}_q$. By the exactness
 283 of the volume quadrature in Assumption 1,

$$284 \quad \mathbf{v}_q^T \mathbf{Q}_i \mathbf{u}_q = \mathbf{v}_q^T \mathbf{W} \mathbf{D}^i \mathbf{P}_q \mathbf{u}_q = \int_{\widehat{D}} \frac{\partial \Pi_N u}{\partial \widehat{x}_i} v = \int_{\partial \widehat{D}} (\Pi_N u) v \widehat{n}_i - \int_{\widehat{D}} (\Pi_N u) \frac{\partial v}{\partial \widehat{x}_i}.$$

286 By the exactness of the surface quadrature in Assumption 1,

$$\begin{aligned} 287 \quad & \int_{\partial \widehat{D}} (\Pi_N u) v \widehat{n}_i - \int_{\widehat{D}} (\Pi_N u) \frac{\partial v}{\partial \widehat{x}_i} = \\ 288 \quad & \mathbf{v}_f^T \mathbf{B}^i \mathbf{E} \mathbf{u}_q - (\mathbf{V}_q \mathbf{D}^i \mathbf{P}_q \mathbf{v}_q)^T \mathbf{W} \mathbf{V}_q \mathbf{P}_q \mathbf{u}_q \\ 289 \quad & = \mathbf{v}_q^T \mathbf{E}^T \mathbf{B}^i \mathbf{E} \mathbf{u}_q - \mathbf{v}_q^T \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{V}_q^T \mathbf{W} \mathbf{V}_q \mathbf{P}_q \mathbf{u}_q, \end{aligned}$$

291 where we have used that, since $v \in V^N$, $\mathbf{v}_f = \mathbf{V}_f \mathbf{P}_q \mathbf{v}_q = \mathbf{E} \mathbf{v}_q$. The proof is completed
 292 by noting that

$$293 \quad \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{V}_q^T \mathbf{W} \mathbf{V}_q \mathbf{P}_q = \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{M} \mathbf{P}_q = \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{V}_q^T \mathbf{W} = \mathbf{Q}_i^T. \quad \square$$

294 The variational SBP property in Lemma 2 can be used to prove a similar property
 295 for the decoupled SBP operator. This property, along with the exact differentiation
 296 of constants, is necessary for the proof of entropy stability.

297 **LEMMA 3.** Suppose Assumption 1 holds. Let \mathbf{D}_N^i be a decoupled SBP operator on
 298 the reference element \widehat{D} . Then, the following variational SBP property holds:

$$299 \quad \mathbf{v}^T \mathbf{Q}_N^i \mathbf{u} = \mathbf{v}^T \left(\mathbf{B}_N^i - (\mathbf{Q}_N^i)^T \right) \mathbf{u}.$$

300 where \mathbf{v}, \mathbf{u} denotes the values of v and u at both volume and surface quadrature points.

301 *Proof.* For convenience, let $\mathbf{u}_q, \mathbf{u}_f$ and $\mathbf{v}_q, \mathbf{v}_f$ denote evaluations of u and v at
 302 volume and surface points, such that

$$304 \quad \mathbf{u} = \begin{pmatrix} \mathbf{u}_q \\ \mathbf{u}_f \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} \mathbf{v}_q \\ \mathbf{v}_f \end{pmatrix}.$$

305 The proof of the variational summation by parts property uses the definition of \mathbf{Q}_N^i
 306 (6),

$$\begin{aligned} 307 \quad \mathbf{v}^T \mathbf{Q}_N^i \mathbf{u} &= \mathbf{v}_q^T \mathbf{Q}_i \mathbf{u}_q - \frac{1}{2} (\mathbf{E} \mathbf{v}_q)^T \mathbf{B}^i (\mathbf{E} \mathbf{u}_q) + \frac{1}{2} (\mathbf{E} \mathbf{v}_q)^T \mathbf{B}^i \mathbf{u}_f \\ 308 \quad & - \frac{1}{2} \mathbf{v}_f^T \mathbf{B}^i (\mathbf{E} \mathbf{u}_q) + \frac{1}{2} \mathbf{v}_f^T \mathbf{B}^i \mathbf{u}_f. \end{aligned}$$

310 Applying Lemma 2 then yields

$$\begin{aligned}
311 \quad \mathbf{v}^T \mathbf{Q}_N^i \mathbf{u} &= -\mathbf{v}_q^T \mathbf{Q}_i^T \mathbf{u}_q + \frac{1}{2} (\mathbf{E} \mathbf{v}_q)^T \mathbf{B}^i (\mathbf{E} \mathbf{u}_q) + \frac{1}{2} (\mathbf{E} \mathbf{v}_q)^T \mathbf{B}^i \mathbf{u}_f \\
312 \quad &\quad - \frac{1}{2} \mathbf{v}_f^T \mathbf{B}^i (\mathbf{E} \mathbf{u}_q) - \frac{1}{2} \mathbf{v}_f^T \mathbf{B}^i \mathbf{u}_f + \mathbf{v}_f^T \mathbf{B}^i \mathbf{u}_f \\
313 \quad &= \begin{pmatrix} \mathbf{v}_q \\ \mathbf{v}_f \end{pmatrix}^T \left(\begin{pmatrix} \mathbf{0} & \mathbf{B}^i \end{pmatrix} + \begin{pmatrix} -\mathbf{Q}_i^T + \frac{1}{2} \mathbf{E}^T \mathbf{B}^i \mathbf{E} & -\frac{1}{2} (\mathbf{B}^i \mathbf{E})^T \\ \frac{1}{2} \mathbf{B}^i \mathbf{E} & -\frac{1}{2} \mathbf{B}^i \end{pmatrix} \right) \begin{pmatrix} \mathbf{u}_q \\ \mathbf{u}_f \end{pmatrix} \\
314 \quad &= \mathbf{v}^T \left(\mathbf{B}_N^i - (\mathbf{Q}_N^i)^T \right) \mathbf{u}. \quad \square \\
315
\end{aligned}$$

316 **5.3. Entropy stability on a reference element.** In this section, we construct
317 so-called “entropy stable” schemes on the reference element \widehat{D} . These methods ensure
318 that the entropy inequality (3) is satisfied discretely by avoiding the use of the chain
319 rule in the proof of entropy dissipation. Entropy stable schemes rely on two main
320 ingredients: an entropy stable numerical flux as defined by Tadmor [28] and a concept
321 referred to as “flux differencing”. Let $\mathbf{f}_S(\mathbf{u}_L, \mathbf{u}_R)$ be a numerical flux function which
322 is a function of “left” and “right” states $\mathbf{u}_L, \mathbf{u}_R$. The numerical flux \mathbf{f}_S is *entropy*
323 *conservative* if it satisfies the following three conditions:

$$\begin{aligned}
324 \quad (9) \quad \mathbf{f}_S^i(\mathbf{u}, \mathbf{u}) &= \mathbf{f}_i(\mathbf{u}), \quad (\text{consistency}) \\
325 \quad \mathbf{f}_S^i(\mathbf{u}_L, \mathbf{u}_R) &= \mathbf{f}_S^i(\mathbf{u}_R, \mathbf{u}_L), \quad (\text{symmetry}) \\
326 \quad (\mathbf{v}_L - \mathbf{v}_R)^T \mathbf{f}_S^i(\mathbf{u}_L, \mathbf{u}_R) &= \psi_i(\mathbf{u}_L) - \psi_i(\mathbf{u}_R), \quad (\text{conservation}) \\
327
\end{aligned}$$

328 for $j = 1, \dots, d$. The construction of entropy stable schemes will utilize (5.3) in
329 discretizations of both volume and surface terms in a DG formulation.

330 Proving a semi-discrete entropy inequality typically involves the “strong” SBP
331 condition. The schemes presented in this work will relax this, and use only two
332 (weaker) conditions: that the decoupled SBP operator \mathbf{D}_N^i exactly differentiates con-
333 stants and that a discrete form of the fundamental theorem of calculus (FTC) holds.
334 The latter property can be derived from the variational SBP property in Lemma 3.

335 **LEMMA 4.** (*Discrete FTC and exact differentiation of constants*) Suppose As-
336 sumption 1 holds for $v(\mathbf{x}) = 1$. Let $u(\mathbf{x})$ be some L^2 integrable function, and let \mathbf{u}
337 denote the vector of values of $u(\mathbf{x})$ at volume and surface quadrature points. Then,

$$338 \quad \mathbf{1}^T \mathbf{Q}_N^i \mathbf{u} = \mathbf{1}^T \mathbf{B}_N^i \mathbf{u}, \quad \mathbf{Q}_N^i \mathbf{1} = \mathbf{0}.$$

339 *Proof.* The proof that $\mathbf{Q}_N^i \mathbf{1} = \mathbf{0}$ follows from the property that polynomials are
340 equal to their L^2 projection, and is identical to that of [5, 6]. The proof of the first
341 equality follows from $\mathbf{Q}_N^i \mathbf{1} = \mathbf{0}$ and Lemma 3

$$342 \quad \mathbf{1}^T (\mathbf{Q}_N^i) \mathbf{u} = \mathbf{1}^T \left(\mathbf{B}_N^i - (\mathbf{Q}_N^i)^T \right) \mathbf{u} = \mathbf{1}^T \mathbf{B}_N^i \mathbf{u}. \quad \square$$

343 We can now construct a skew-symmetric formulation on the reference element
344 \widehat{D} and show that it is semi-discretely entropy conservative. This formulation can
345 be made entropy stable by adding interface dissipation. Let \mathbf{u}_h denote the discrete
346 solution, and let \mathbf{u}_q denote the values of the solution at volume quadrature points.
347 We define the auxiliary conservative variables $\tilde{\mathbf{u}}$ in terms of the L^2 projections of the
348 entropy variables

$$349 \quad (10) \quad \mathbf{v}_q = \mathbf{v}(\mathbf{u}_q), \quad \tilde{\mathbf{v}} = \begin{bmatrix} \mathbf{V}_q \\ \mathbf{V}_f \end{bmatrix} \mathbf{P}_q \mathbf{v}_q, \quad \tilde{\mathbf{u}} = \mathbf{u}(\tilde{\mathbf{v}}).$$

351 The skew-symmetric formulation for (1) on \hat{D} is given in terms of $\tilde{\mathbf{u}}$

$$352 \quad (11) \quad \mathbf{M} \frac{\partial \mathbf{u}_h}{\partial t} + \sum_{i=1}^d \begin{bmatrix} \mathbf{V}_q \\ \mathbf{V}_f \end{bmatrix}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S^i \right) \mathbf{1} + \mathbf{V}_f^T \mathbf{B}^i \mathbf{f}_i^* = 0,$$

$$353 \quad (\mathbf{F}_S^i)_{ij} = \mathbf{f}_S^i(\tilde{\mathbf{u}}_i, \tilde{\mathbf{u}}_j), \quad 1 \leq i, j \leq N_q + N_q^f,$$

355 where \mathbf{f}^* is some numerical flux, and the matrix $(\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T)$ possesses the fol-
356 lowing block structure:

$$357 \quad \left(\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T \right) = \begin{pmatrix} \mathbf{Q}_i - \mathbf{Q}_i^T & \mathbf{E}^T \mathbf{B}^i \\ -\mathbf{B}^i \mathbf{E} & \mathbf{0} \end{pmatrix}.$$

358 Multiplying the formulation (11) by \mathbf{M}^{-1} on both sides yields a strong form

$$359 \quad (12) \quad \frac{\partial \mathbf{u}_h}{\partial t} + \sum_{i=1}^d \begin{bmatrix} \mathbf{P}_q & \mathbf{L}_f \end{bmatrix} \left((\mathbf{D}_N^i - \mathbf{W}_N^{-1} (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S^i \right) \mathbf{1} + \mathbf{L}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{f}_i^* = 0,$$

361 We can now show that the skew-symmetric formulation is entropy conservative.

362 **THEOREM 5.** *Suppose Assumption 1 holds for $v(\mathbf{x}) = 1$. Then, the formulation*
363 *(11) is entropy conservative such that*

$$364 \quad (13) \quad \mathbf{1}^T \mathbf{W} \frac{\partial U(\mathbf{u}_q)}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \mathbf{B}^i (\psi_i(\tilde{\mathbf{u}}_f) - \tilde{\mathbf{v}}_f^T \mathbf{f}_i^*) = 0, \quad \mathbf{u}_q = \mathbf{V}_q \mathbf{u}.$$

365

366 *Proof.* Testing (11) by $\mathbf{v}_h = \mathbf{P}_q \mathbf{v}_q$ yields

$$367 \quad (14) \quad \mathbf{v}_q^T \mathbf{W} \frac{\partial (\mathbf{V}_q \mathbf{u})}{\partial t} + \sum_{i=1}^d \tilde{\mathbf{v}}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S^i \right) \mathbf{1} + \tilde{\mathbf{v}}_f^T \mathbf{B}^i \mathbf{f}_i^* = 0.$$

368

369 One can show that [5]

$$370 \quad \tilde{\mathbf{v}}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S^i \right) \mathbf{1} = \tilde{\mathbf{v}}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S^i) \mathbf{1} - \tilde{\mathbf{v}}^T ((\mathbf{Q}_N^i)^T \circ \mathbf{F}_S^i) \mathbf{1}$$

$$371 \quad = \tilde{\mathbf{v}}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S^i) \mathbf{1} - \mathbf{1}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S^i) \tilde{\mathbf{v}}.$$

373 Where we have used that \mathbf{F}_S^i is symmetric and that the Hadamard product commutes.

374 Applying the conservation condition on \mathbf{f}_S^i in (5.3) then yields

$$375 \quad \tilde{\mathbf{v}}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S^i) \mathbf{1} - \mathbf{1}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S^i) \tilde{\mathbf{v}}$$

$$376 \quad = \sum_{jk} (\mathbf{Q}_N^i)_{jk} (\tilde{\mathbf{v}}_j - \tilde{\mathbf{v}}_k) \mathbf{f}_S^i(\tilde{\mathbf{u}}_j, \tilde{\mathbf{u}}_k) = \sum_{jk} (\mathbf{Q}_N^i)_{jk} (\psi_i(\tilde{\mathbf{u}}_j) - \psi_i(\tilde{\mathbf{u}}_k))$$

$$377 \quad = \mathbf{1}^T (\mathbf{Q}_N^i) \psi_i(\tilde{\mathbf{u}}) - \psi_i(\tilde{\mathbf{u}})^T (\mathbf{Q}_N^i) \mathbf{1} = \mathbf{1}^T (\mathbf{Q}_N^i) \psi_i(\tilde{\mathbf{u}}) = \mathbf{1}^T \mathbf{B}_N^i \psi_i(\tilde{\mathbf{u}}),$$

379 where we have used Corollary 4 in the last equality. Noting that $\mathbf{1}^T \mathbf{B}_N^i \psi_i(\tilde{\mathbf{u}}) =$
380 $\mathbf{1}^T \mathbf{B}^i \psi_i(\tilde{\mathbf{u}}_f)$ completes the proof. \square

381 **REMARK.** *The primary difference between this formulation and the formulation*
382 *presented in [5] is the use of the skew-symmetric volume operator $(\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T)$.*
383 *When the strong SBP property (8) holds, this formulation is exactly equivalent to*
384 *the strong formulation presented in [5]. However, the only result necessary to prove*
385 *Theorem 5 is Corollary 4. The decoupled SBP property (8) is not utilized, and is not*
386 *necessary to guarantee a semi-discrete conservation of entropy for the skew-symmetric*
387 *formulation.*

The skew symmetric formulation can also be shown to be locally conservative in the sense of [29], which is sufficient to show the numerical solution convergences to the weak solution under mesh refinement.

THEOREM 6. *The formulation (11) is locally conservative such that*

$$(15) \quad \mathbf{1}^T \mathbf{W} \frac{\partial (\mathbf{V}_q \mathbf{u})}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \mathbf{B}^i \mathbf{f}_i^* = 0.$$

Proof. To show local conservation, we test (11) with 1

$$(16) \quad \mathbf{1}^T \mathbf{W} \mathbf{V}_q \frac{\partial \mathbf{u}_h}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} + \mathbf{1}^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) \mathbf{f}_i^* = 0.$$

Because \mathbf{F}_S is symmetric and $(\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T)$ is skew-symmetric, their Hadamard product is also skew-symmetric. Using that $\mathbf{x}^T \mathbf{A} \mathbf{x} = 0$ for any skew symmetric matrix \mathbf{A} , the volume term vanishes

$$\mathbf{1}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} = 0. \quad \square$$

5.4. On quadrature conditions for Assumption 1 with $v = 1$. Apart from algebraic manipulations, only Corollary 4 is necessary to prove entropy conservation for (12). Corollary 4 requires that Assumption 1 holds for $v = 1$. Thus, the volume and surface quadratures must be sufficiently accurate to guarantee that the mass matrix is positive definite and to integrate

$$(17) \quad \int_{\hat{D}} \frac{\partial u}{\partial x_i}, \quad \int_{\partial \hat{D}} u n_i.$$

On simplicial elements, the mass matrix is guaranteed to be positive definite for any volume quadrature which is exact for degree $2N$ polynomial integrands. This choice of volume quadrature also guarantees that the volume term in (17) is integrated exactly. The surface quadrature can thus be taken to be any quadrature rule which is exact for only degree N integrands on faces. In contrast, the construction of simplicial decoupled SBP operators has required face quadratures which are accurate for degree $2N$ polynomials [5, 6].

On tensor product elements, we can take any degree $(2N - 1)$ quadrature rule which ensures a positive definite mass matrix (e.g. a $(N + 1)$ -point GLL quadrature), as a quadrature of this accuracy is sufficient to exactly integrate the volume term in (17). For the surface quadrature, we can again take any quadrature rule which is exact for degree N polynomial integrands. For example, on a quadrilateral element, one can use $\lceil \frac{N+1}{2} \rceil$ -point Gauss quadrature rule or a $\lceil \frac{N+3}{2} \rceil$ -point GLL rule as face quadratures for a degree N scheme.

5.5. On the accuracy of the skew-symmetric formulation. We note that, while the proofs in this Section show that it is possible to maintain entropy stability under reduced accuracy surface quadrature rules, the accuracy of the resulting schemes depend on the accuracy of the volume and surface quadrature.

It was shown in [5] that, when combined with quadrature-based L^2 projection and lifting operators, the decoupled SBP operator (8) is a degree N approximation

to the derivative. This approximation can be interpreted as augmenting a volume approximation of the derivative with boundary correction terms [4]. Let $f(\mathbf{x}), g(\mathbf{x})$ denote two L^2 integrable functions, and let $\mathbf{f}_N, \mathbf{g}_N$ denote the vectors of values of f, g at both volume and surface quadrature points. A degree N approximation $u \in V^N$ to $f \frac{\partial g}{\partial x_i}$ can be constructed via

$$\mathbf{M}\mathbf{u} = \begin{bmatrix} \mathbf{V}_q \\ \mathbf{V}_f \end{bmatrix}^T \text{diag}(\mathbf{f}_N) \mathbf{Q}_N^i \mathbf{g}_N, \quad \mathbf{f}_N = \begin{bmatrix} \mathbf{f}_q \\ \mathbf{f}_f \end{bmatrix}, \quad \mathbf{g}_N = \begin{bmatrix} \mathbf{g}_q \\ \mathbf{g}_f \end{bmatrix},$$

where \mathbf{u} denotes the vector of coefficients for u . Expanding terms using (8) yields

$$\begin{aligned} \mathbf{M}\mathbf{u} &= \mathbf{V}_q^T \text{diag}(\mathbf{f}_q) \mathbf{Q}^i \mathbf{g}_q + \frac{1}{2} \mathbf{V}_q^T \text{diag}(\mathbf{f}_q) \mathbf{E}^T \mathbf{B}^i (\mathbf{g}_f - \mathbf{E} \mathbf{g}_q) \\ &\quad + \frac{1}{2} \mathbf{V}_f^T \text{diag}(\mathbf{f}_f) \mathbf{B}^i (\mathbf{g}_f - \mathbf{E} \mathbf{g}_q). \end{aligned}$$

where we have used that \mathbf{B}^i is diagonal and that diagonal matrices commute. Let \mathbf{v} are coefficients of some test function $v \in V^N$, and let $\mathbf{v}_q = \mathbf{V}_q \mathbf{v}$ and $\mathbf{v}_f = \mathbf{V}_f \mathbf{v}$. Multiplying by \mathbf{v}^T on both sides yields that $\mathbf{v}^T \mathbf{M}\mathbf{u}$ is equal to

$$(\text{diag}(\mathbf{f}_q) \mathbf{v}_q)^T \mathbf{Q}^i \mathbf{g}_q + \frac{1}{2} (\text{diag}(\mathbf{f}_f) \mathbf{v}_f + \mathbf{E} \text{diag}(\mathbf{f}_q) \mathbf{v}_q)^T \mathbf{B}^i (\mathbf{g}_f - \mathbf{E} \mathbf{g}_q).$$

This algebraic expression can be reinterpreted as a quadrature approximation of the following variational problem

$$(18) \quad \int_{\hat{D}} uv = \int_{\hat{D}} f \frac{\partial \Pi_N g}{\partial x_i} v + \int_{\partial \hat{D}} (g - \Pi_N g) \left(\frac{fv + \Pi_N(fv)}{2} \right) \hat{n}_i, \quad \forall v \in V^N.$$

The skew-symmetric formulation can be interpreted as approximating a skew symmetric form of (18). Integrating half of the volume term by parts yields

$$\begin{aligned} (19) \quad & \frac{1}{2} \int_{\hat{D}} \left(f \frac{\partial \Pi_N g}{\partial x_k} v - g \frac{\partial \Pi_N(fv)}{\partial x_i} \right) \\ & + \frac{1}{2} \int_{\partial \hat{D}} \left(f g v \hat{n}_i + (g - \Pi_N g) \left(\frac{fv + \Pi_N(fv)}{2} \right) \hat{n}_i \right) \end{aligned}$$

Approximating integrals using quadrature allows us to express (19) as a matrix equation using \mathbf{B}_N^i in (8)

$$\mathbf{M}\mathbf{u} = \frac{1}{2} \left(\begin{bmatrix} \mathbf{V}_q \\ \mathbf{V}_f \end{bmatrix} \right)^T (\text{diag}(\mathbf{f}_N) (\mathbf{Q}_N - \mathbf{Q}_N^T) \mathbf{g}_N) + \mathbf{V}_f^T \text{diag}(\mathbf{f}_f) \mathbf{B}^i \mathbf{g}_f.$$

The accuracy of this approximation depends on the degree of accuracy of the volume and surface quadrature rules.

The entropy conservative skew-symmetric formulation (11)

Fix this up: if IBP holds for $v \in P^M$, the approximation should be $O(h^M)$.

Add in continuous formulation and note that skew-symmetry approximates it.

6. Skew-symmetric entropy conservative formulations on mapped elements. We now construct skew-symmetric formulations on mapped elements. We assume some domain Ω is decomposed into non-overlapping elements D^k , such that D^k is the image of the reference element \hat{D} under an isoparametric mapping Φ^k .

We define geometric change of variables terms G_{ij}^k as scaled derivatives of reference coordinates $\hat{\mathbf{x}}$ w.r.t. physical coordinates \mathbf{x}

$$(20) \quad \frac{\partial u}{\partial x_i} = \sum_{ij} G_{ij}^k \frac{\partial u}{\partial \hat{x}_j}, \quad G_{ij}^k = J^k \frac{\partial \hat{x}_j}{\partial x_i},$$

where J^k is the determinant of the Jacobian of the geometric mapping on the element D^k . We also introduce the scaled outward normal components $n_i J_f^k$, which can be computed in terms of (20) and the reference normals $\hat{\mathbf{n}}$ on \hat{D}

$$(21) \quad n_i^k J_f^k = \sum_{j=1}^d G_{ij}^k \hat{n}_j.$$

We also define \mathbf{n}_i^k as the vector containing concatenated values of the scaled outward normals $n_i^k J_f^k$ at surface quadrature nodes. For the remainder of the work, we assume that the mesh is watertight [6] such that at all points on any internal face, the scaled outward normals $n_i^k J_f^k$ on the two elements sharing this face are equal and opposite.

On a single element (and on affine meshes), it is possible to show entropy stability of the skew-symmetric formulation (11) under a surface quadrature which is only exact for degree N polynomials. However, on curved meshes, stronger conditions are required to guarantee entropy stability. This is due to the fact that the geometric terms are now high order polynomials which vary spatially over each element. Moreover, Corollary 4 assumes affine geometric mappings, and does not hold on curved elements. In this section, we discuss how to extend Corollary 4 to curved simplicial and tensor product elements.

6.1. Curved elements and the geometric conservation law. In this section, we describe how to construct appropriate decoupled SBP operators on curved meshes, and give conditions on the volume and surface quadrature rules under which a semi-discretely entropy stable scheme can be constructed.

We first show how to construct decoupled SBP operators on curved elements. Let \mathbf{G}_{ij}^k denote the vector of scaled geometric terms G_{ij}^k evaluated at both volume and surface quadrature points, and let $\hat{\mathbf{D}}_N^j$ denote the decoupled SBP operator for the j th reference coordinate. Decoupled SBP operators on a curved element D^k can be defined as in [6] by

$$(22) \quad \mathbf{D}_k^i = \frac{1}{2} \sum_{j=1}^d \left(\text{diag}(\mathbf{G}_{ij}^k) \hat{\mathbf{D}}_N^j + \hat{\mathbf{D}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \right), \quad \mathbf{Q}_k^i = \mathbf{W} \mathbf{D}_k^i.$$

It can be shown that, if $\hat{\mathbf{D}}_N^j$ satisfies a strong summation by parts property, \mathbf{D}_k^i satisfies an analogous strong summation by parts property.

For curved elements, steps must also be taken to ensure that the analogue of Lemma 3 and Corollary 4 hold at the discrete level. Expanding out the condition $\mathbf{Q}_k^i \mathbf{1} = \mathbf{0}$ in terms of (22) yields

$$(23) \quad \mathbf{Q}_k^i \mathbf{1} = \frac{1}{2} \mathbf{W} \sum_{j=1}^d \text{diag}(\mathbf{G}_{ij}^k) \hat{\mathbf{D}}_N^j \mathbf{1} + \hat{\mathbf{D}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \mathbf{1} = \frac{1}{2} \mathbf{W} \sum_{j=1}^d \hat{\mathbf{D}}_N^j (\mathbf{G}_{ij}^k) = 0,$$

where we have used that $\hat{\mathbf{D}}_N^j \mathbf{1} = 0$. We refer to the condition $\mathbf{Q}_k^i \mathbf{1} = 0$ as the discrete geometric conservation law (GCL) [30, 31]. For degree N isoparametric mappings,

the GCL is automatically satisfied in two dimensions due to the fact that the exact geometric terms G_{ij}^k are polynomials of degree N [31]. However, in three dimensions, the GCL is not automatically satisfied due to the fact that the degree of G_{ij}^k is larger than N . Thus, the geometric terms cannot be represented exactly using degree N polynomials, and (23) must be enforced through an alternative construction of G_{ij}^k .

A common approach is to rewrite the geometric terms as the curl of some quantity \mathbf{r}^i , but to interpolate \mathbf{r}^i before applying the curl [32, 31, 33, 6]:

$$(24) \quad \mathbf{r}^i = \frac{\partial \mathbf{x}}{\partial \hat{x}_i} \times \mathbf{x}, \quad \begin{bmatrix} G_{i1}^k \\ G_{i2}^k \\ G_{i3}^k \end{bmatrix} = -\frac{1}{2} \left(\frac{\partial I_{N_{\text{geo}}} \mathbf{r}^k}{\partial \hat{x}_j} - \frac{\partial I_{N_{\text{geo}}} \mathbf{r}^j}{\partial \hat{x}_k} \right),$$

$$N_{\text{geo}} \leq \begin{cases} N+1 & (\text{tetrahedra}) \\ N & (\text{hexahedra}) \end{cases},$$

where $I_{N_{\text{geo}}}$ denotes a degree N_{geo} polynomial interpolation operator with appropriate interpolation nodes.¹ The restriction on the maximum value of N_{geo} ensures that $G_{ij}^k \in V^N$ (e.g. $G_{ij}^k \in P^N$ on tetrahedral elements and $G_{ij}^k \in Q^N$ on hexahedral elements), which is necessary to guarantee (23).

Because the decoupled SBP operators \mathbf{D}_k^i are now defined through (22), Corollary 4 and the proof of entropy stability no longer hold for curved elements and must be modified. The introduction of curvilinear meshes will impose slightly different conditions on the accuracy of the surface quadrature. We discuss simplicial and tensor product elements separately, as differences in the natural polynomial approximation spaces will result in different assumptions for each proof.

LEMMA 7. (*Discrete FTC and exact constant differentiation on curved elements*)
 Let D^k be a curved element, and let the geometric terms G_{ij}^k be constructed using (6.1). Let Assumption 1 hold for $v = 1$ and $v = G_{ij}^k$ for all $i, j = 1, \dots, d$, and let \mathbf{u} denote a vector of values of some function at volume and surface quadrature points. Then,

$$\mathbf{1}^T \mathbf{Q}_k^i \mathbf{u} = \mathbf{1}^T \mathbf{B}_k^i \mathbf{u}, \quad \mathbf{Q}_k^i \mathbf{1} = \mathbf{0},$$

where the physical boundary matrix \mathbf{B}_k^i is defined as

$$\mathbf{B}_k^i = \sum_{j=1}^d \mathbf{B}_N^j \text{diag}(\mathbf{G}_{ij}^k) = \sum_{j=1}^d \text{diag}(\mathbf{G}_{ij}^k) \mathbf{B}_N^j = \begin{pmatrix} \mathbf{0} & \mathbf{W}_f \text{diag}(\mathbf{n}_i^k) \end{pmatrix}.$$

Proof. The proof of $\mathbf{Q}_k^i \mathbf{1} = \mathbf{0}$ was shown for tensor product elements in [31] and for simplicial elements in [6], and relies only on the fact that $G_{ij}^k \in V^N$. We focus on showing that the discrete fundamental theorem of calculus $\mathbf{1}^T \mathbf{Q}_k^i \mathbf{u} = \mathbf{1}^T \mathbf{B}_k^i \mathbf{u}$ holds. Using (22), we have that

$$\begin{aligned} \mathbf{1}^T \mathbf{Q}_k^i \mathbf{u} &= \frac{1}{2} \sum_{j=1}^d \mathbf{1}^T \left(\text{diag}(\mathbf{G}_{ij}^k) \hat{\mathbf{Q}}_N^j + \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \right) \mathbf{u} \\ &= \frac{1}{2} \sum_{j=1}^d \left((\mathbf{G}_{ij}^k)^T \hat{\mathbf{Q}}_N^j \mathbf{u} + \mathbf{1}^T \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \mathbf{u} \right). \end{aligned}$$

¹This interpolation step must be performed using interpolation points with an appropriate number of nodes on each boundary [6]. These include, for example, GLL nodes on tensor product elements, as well as Warp and Blend nodes on non-tensor product elements [34, 35].

Applying Lemma 3 to $\sum_{j=1}^d \mathbf{1}^T \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \mathbf{u}$ yields

$$\begin{aligned} \sum_{j=1}^d \mathbf{1}^T \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \mathbf{u} &= \sum_{j=1}^d \mathbf{1}^T \left(\mathbf{B}_N^j - \left(\hat{\mathbf{Q}}_N^j \right)^T \right) \text{diag}(\mathbf{G}_{ij}^k) \mathbf{u} \\ &= \sum_{j=1}^d \mathbf{1}^T \mathbf{B}_N^j \text{diag}(\mathbf{G}_{ij}^k) \mathbf{u}, \end{aligned}$$

where we have used Lemma 4 to conclude that $\hat{\mathbf{Q}}_N^j \mathbf{1} = 0$. Repeating the procedure for $\sum_{j=1}^d (\mathbf{G}_{ij}^k)^T \hat{\mathbf{Q}}_N^j \mathbf{u}$ yields

$$\begin{aligned} \sum_{j=1}^d (\mathbf{G}_{ij}^k)^T \hat{\mathbf{Q}}_N^j \mathbf{u} &= \sum_{j=1}^d (\mathbf{G}_{ij}^k)^T \left(\mathbf{B}_N^j - \left(\hat{\mathbf{Q}}_N^j \right)^T \right) \mathbf{u} \\ &= \sum_{j=1}^d (\mathbf{G}_{ij}^k)^T \mathbf{B}_N^j \mathbf{u} - \sum_{j=1}^d \mathbf{u}^T \hat{\mathbf{Q}}_N^j \mathbf{G}_{ij}^k = \sum_{j=1}^d (\mathbf{G}_{ij}^k)^T \mathbf{B}_N^j \mathbf{u}, \end{aligned}$$

where we have used that the \mathbf{G}_{ij}^k satisfies the discrete GCL. The proof is completed by noting that \mathbf{B}_N^j is diagonal and thus commutes with $\text{diag}(\mathbf{G}_{ij}^k)$, such that

$$\mathbf{1}^T \mathbf{Q}_k^i \mathbf{u} = \sum_{j=1}^d \mathbf{1}^T \left(\mathbf{B}_N^j \text{diag}(\mathbf{G}_{ij}^k) \right) \mathbf{u} = \mathbf{1}^T \mathbf{B}_k^i \mathbf{u}.$$

The property $\mathbf{Q}_k^i \mathbf{1} = 0$ is also related to free-stream preservation [31]. Constant solutions are stationary solutions of systems of conservation laws. However, on curved meshes, the presence of spatially varying geometric terms can result in the production of spurious transient waves. The construction of geometric terms through (6.1) guarantees that the resulting methods are free-stream preserving, and that constant solutions remain stationary solutions of discretizations of (1).

We can now construct and prove entropy conservation and free stream preservation for a skew-symmetric formulation on a curved element D^k . Let \mathbf{Q}_k^i be given by (22), and define the curved mass matrix $\mathbf{M}^k = \mathbf{V}_q^T \mathbf{W} \text{diag}(\mathbf{J}^k) \mathbf{V}_q$. Note that \mathbf{M}^k is positive-definite so long as \mathbf{J}^k is positive at all quadrature points. We define the auxiliary quantities $\tilde{\mathbf{u}}$

$$\mathbf{v}_q = \mathbf{v}(\mathbf{u}_q), \quad \tilde{\mathbf{v}} = \begin{bmatrix} \mathbf{V}_q \\ \mathbf{V}_f \end{bmatrix} \mathbf{P}_q^k \mathbf{v}_q, \quad \tilde{\mathbf{u}} = \mathbf{u}(\tilde{\mathbf{v}}).$$

where $\mathbf{P}_q^k = (\mathbf{M}^k)^{-1} \mathbf{V}_q^T \mathbf{W} \text{diag}(\mathbf{J}^k)$. Then, we have the following theorem:

THEOREM 8. *Let the geometric terms \mathbf{G}_{ij}^k be computed from (6.1), let the scaled outward normals \mathbf{n}_i^k be computed pointwise from (21), and suppose Assumption 1 holds for $v = 1$ and $v = \mathbf{G}_{ij}^k$. Let $\tilde{\mathbf{u}}_f^+$ denote the face value of the entropy-projected conservative variables $\tilde{\mathbf{u}}_f$ on the neighboring element. Then, the formulation*

$$\begin{aligned} (25) \quad \mathbf{M}^k \frac{\partial \mathbf{u}_h}{\partial t} + \sum_{i=1}^d \begin{bmatrix} \mathbf{V}_q \\ \mathbf{V}_f \end{bmatrix}^T \left(\left(\mathbf{Q}_k^i - (\mathbf{Q}_k^i)^T \right) \circ \mathbf{F}_S^i \right) \mathbf{1} + \mathbf{V}_f^T \mathbf{W}_f \text{diag}(\mathbf{n}_i^k) \mathbf{f}_i^* &= 0, \\ (\mathbf{F}_S^i)_{ij} &= \mathbf{f}_S^i(\tilde{\mathbf{u}}_i, \tilde{\mathbf{u}}_j), \quad 1 \leq i, j \leq N_q + N_q^f, \\ \mathbf{f}_i^* &= \mathbf{f}_S^i(\tilde{\mathbf{u}}_f^+, \tilde{\mathbf{u}}_f), \quad \text{on interior interfaces,} \end{aligned}$$

572 is semi-discretely entropy conservative on D^k such that for $\mathbf{u}_q = \mathbf{V}_q \mathbf{u}$,

$$573 \quad \mathbf{1}^T \mathbf{W} \text{diag}(\mathbf{J}^k) \frac{\partial U(\mathbf{u}_q)}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \mathbf{W}_f \text{diag}(\mathbf{n}_i^k) (\psi_i(\tilde{\mathbf{u}}_f) - \tilde{\mathbf{v}}_f^T \mathbf{f}_i^*) = 0.$$

574
575 Additionally, the method is free-stream preserving such that $\frac{\partial \mathbf{u}_h}{\partial t} = 0$ for constant
576 solutions.

577 *Proof.* The proof of entropy stability is identical to that of Theorem 5, except that
578 Lemma 7 is used instead of Lemma 4. However, the proof of free-stream preservation
579 differs from proofs found in the literature, which typically show that the right hand
580 side is zero for \mathbf{u}_h constant [14, 6]. This strategy for is appropriate for “strong”
581 formulations, but must be modified for the skew-symmetric formulation (25).

582 We have that $\mathbf{f}_i^* = \mathbf{f}_S^i(\tilde{\mathbf{u}}_f^+, \tilde{\mathbf{u}}_f) = \mathbf{f}_i(\mathbf{u}_f)$ using the consistency property of the
583 Tadmor flux (5.3) and the fact that the solution is globally constant. For \mathbf{u}_h constant,
584 \mathbf{F}_S is also constant. We assume (without loss of generality) that \mathbf{F}_S and $\mathbf{f}_i(\mathbf{u}_f)$ are
585 the matrix and vector of ones, respectively. Then,

$$586 \quad \left((\mathbf{Q}_k^i - (\mathbf{Q}_k^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} = \left(\mathbf{Q}_k^i - (\mathbf{Q}_k^i)^T \right) \mathbf{1} = (\mathbf{Q}_k^i)^T \mathbf{1},$$

587 where we have used that $\mathbf{Q}_k^i \mathbf{1} = 0$ by Lemma 7. Thus, the right hand side reduces to

$$588 \quad \sum_{j=1}^d \left[\begin{array}{c} \mathbf{V}_q \\ \mathbf{V}_f \end{array} \right]^T \left(-(\mathbf{Q}_k^i)^T + \mathbf{V}_f^T \mathbf{W}_f \text{diag}(\mathbf{n}_i^k) \right) \mathbf{1}$$

$$589 \quad = \sum_{j=1}^d \left[\begin{array}{c} \mathbf{V}_q \\ \mathbf{V}_f \end{array} \right]^T \left(\left(-\frac{1}{2} \left(\text{diag}(\mathbf{G}_{ij}^k) \hat{\mathbf{Q}}_N^j + \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \right)^T + \hat{\mathbf{B}}_N^j \text{diag}(\mathbf{G}_{ij}^k) \right) \right) \mathbf{1},$$

591 where we have used the definition of \mathbf{Q}_k^i in (22). Rearranging terms gives that, for a
592 constant solution, (25) reduces to

$$593 \quad \mathbf{M}^k \frac{\partial \mathbf{u}_h}{\partial t} + \frac{1}{2} \sum_{j=1}^d \left[\begin{array}{c} \mathbf{V}_q \\ \mathbf{V}_f \end{array} \right]^T \left(-(\hat{\mathbf{Q}}_N^j)^T + \hat{\mathbf{B}}_N^j \right) \mathbf{G}_{ij}^k$$

$$594 \quad + \frac{1}{2} \sum_{j=1}^d \left[\begin{array}{c} \mathbf{V}_q \\ \mathbf{V}_f \end{array} \right]^T \text{diag}(\mathbf{G}_{ij}^k) \left(-(\hat{\mathbf{Q}}_N^j)^T + \hat{\mathbf{B}}_N^j \right) \mathbf{1} = 0.$$

596 Let u denote the constant value of the solution such that the values of the solution at
597 quadrature points is $u\mathbf{1}$. Multiplying both sides by the constant solution \mathbf{u}_h^T yields

$$598 \quad \frac{1}{2} \frac{\partial}{\partial t} (\mathbf{u}_h^T \mathbf{M}^k \mathbf{u}_h) +$$

$$599 \quad \frac{u}{2} \sum_{j=1}^d \left(\mathbf{1}^T \left(-(\hat{\mathbf{Q}}_N^j)^T + \hat{\mathbf{B}}_N^j \right) \mathbf{G}_{ij}^k + (\mathbf{G}_{ij}^k)^T \left(-(\hat{\mathbf{Q}}_N^j)^T + \hat{\mathbf{B}}_N^j \right) \mathbf{1} \right) = 0.$$

601 By assumption, Lemma 3 holds for test functions $\mathbf{1}$ and \mathbf{G}_{ij}^k , and the formulation
602 reduces to

$$603 \quad \frac{1}{2} \frac{\partial}{\partial t} (\mathbf{u}_h^T \mathbf{M}^k \mathbf{u}_h) + \frac{u}{2} \sum_{j=1}^d \left(\mathbf{1}^T \hat{\mathbf{Q}}_N^j \mathbf{G}_{ij}^k + (\mathbf{G}_{ij}^k)^T \hat{\mathbf{Q}}_N^j \mathbf{1} \right) = 0.$$

We have that $\widehat{Q}_N^j \mathbf{1} = 0$ by construction, and $\sum_{j=1}^d \mathbf{1}^T \widehat{Q}_N^j G_{ij}^k = 0$ since G_{ij}^k satisfies the GCL. This implies that $\frac{\partial}{\partial t} (\mathbf{u}_h^T \mathbf{M}^k \mathbf{u}_h) = 0$ on D^k . Since \mathbf{M}^k is symmetric and positive-definite, $\mathbf{u}_h^T \mathbf{M}^k \mathbf{u}_h$ is a norm on \mathbf{u}_h . Summing $\frac{\partial}{\partial t} (\mathbf{u}_h^T \mathbf{M}^k \mathbf{u}_h) = 0$ over all elements ensures that global constants are stationary solutions. \square

The proof of global entropy conservation follows from summing up (13) over all elements and noting that the surface terms cancel due to the symmetry and conservation properties of the Tadmor flux (5.3) [5]. The entropy conservative formulations presented in this work can be made entropy stable by adding appropriate interface dissipation, such as Lax-Friedrichs or matrix-based penalization terms [36, 13, 5].

REMARK. *It is also possible to replace the curved mass matrix \mathbf{M}^k with a more easily invertible weight-adjusted approximation while maintaining high order accuracy, entropy stability, and local conservation [6]. This approximation avoids the inversion of dense weighted L^2 mass matrices \mathbf{M}^k on curved simplicial elements. On tensor product elements, this approximation is generally unnecessary as common choices of volume quadrature result in a diagonal (lumped) mass matrix [3, 15, 4].*

6.2. On quadrature conditions for Assumption 1 for $v = 1$ and $v = G_{ij}^k$.

Semi-discrete entropy conservation on curved meshes requires that Assumption 1 holds for $v = 1$ and $v = G_{ij}^k$. We discuss specific choices of volume and surface quadrature on for which this assumption is valid. *Emphasize we're interested in taking N_{geo} as large as possible for a given quadrature rule, and that there is a focus on degree $(2N - 1)$ quadrature rules.*

6.2.1. Simplicial elements.

On simplicial elements, for $u \in P^N$, $\frac{\partial u}{\partial \hat{x}_j} \in P^{N-1}$, and for $N_{\text{geo}} \leq (N + 1)$, $G_{ij}^k \in P^{N_{\text{geo}}-1}$. Thus, the integrands in Assumption 1 $\frac{\partial u}{\partial \hat{x}_j} v \in P^{N+N_{\text{geo}}-1}$ and $u v n_i \in P^{N+N_{\text{geo}}}$ for $v = G_{ij}^k$.

Thus, a volume quadrature rule of degree $(2N - 1)$ and surface quadrature of degree $2N$ are sufficient to ensure semi-discrete entropy conservation for $N_{\text{geo}} \leq N$. *Add discussion on triangular elements.*

6.2.2. Tensor product elements.

It was shown in Section 5.4 that tensor product quadratures of degree $(2N - 1)$ satisfy Assumption 1 holds for $v = 1$. However, in contrast to the simplicial case, it is not immediately clear that degree $(2N - 1)$ volume quadratures exactly integrate $\int_{\widehat{D}} \frac{\partial u}{\partial \hat{x}_j} v$ for $v = G_{ij}^k$ for tensor product elements.

The key difference between simplicial and tensor product elements is the polynomial space in which the derivative lies. In contrast to the simplicial case, if $u \in Q^N$, $\frac{\partial u}{\partial \hat{x}_j} \notin Q^{N-1}$. Consider the three-dimensional case with $i = 1$, $u, v \in Q^N$. Then, $\frac{\partial u}{\partial \hat{x}_1} \in Q^{N-1, N, N}$, and differentiation reduces the polynomial degree in one coordinate but not others. As a result, $\frac{\partial u}{\partial \hat{x}_j} v \notin Q^{2N-1}$, and a tensor product quadrature of degree $(2N - 1)$ does not exactly integrate $\int_{\widehat{D}} \frac{\partial u}{\partial \hat{x}_j} v$ for general $v \in Q^N$.

We first consider the two dimensional case, and show that Assumption 1 holds with $v = G_{ij}^k$ for all $N_{\text{geo}} \leq N$. On quadrilateral elements, G_{ij}^k is

$$\begin{aligned} G_{11}^k &= \frac{\partial x_2}{\partial \hat{x}_2} \in Q^{N_{\text{geo}}, N_{\text{geo}}-1}, & G_{12}^k &= -\frac{\partial x_2}{\partial \hat{x}_1} \in Q^{N_{\text{geo}}-1, N_{\text{geo}}}, \\ G_{21}^k &= -\frac{\partial x_1}{\partial \hat{x}_2} \in Q^{N_{\text{geo}}, N_{\text{geo}}-1}, & G_{22}^k &= \frac{\partial x_1}{\partial \hat{x}_1} \in Q^{N_{\text{geo}}-1, N_{\text{geo}}}. \end{aligned}$$

647 Since $\frac{\partial u}{\partial \widehat{x}_1} \in Q^{N-1,N}$ and $\frac{\partial u}{\partial \widehat{x}_2} \in Q^{N,N-1}$

$$648 \quad \frac{\partial u}{\partial \widehat{x}_i} G_{ij}^k \in Q^{N+N_{\text{geo}}-1} \subset Q^{2N-1}, \quad \forall N_{\text{geo}} \leq N.$$

649 Thus, any quadrature rule of degree $(2N-1)$ exactly integrates $\int_{\widehat{D}} \frac{\partial u}{\partial \widehat{x}_i} v$ for $v = G_{ij}^k$.

650 We now consider the condition in Assumption 1 on the surface integrals $\int_{\partial \widehat{D}} u v n_i$
651 for $v = G_{ij}^k$. **Finish**

652 We now consider the three-dimensional case. Unlike quadrilateral elements, As-
653 sumption 1 does not hold for all $N_{\text{geo}} \leq N$ on general curved hexahedra. Expanding
654 out (6.1), one can show that $G_{ij}^k \in Q^{N_{\text{geo}}}$ on hexahedral elements. Thus, if $N_{\text{geo}} = N$
655 and $u \in Q^N$, $\frac{\partial u}{\partial \widehat{x}_1} G_{i1}^k \in Q^{N-1,N,N}$. **Finish.**

656 **6.3. Sufficient conditions on quadrature and N_{geo} to guarantee entropy**
657 **conservation.** For convenience, this section summarizes conditions on the accuracy
658 of volume and surface quadratures under which one can guarantee entropy conserva-
659 tion for a given degree N_{geo} of a geometric mapping.

660 **finish**

661 **7. A mortar-based implementation.** For hybrid meshes, the choice of sur-
662 face quadrature for an interface may vary depending on the neighboring elements.
663 Consider, for example, a mixed triangular-quadrilateral mesh where the quadrilater-
664 als utilize GLL quadrature on each face. While the skew-symmetric formulation allows
665 for the stable coupling of quadrilaterals and triangles using a GLL face quadrature,
666 one may wish to utilize more accurate Gauss face quadratures for triangle-triangle
667 interfaces. To accommodate arbitrary combinations of face quadrature, we introduce a
668 mortar-based implementation.

669 We assume that each element

670 **Finish - figure out notation for face interp!**

671 **8. Numerical experiments.** In this section, we present two-dimensional ex-
672 periments which verify the theoretical results presented.

673 **8.1. Triangular meshes.** Affine and curved triangles with GLL surface quadra-
674 ture (under-integrated).

675 **8.2. Quadrilateral meshes.** GLL quadrilaterals with Gauss surface quadra-
676 ture. Explain that for GLL quadratures, the decoupled SBP property doesn't hold
677 when Gauss points are used.

678 **8.3. Hybrid quadrilateral-triangular meshes.** It is not immediately clear
679 that the outward normals as defined by (21) will be equal and opposite across inter-
680 faces shared by elements of different types. Thus, we restrict ourselves to straight-
681 sided meshes in this case, and will investigate the effect of curvilinear mappings on
682 hybrid meshes in future work.

683 **9. Conclusions.**

684 **10. Acknowledgments.**

REFERENCES

- 686 [1] Gregor J Gassner. A skew-symmetric discontinuous Galerkin spectral element discretization and
687 its relation to SBP-SAT finite difference methods. *SIAM Journal on Scientific Computing*,
688 35(3):A1233–A1253, 2013.

- [2] Travis C Fisher and Mark H Carpenter. High-order entropy stable finite difference schemes for nonlinear conservation laws: Finite domains. *Journal of Computational Physics*, 252:518–557, 2013.
- [3] Mark H Carpenter, Travis C Fisher, Eric J Nielsen, and Steven H Frankel. Entropy Stable Spectral Collocation Schemes for the Navier–Stokes Equations: Discontinuous Interfaces. *SIAM Journal on Scientific Computing*, 36(5):B835–B867, 2014.
- [4] Jesse Chan, David C Del Rey Fernandez, and Mark Carpenter. Efficient entropy stable gauss collocation schemes. 2018. In preparation.
- [5] Jesse Chan. On discretely entropy conservative and entropy stable discontinuous Galerkin methods. *Journal of Computational Physics*, 362:346 – 374, 2018.
- [6] Jesse Chan and Lucas C Wilcox. Discretely entropy stable weight-adjusted discontinuous Galerkin methods on curvilinear meshes. *arXiv preprint arXiv:1805.10934*, 2018.
- [7] Zhijian J Wang, Krzysztof Fidkowski, Rémi Abgrall, Francesco Bassi, Doru Caraeni, Andrew Cary, Herman Deconinck, Ralf Hartmann, Koen Hillewaert, Hung T Huynh, et al. High-order CFD methods: current status and perspective. *International Journal for Numerical Methods in Fluids*, 72(8):811–845, 2013.
- [8] Mark Ainsworth. Dispersive and dissipative behaviour of high order discontinuous Galerkin finite element methods. *Journal of Computational Physics*, 198(1):106–130, 2004.
- [9] Miguel R Visbal and Datta V Gaitonde. High-order-accurate methods for complex unsteady subsonic flows. *AIAA journal*, 37(10):1231–1239, 1999.
- [10] Gregor J Gassner, Andrew R Winters, and David A Kopriva. Split form nodal discontinuous Galerkin schemes with summation-by-parts property for the compressible Euler equations. *Journal of Computational Physics*, 327:39–66, 2016.
- [11] Gregor J Gassner, Andrew R Winters, Florian J Hindenlang, and David A Kopriva. The BR1 scheme is stable for the compressible Navier–Stokes equations. *Journal of Scientific Computing*, pages 1–47, 2017.
- [12] Jared Crean, Jason E Hicken, David C Del Rey Fernández, David W Zingg, and Mark H Carpenter. High-Order, Entropy-Stable Discretizations of the Euler Equations for Complex Geometries. In *23rd AIAA Computational Fluid Dynamics Conference*. American Institute of Aeronautics and Astronautics, 2017.
- [13] Tianheng Chen and Chi-Wang Shu. Entropy stable high order discontinuous Galerkin methods with suitable quadrature rules for hyperbolic conservation laws. *Journal of Computational Physics*, 345:427–461, 2017.
- [14] Jared Crean, Jason E Hicken, David C Del Rey Fernández, David W Zingg, and Mark H Carpenter. Entropy-stable summation-by-parts discretization of the Euler equations on general curved elements. *Journal of Computational Physics*, 356:410–438, 2018.
- [15] Matteo Parsani, Mark H Carpenter, Travis C Fisher, and Eric J Nielsen. Entropy Stable Staggered Grid Discontinuous Spectral Collocation Methods of any Order for the Compressible Navier–Stokes Equations. *SIAM Journal on Scientific Computing*, 38(5):A3129–A3162, 2016.
- [16] Lucas Friedrich, Andrew R Winters, David C Fernández, Gregor J Gassner, Matteo Parsani, and Mark H Carpenter. An Entropy Stable h/p Non-Conforming Discontinuous Galerkin Method with the Summation-by-Parts Property. *arXiv preprint arXiv:1712.10234*, 2017.
- [17] Jeremy E Kozdon and Lucas C Wilcox. An energy stable approach for discretizing hyperbolic equations with nonconforming discontinuous Galerkin methods. *Journal of Scientific Computing*, 76(3):1742–1784, 2018.
- [18] Thomas JR Hughes, LP Franca, and M Mallet. A new finite element formulation for computational fluid dynamics: I. Symmetric forms of the compressible Euler and Navier-Stokes equations and the second law of thermodynamics. *Computer Methods in Applied Mechanics and Engineering*, 54(2):223–234, 1986.
- [19] Michael S Mock. Systems of conservation laws of mixed type. *Journal of Differential equations*, 37(1):70–88, 1980.
- [20] Amiram Harten. On the symmetric form of systems of conservation laws with entropy. *Journal of computational physics*, 49(1):151–164, 1983.
- [21] Constantine M Dafermos. *Hyperbolic conservation laws in continuum physics*. Springer, 2005.
- [22] Jesse Chan, Zheng Wang, Axel Modave, Jean-Francois Remacle, and T Warburton. GPU-accelerated discontinuous Galerkin methods on hybrid meshes. *Journal of Computational Physics*, 318:142–168, 2016.
- [23] Jason E Hicken, David C Del Rey Fernández, and David W Zingg. Multidimensional summation-by-parts operators: General theory and application to simplex elements. *SIAM Journal on Scientific Computing*, 38(4):A1935–A1958, 2016.
- [24] David C Del Rey Fernández, Pieter D Boom, and David W Zingg. A generalized framework for

- nodal first derivative summation-by-parts operators. *Journal of Computational Physics*, 266:214–239, 2014.
- [25] Hendrik Ranocha. Generalised summation-by-parts operators and variable coefficients. *Journal of Computational Physics*, 362:20 – 48, 2018.
- [26] H Xiao and Zydrunas Gimbutas. A numerical algorithm for the construction of efficient quadrature rules in two and higher dimensions. *Comput. Math. Appl.*, 59:663–676, 2010.
- [27] Claudio Canuto, M Yousuff Hussaini, Alfio Quarteroni, and Thomas A Zang. *Spectral Methods: Fundamentals in Single Domains*. Springer Science & Business Media, 2007.
- [28] Eitan Tadmor. The numerical viscosity of entropy stable schemes for systems of conservation laws. I. *Mathematics of Computation*, 49(179):91–103, 1987.
- [29] Cengke Shi and Chi-Wang Shu. On local conservation of numerical methods for conservation laws. *Computers and Fluids*, 2017.
- [30] PD Thomas and CK Lombard. Geometric conservation law and its application to flow computations on moving grids. *AIAA journal*, 17(10):1030–1037, 1979.
- [31] David A Kopriva. Metric identities and the discontinuous spectral element method on curvilinear meshes. *Journal of Scientific Computing*, 26(3):301–327, 2006.
- [32] Miguel R Visbal and Datta V Gaitonde. On the use of higher-order finite-difference schemes on curvilinear and deforming meshes. *Journal of Computational Physics*, 181(1):155–185, 2002.
- [33] Florian Hindenlang, Gregor J Gassner, Christoph Altmann, Andrea Beck, Marc Staudenmaier, and Claus-Dieter Munz. Explicit discontinuous Galerkin methods for unsteady problems. *Computers & Fluids*, 61:86–93, 2012.
- [34] T Warburton. An explicit construction of interpolation nodes on the simplex. *Journal of engineering mathematics*, 56(3):247–262, 2006.
- [35] Jesse Chan and T Warburton. A comparison of high order interpolation nodes for the pyramid. *SIAM Journal on Scientific Computing*, 37(5):A2151–A2170, 2015.
- [36] Andrew R Winters, Dominik Derigs, Gregor J Gassner, and Stefanie Walch. A uniquely defined entropy stable matrix dissipation operator for high Mach number ideal MHD and compressible Euler simulations. *Journal of Computational Physics*, 332:274–289, 2017.