

1 **SKIEW-SYMMETRIC ENTROPY STABLE DISCONTINUOUS** 2 **GALERKIN FORMULATIONS WITH APPLICATION TO HYBRID** 3 **MESHES**

4 JESSE CHAN

5 **1. Introduction.** High order methods for the simulation of time-dependent
6 compressible flow have the potential to achieve higher levels of accuracy at lower
7 costs compared to current low order schemes [1]. In addition to superior accuracy
8 and efficiency on modern computing architectures, the low numerical dispersion and
9 dissipation of high order methods [2] enables the accurate propagation of waves over
10 long distances and time scales. The same properties also make high order methods
11 attractive for the resolution of unsteady phenomena such as vorticular and turbulent
12 flows, which are sensitive to numerical dissipation [3, 1]. However, while high order
13 methods are often provably stable for wave problems, high order schemes for nonlinear
14 conservation laws have consistently been hampered by problems of instability.

15 When applied to nonlinear conservation laws, high order methods can experience
16 artificial growth and blow-up near under-resolved features such as shocks, turbulence,
17 or boundary layers. In practice, the application of high order methods to practical
18 problems requires shock capturing and stabilization techniques (such as artificial vis-
19 cosity) or solution regularization (such as filtering or limiting) to prevent solution
20 blow-up. The resulting schemes for nonlinear conservation laws walk a fine line be-
21 tween stability, robustness, and accuracy. Aggressive stabilization or regularization
22 can result in the loss of high order accuracy, while too little can result in instability
23 [1]. Moreover, it can be difficult to determine robust expressions for stabilization
24 paramaters, as parameters which work for one simulation can fail when applied to a
25 different physical regime or discretization setting.

26 These issues have motivated the introduction of high order *entropy stable* dis-
27 cretizations, which satisfy a semi-discrete entropy inequality while maintaining high
28 order accuracy in smooth regions. Proofs of continuous entropy inequalities rely on
29 the chain rule. In contrast, these discrete entropy inequalities account for the loss of
30 the chain rule due to effects such as quadrature errors, which are incurred when apply-
31 ing polynomially exact quadrature rules to nonlinear and rational integrands within
32 the DG formulation. These schemes were first introduced as high order collocation
33 methods on tensor product elements in [4, 5, 6, 7], and were extended to simplicial
34 elements in [8, 9, 10, 11]. These methods have also been extended to staggered grid
35 methods [12] and non-conforming meshes [13]. Entropy stable boundary conditions
36 have also been determined for the compressible Euler and Navier-Stokes equations
37 [14, 15].

38 **2. Entropy stability for systems of nonlinear conservation laws.** We
39 begin by reviewing the dissipation of entropy for a d -dimensional system of nonlinear
40 conservation laws on a domain Ω

$$41 \quad (1) \quad \frac{\partial \mathbf{u}}{\partial t} + \sum_{j=1}^d \frac{\partial \mathbf{f}_j(\mathbf{u})}{\partial x_j} = 0, \quad \mathbf{u} \in \mathbb{R}^n, \quad \mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n,$$

42 where \mathbf{u} are the conservative variables and $\mathbf{f}(\mathbf{u})$ is a vector-valued nonlinear flux
43 function. We are interested in nonlinear conservation laws for which a convex entropy
44 function $U(\mathbf{u})$ exists. For such systems, the *entropy variables* are an invertible map-
45 ping $\mathbf{v}(\mathbf{u}) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined as the derivative of the entropy function with respect

to the conservative variables

$$(2) \quad \mathbf{v}(\mathbf{u}) = \frac{\partial U}{\partial \mathbf{u}}.$$

Several widely used equations in fluid modeling (Burgers, shallow water, compressible Euler and Navier-Stokes equations) yield convex entropy functions $U(\mathbf{u})$ [16, 9]. Let $\partial\Omega$ be the boundary of Ω with outward unit normal \mathbf{n} . By multiplying the equation (1) with $\mathbf{v}(\mathbf{u})^T$, the solutions \mathbf{u} of (1) can be shown to satisfy an entropy inequality (3)

$$\int_{\Omega} \frac{\partial U(\mathbf{u})}{\partial t} dx + \int_{\partial\Omega} \sum_{j=1}^d (\mathbf{v}(\mathbf{u})^T \mathbf{f}_j(\mathbf{u}) - \psi_j(\mathbf{v}(\mathbf{u}))) n_j dx \leq 0, \quad \mathbf{n} = (n_1, \dots, n_d),$$

where $\psi_j(\mathbf{u})$ is a function referred to as the entropy potential.

The proof of (3) requires the use of the chain rule [17, 18, 19]. The instability-in-practice of high order schemes for (1) can be attributed in part to the fact that the discrete form of the equations do not satisfy the chain rule, and thus do not satisfy (3). As a result, discretizations of (1) do not typically possess an underlying statement of stability. For low order schemes, this can be offset in practice by the inherent numerical dissipation. However, because high order discretizations possess low numerical dissipation, the lack of an underlying discrete stability has been conflated with the idea that high order methods are inherently less stable than low order methods.

3. Approximation spaces. We define an approximation space using degree N polynomials on the reference element. In this work, we consider either simplicial reference elements (triangles and tetrahedra) or tensor product reference elements (quadrilaterals and hexahedra). The natural polynomial approximation space differs depending on the element type [20].

On a d -dimensional reference simplex, the natural polynomial space are total degree N polynomials

$$P^N(\widehat{D}) = \left\{ \widehat{x}_1^{i_1} \dots \widehat{x}_d^{i_d}, \quad \widehat{\mathbf{x}} \in \widehat{D}, \quad 0 \leq \sum_{k=1}^d i_k \leq N \right\}.$$

In contrast, the natural polynomial space on a d -dimensional tensor product element is

$$Q^N(\widehat{D}) = \left\{ \widehat{x}_1^{i_1} \dots \widehat{x}_d^{i_d}, \quad \widehat{\mathbf{x}} \in \widehat{D}, \quad 0 \leq i_k \leq N, \quad k = 1, \dots, d \right\}.$$

We take V^N to denote the natural approximation space on a given element. Furthermore, we denote the dimension of V^N as $N_p = \dim(V^N(\widehat{D}))$.

We also define trace spaces for each reference element. Let \widehat{f} be a face of the reference element \widehat{D} . The trace space $V^N(\widehat{f})$ is defined as the restrictions of functions in $V^N(\widehat{D})$ to \widehat{f}

$$V^N(\widehat{f}) = \left\{ u|_{\widehat{f}}, \quad u \in V^N(\widehat{D}) \right\}, \quad \widehat{f} \in \partial\widehat{D}.$$

For example, the trace space for degree N polynomials on a quadrilateral face \widehat{f} of the bi-unit hexahedral element $[-1, 1]^3$ is $Q^N(\widehat{f})$, while the trace space for degree

82 N polynomials on a triangular face f of the tetrahedron is $P^N(\hat{f})$. We denote the
 83 dimension of the trace space as $\dim(V^N(\hat{f})) = N_p^f$.

84 We similarly define the trace space for the surface $\partial\hat{D}$ of \hat{D}

$$85 \quad V^N(\partial\hat{D}) = \left\{ u|_{\partial\hat{D}}, \quad u \in V^N(\hat{D}) \right\}.$$

86 4. L^2 projection and lifting matrices and a decoupled SBP operator.

87 The schemes in [11] begin on the reference element \hat{D} by approximating the solution
 88 in a degree N polynomial basis $\{\phi_j(\mathbf{x})\}_{i=1}^{N_p}$. It is assumed that volume and surface
 89 quadrature rules (\mathbf{x}_i, w_i) , (\mathbf{x}_i^f, w_i^f) are provided for \hat{D} and the surface $\partial\hat{D}$ of \hat{D} ,
 90 respectively. We make the following assumption on the volume quadrature rule:

91 ASSUMPTION 1. *The volume quadrature rule integrates exactly all polynomials up*
 92 *to degree $(2N - 1)$.*

93 In addition to being related to the positive definiteness of the mass matrix, Assump-
 94 tion 1 ensures that, for any $u, v \in V^N$, the following integral is exactly approximated
 95 using quadrature

$$96 \quad \int_{\hat{D}} \frac{\partial u(\hat{\mathbf{x}})}{\partial \hat{x}_i} v(\hat{\mathbf{x}}) d\hat{\mathbf{x}} = \sum_{k=1}^{N_q} w_k \frac{\partial u(\hat{\mathbf{x}}_k)}{\partial \hat{x}_i} v(\hat{\mathbf{x}}_k).$$

97 In other words, evaluating the integral of the derivative of u and v using quadrature
 98 is exact because the integrand is a polynomial of degree $(2N - 1)$.

99 **TODO: Standardize hat notation for operators/normals: can refer to operators on**
 100 **reference element and do k superscript for physical elements, or do hats on reference**
 101 **element.**

102 Let $\mathbf{V}_q, \mathbf{V}_f$ denote interpolation matrices, and let \mathbf{D}^i be the differentiation matrix
 103 with respect to the i th coordinate such that

$$104 \quad (4) \quad (\mathbf{V}_q)_{ij} = \phi_j(\mathbf{x}_i), \quad (\mathbf{V}_f)_{ij} = \phi_j(\mathbf{x}_i^f), \quad \frac{\partial \phi_j(\mathbf{x})}{\partial x_i} = \sum_{k=1}^{N_p} (\mathbf{D}_{jk}^i) \phi_k(\mathbf{x}).$$

106 The interpolation matrices $\mathbf{V}_q, \mathbf{V}_f$ map basis coefficients to evaluations at volume and
 107 surface quadrature points respectively, while the differentiation matrix \mathbf{D}_i maps basis
 108 coefficients of a function to the basis coefficients of its derivative with respect to x_k .
 109 The interpolation matrices are used to assemble the mass matrix \mathbf{M} , the quadrature-
 110 based projection matrix \mathbf{P}_q , and lifting matrix \mathbf{L}_f

$$111 \quad (5) \quad \mathbf{M} = \mathbf{V}_q^T \mathbf{W} \mathbf{V}_q, \quad \mathbf{P}_q = \mathbf{M}^{-1} \mathbf{V}_q^T \mathbf{W}, \quad \mathbf{L}_f = \mathbf{M}^{-1} \mathbf{V}_f^T \mathbf{W}_f,$$

113 where \mathbf{W}, \mathbf{W}_f are diagonal matrices of volume and surface quadrature weights, res-
 114 pectively. The matrix \mathbf{P}_q is a quadrature-based discretization of the L^2 projection
 115 operator Π_N onto degree N polynomials.

116 Interpolation, differentiation, and L^2 projection matrices can be combined to con-
 117 struct finite difference operators. For example, the matrix $\mathbf{D}_q = \mathbf{V}_q \mathbf{D}^i \mathbf{P}_q$ maps func-
 118 tion values at quadrature points to approximate values of the derivative at quadrature
 119 points. By choosing specific quadrature rules, \mathbf{D}_q recovers high order summation-by-
 120 parts finite difference operators in [21, 22, 23] and certain operators in [24]. However,
 121 to address difficulties in designing efficient entropy stable interface terms for nonlinear
 122 conservation laws, the PI introduced in [11] a new “decoupled” summation by parts
 123 matrix which builds interface terms directly into the approximation of the derivative.

Let \mathbf{n} denote the outward unit normal \mathbf{n} on \widehat{D} , and let \mathbf{n}_i denote the vector containing values of the i th component at all surface quadrature points. The “decoupled” summation by parts operator \mathbf{D}_N^i is defined as the block matrix involving both volume and surface quadratures

$$(6) \quad \mathbf{D}_N^i = \begin{bmatrix} \mathbf{D}_q^i - \frac{1}{2} \mathbf{V}_q \mathbf{L}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q & \frac{1}{2} \mathbf{V}_q \mathbf{L}_f \text{diag}(\mathbf{n}_i) \\ -\frac{1}{2} \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q & \frac{1}{2} \text{diag}(\mathbf{n}_i) \end{bmatrix}.$$

When combined with projection and lifting matrices, \mathbf{D}_N^i produces a high order polynomial approximation of $f \frac{\partial g}{\partial x_i}$. Let f, g be differentiable functions, and let $\mathbf{f}_i = f(\mathbf{x}_i)$, $\mathbf{g}_i = g(\mathbf{x}_i)$ denote values of f, g at volume and surface quadrature points. Then,

$$(7) \quad f \frac{\partial g}{\partial x_i} \approx [\mathbf{P}_q \quad \mathbf{L}_f] \text{diag}(\mathbf{f}) \mathbf{D}_N^i \mathbf{g}.$$

The approximation can also be interpreted as solving a variational problem. Let n_i be the i th component of the unit normal on \widehat{D} . Then (7) is equivalent to finding $u(\mathbf{x}) \approx f \frac{\partial g}{\partial x_i}$ such that

$$(8) \quad \int_{\widehat{D}} u(\mathbf{x}) v(\mathbf{x}) = \int_{\widehat{D}} g \frac{\partial \Pi_N f}{\partial x_i} v + \int_{\partial \widehat{D}} (f - \Pi_N f) \frac{(gv + \Pi_N(gv))}{2} n_i, \quad \forall v \in V^N(\widehat{D}).$$

If the surface quadrature rule is exact for polynomials of degree $2N$, then the decoupled SBP operator \mathbf{D}_N^i also satisfies a summation by parts property.

$$(9) \quad \mathbf{Q}_N^i = \mathbf{W}_N \mathbf{D}_N^i, \quad \mathbf{W}_N = \begin{pmatrix} \mathbf{W} & \\ & \mathbf{W}_f \end{pmatrix},$$

$$(10) \quad \mathbf{Q}_N^i + (\mathbf{Q}_N^i)^T = \begin{pmatrix} \mathbf{0} & \\ & \mathbf{W}_f \text{diag}(\mathbf{n}_i) \end{pmatrix} = \mathbf{B}_N$$

By noting that $\mathbf{W} \mathbf{V}_q \mathbf{L}_q = \mathbf{P}_q^T \mathbf{V}_f^T \mathbf{W}_f$, the entries of \mathbf{Q}_N^i can be re-written as

$$(11) \quad \mathbf{Q}_i = \mathbf{W} \mathbf{V}_q \mathbf{D}_q^i \mathbf{P}_q,$$

$$(12) \quad \mathbf{Q}_N^i = \begin{bmatrix} \mathbf{Q}_i - \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q)^T \mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q & \frac{1}{2} (\mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q)^T \\ -\frac{1}{2} \mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q & \frac{1}{2} \mathbf{W}_f \text{diag}(\mathbf{n}_i) \end{bmatrix}.$$

from which the SBP property (8) is more apparent.

5. Skew-symmetric entropy conservative formulations.

5.1. Loss of the SBP property for hybrid and non-conforming meshes.

While the SBP property has been used to derive entropy stable schemes, the SBP property is difficult or impossible to enforce in certain discretization settings, such as hybrid and non-conforming meshes arises. This difficulty is a result of the choices of surface quadrature which naturally arise for these settings.

5.2. A variational SBP property.

It can be shown that the SBP property holds for \mathbf{D}_N^i if the surface quadrature is exact for polynomials of degree $2N$ [11]. The SBP property is lost if the surface quadrature is only exact for polynomial integrands of degree less than $2N$. However, one can still show that a variational version of the SBP property is satisfied.




Fig. 1: Examples of interface couplings which do not result in an SBP property and are not entropy stable. On the left, the surface quadrature is a $(N + 1)$ point GLL rule, and results in a loss of the SBP property on the triangle. On the right, the surface quadrature is a $(N + 1)$ point Gauss-Legendre rule, and results in a loss of the SBP property on the quadrilateral.




Fig. 2: Illustration of a 3D coupling between a GLL hexahedral element and a pyramid. The SBP property does not hold on the pyramid due to the use of GLL quadrature on the quadrilateral face.




Fig. 3: Illustration of two different entropy stable inter-element couplings for h non-conforming meshes. Each dashed red line indicates a dyadic flux computation required between two nodes.

$M \leq N$. If the surface quadrature is exact for polynomials of degree $N + M$, then for an arbitrary vector \mathbf{u}_q ,

$$\mathbf{v}_q^T \mathbf{Q}_i \mathbf{u}_q = \mathbf{v}_q^T \left((\mathbf{V}_f \mathbf{P}_q)^T \mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q - \mathbf{Q}_i^T \right) \mathbf{u}_q$$

where \mathbf{v} denotes the vector of values of v of at both volume and surface quadrature points.

Proof. Recall that $\mathbf{Q}_i = \mathbf{W} \mathbf{V}_q \mathbf{D}^i \mathbf{P}_q$. Since the volume quadrature is exact for degree $2N - 1$ polynomials, we have that

$$\mathbf{v}_q^T \mathbf{Q}_i \mathbf{u}_q = \mathbf{v}_q^T \mathbf{W} \mathbf{D}^i \mathbf{P}_q \mathbf{u}_q = \int_{\widehat{D}} \frac{\partial \Pi_N u}{\partial \widehat{x}_i} v = \int_{\partial \widehat{D}} (\Pi_N u) v \widehat{n}_i - \int_{\widehat{D}} (\Pi_N u) \frac{\partial v}{\partial \widehat{x}_i}.$$

Since the volume integrand $(\Pi_N u) \frac{\partial v}{\partial \widehat{x}_i} \in V^{2N-1}$, it is exactly integrated using the volume quadrature. Additionally, since $\Pi_N u \in V^N$ and $v \in V^M$ on the surface of \widehat{D} , the surface quadrature is exact for the surface integral and

$$\begin{aligned} & \int_{\partial \widehat{D}} (\Pi_N u) v \widehat{n}_i - \int_{\widehat{D}} (\Pi_N u) \frac{\partial v}{\partial \widehat{x}_i} = \\ & \mathbf{v}_f^T \mathbf{W}_f \text{diag}(\widehat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q \mathbf{u}_q - (\mathbf{V}_q \mathbf{D}^i \mathbf{P}_q \mathbf{v}_q)^T \mathbf{W} \mathbf{V}_q \mathbf{P}_q \mathbf{u}_q \\ & = \mathbf{v}_q^T (\mathbf{V}_f \mathbf{P}_q)^T \mathbf{W}_f \text{diag}(\widehat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q \mathbf{u}_q - \mathbf{v}_q^T \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{V}_q^T \mathbf{W} \mathbf{V}_q \mathbf{P}_q \mathbf{u}_q, \end{aligned}$$

where we have used that, since $v \in V^N$, $\mathbf{v}_f = \mathbf{V}_f \mathbf{P}_q \mathbf{v}_q$. The proof is completed by noting that

$$\mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{V}_q^T \mathbf{W} \mathbf{V}_q \mathbf{P}_q = \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{M} \mathbf{P}_q = \mathbf{P}_q^T (\mathbf{D}^i)^T \mathbf{V}_q^T \mathbf{W} = \mathbf{Q}_i^T. \quad \square$$

The variational SBP property also extends to the decoupled SBP operator. This property, along with the exact differentiation of constants, is necessary for the proof of entropy stability.

LEMMA 2. Let \mathbf{D}_N^i be a decoupled SBP operator on the reference element \widehat{D} , and let the surface quadrature be exact for polynomials of degree $N + M$ for some $M \leq N$. Suppose $v \in V^N$ and that $v|_{\partial \widehat{D}} \in P^M(\partial \widehat{D})$, Then,

$$\mathbf{v}^T \mathbf{Q}_N^i \mathbf{u} = \mathbf{v}^T \left(\mathbf{B}_N^i - (\mathbf{Q}_N^i)^T \right) \mathbf{u}.$$

where \mathbf{v} denotes the values of v of at both volume and surface quadrature points.

Proof. For convenience, let $\mathbf{u}_q, \mathbf{u}_f$ denote evaluations of u at volume and surface points, such that

$$\mathbf{u} = \begin{pmatrix} \mathbf{u}_q \\ \mathbf{u}_f \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} \mathbf{v}_q \\ \mathbf{v}_f \end{pmatrix}.$$

The proof of the variational summation by parts property uses the definition of \mathbf{Q}_N^i (10),

$$\begin{aligned} \mathbf{v}^T \mathbf{Q}_N^i \mathbf{u} &= \mathbf{v}_q^T \mathbf{Q}_i \mathbf{u}_q - \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q \mathbf{v}_q)^T \mathbf{W}_f \widehat{\mathbf{n}} (\mathbf{V}_f \mathbf{P}_q \mathbf{u}_q) + \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q \mathbf{v}_q)^T \mathbf{W}_f \widehat{\mathbf{n}} \mathbf{u}_f \\ &\quad - \frac{1}{2} \mathbf{v}_f^T \mathbf{W}_f \widehat{\mathbf{n}} (\mathbf{V}_f \mathbf{P}_q \mathbf{u}_q) + \frac{1}{2} \mathbf{v}_f^T \mathbf{W}_f \widehat{\mathbf{n}} \mathbf{u}_f. \end{aligned}$$

196 Applying Lemma 1 then yields

$$\begin{aligned}
197 \quad \mathbf{v}^T \mathbf{Q}_N^i \mathbf{u} &= -\mathbf{v}_q^T \mathbf{Q}_i^T \mathbf{u}_q + \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q \mathbf{v}_q)^T \mathbf{W}_f \hat{\mathbf{n}} (\mathbf{V}_f \mathbf{P}_q \mathbf{u}_q) + \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q \mathbf{v}_q)^T \mathbf{W}_f \hat{\mathbf{n}} \mathbf{u}_f \\
198 \quad &\quad - \frac{1}{2} \mathbf{v}_f^T \mathbf{W}_f \hat{\mathbf{n}} (\mathbf{V}_f \mathbf{P}_q \mathbf{u}_q) - \frac{1}{2} \mathbf{v}_f^T \mathbf{W}_f \hat{\mathbf{n}} \mathbf{u}_f + \mathbf{v}_f^T \mathbf{W}_f \hat{\mathbf{n}} \mathbf{u}_f \\
199 \quad &= \begin{pmatrix} \mathbf{v}_q \\ \mathbf{v}_f \end{pmatrix}^T \left(\begin{pmatrix} \mathbf{0} & \\ & \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) \end{pmatrix} \right. \\
200 \quad &\quad \left. + \begin{pmatrix} -\mathbf{Q}_i^T + \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q)^T \mathbf{W}_f \hat{\mathbf{n}} \mathbf{V}_f \mathbf{P}_q & -\frac{1}{2} (\mathbf{W}_f \hat{\mathbf{n}} \mathbf{V}_f \mathbf{P}_q)^T \\ \frac{1}{2} \mathbf{W}_f \hat{\mathbf{n}} \mathbf{V}_f \mathbf{P}_q & -\frac{1}{2} \mathbf{W}_f \hat{\mathbf{n}} \end{pmatrix} \right) \begin{pmatrix} \mathbf{u}_q \\ \mathbf{u}_f \end{pmatrix} \\
201 \quad &= \mathbf{v}^T \left(\mathbf{B}_N - (\mathbf{Q}_N^i)^T \right) \mathbf{u}. \quad \square
\end{aligned}$$

203 Using Lemma 2, we will determine the accuracy of the surface quadrature necessary
204 to ensure the proof of semi-discrete entropy stability is valid. This minimal degree
205 of surface quadrature accuracy will depend on the nature of the reference-to-physical
206 mapping (e.g. affine vs curved elements).

207 **COROLLARY 3.** *Let the surface quadrature be exact for polynomials of degree N .*
208 *Then,*

$$209 \quad \mathbf{1}^T \mathbf{Q}_N^i \mathbf{u} = \mathbf{1}^T \mathbf{B}_N \mathbf{u}, \quad \mathbf{Q}_N^i \mathbf{1} = \mathbf{0},$$

210 where \mathbf{u} is a vector of values of some function at volume and surface quadrature points.

211 *Proof.* The proof that $\mathbf{Q}_N^i \mathbf{1} = \mathbf{0}$ follows from the property that polynomials are
212 equal to their L^2 projection, and is identical to that of [11, 25]. The proof of the first
213 equality follows from $\mathbf{Q}_N^i \mathbf{1} = \mathbf{0}$ and Lemma 2 with $M = 0$

$$214 \quad \mathbf{1}^T (\mathbf{Q}_N^i) \mathbf{u} = \mathbf{1}^T \left(\mathbf{B}_N - (\mathbf{Q}_N^i)^T \right) \mathbf{u} = \mathbf{1}^T \mathbf{B}_N \mathbf{u}. \quad \square$$

215 **5.3. Entropy stability on affine meshes.** The high order methods in this
216 work ensure that (3) is satisfied discretely by avoiding the use of the chain rule in
217 the proof of entropy dissipation. These “entropy stable” schemes rely on two main
218 ingredients: an entropy stable numerical flux as defined by Tadmor [26] and a concept
219 referred to as “flux differencing”. Let $\mathbf{f}_S(\mathbf{u}_L, \mathbf{u}_R)$ be a numerical flux function which
220 is a function of “left” and “right” states $\mathbf{u}_L, \mathbf{u}_R$. The numerical flux \mathbf{f}_S is *entropy*
221 *conservative* if it satisfies the following three conditions:

$$\begin{aligned}
222 \quad (11) \quad &\mathbf{f}_S(\mathbf{u}, \mathbf{u}) = \mathbf{f}(\mathbf{u}), \quad (\text{consistency}) \\
223 \quad &\mathbf{f}_S(\mathbf{u}_L, \mathbf{u}_R) = \mathbf{f}_S(\mathbf{u}_R, \mathbf{u}_L), \quad (\text{symmetry}) \\
224 \quad &(\mathbf{v}_L - \mathbf{v}_R)^T \mathbf{f}_S(\mathbf{u}_L, \mathbf{u}_R) = \psi(\mathbf{u}_L) - \psi(\mathbf{u}_R), \quad (\text{conservation})
\end{aligned}$$

226 We begin by deriving a skew-symmetric formulation on the reference element \hat{D} ,
227 and showing it is entropy conservative. This formulation is then made entropy stable
228 by adding interface dissipation, and can be extended in a straightforward manner to
229 affine elements. The skew-symmetric formulation for (1) on \hat{D} is

$$230 \quad (12) \quad \frac{\partial \mathbf{u}}{\partial t} + \sum_{i=1}^d \begin{bmatrix} \mathbf{P}_q & \mathbf{L}_f \end{bmatrix} \left(\left(\mathbf{D}_N^i - \mathbf{W}_N^{-1} (\mathbf{Q}_N^i)^T \right) \circ \mathbf{F}_S \right) \mathbf{1} + \mathbf{L}_f \text{diag}(\hat{\mathbf{n}}) \mathbf{f}_i^* = \mathbf{0},$$

$$231 \quad (13) \quad (\mathbf{F}_S)_{ij} = \mathbf{f}_S(\tilde{\mathbf{u}}_i, \tilde{\mathbf{u}}_j), \quad 1 \leq i, j \leq N_q + N_q^f,$$

where \mathbf{f}^* is some numerical flux. Multiplying the formulation (13) by \mathbf{M} on both sides yields a weak form

$$(14) \quad \mathbf{M} \frac{\partial \mathbf{u}}{\partial t} + \sum_{i=1}^d \left[\begin{array}{c} \mathbf{V}_q \\ \mathbf{V}_f \end{array} \right]^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} + \mathbf{V}_f^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) \mathbf{f}_i^* = 0.$$

where the matrix $(\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T)$ possesses the following structure

$$(15) \quad (\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) = \begin{pmatrix} \mathbf{Q}_i - \mathbf{Q}_i^T & (\mathbf{V}_f \mathbf{P}_q)^T \mathbf{W}_f \text{diag}(\mathbf{n}_i) \\ \mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q & \mathbf{0} \end{pmatrix}.$$

We can now show that the skew-symmetric formulation is semi-discretely entropy conservative.

THEOREM 4. *Let the surface quadrature be exact for polynomials of degree N . Then, the formulation (13) is entropy conservative such that*

$$(16) \quad \mathbf{1}^T \mathbf{W} \frac{\partial U(\mathbf{u}_q)}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) (\psi_i(\tilde{\mathbf{u}}_f) - \tilde{\mathbf{v}}_f^T \mathbf{f}_i^*) = 0, \quad \mathbf{u}_q = \mathbf{V}_q \mathbf{u}.$$

244

Proof. Testing (14) by $\mathbf{v}_h = \mathbf{P}_q \mathbf{v}_q$ yields

$$(17) \quad \mathbf{v}_q^T \mathbf{W} \frac{\partial (\mathbf{V}_q \mathbf{u})}{\partial t} + \sum_{i=1}^d \tilde{\mathbf{v}}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} + \tilde{\mathbf{v}}_f^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) \mathbf{f}_i^* = 0.$$

One can show that [11]

$$(18) \quad \begin{aligned} \tilde{\mathbf{v}}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} &= \tilde{\mathbf{v}}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S) \mathbf{1} - \tilde{\mathbf{v}}^T ((\mathbf{Q}_N^i)^T \circ \mathbf{F}_S) \mathbf{1} \\ &= \tilde{\mathbf{v}}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S) \mathbf{1} - \mathbf{1}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S) \tilde{\mathbf{v}}. \end{aligned}$$

Where we have used that \mathbf{F}_S is symmetric and that the Hadamard product commutes. Applying the conservation condition on \mathbf{f}_S in (5.3) then yields

$$(19) \quad \begin{aligned} \tilde{\mathbf{v}}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S) \mathbf{1} - \mathbf{1}^T (\mathbf{Q}_N^i \circ \mathbf{F}_S) \tilde{\mathbf{v}} &= \sum_{jk} (\mathbf{Q}_N^i)_{jk} (\tilde{\mathbf{v}}_j - \tilde{\mathbf{v}}_k) \mathbf{f}_S(\tilde{\mathbf{u}}_j, \tilde{\mathbf{u}}_k) \\ &= \sum_{ij} (\mathbf{Q}_N^i)_{jk} (\psi_i(\tilde{\mathbf{u}}_j) - \psi_i(\tilde{\mathbf{u}}_k)) \\ &= \mathbf{1}^T (\mathbf{Q}_N^i) \psi_i(\tilde{\mathbf{u}}) - \psi_i(\tilde{\mathbf{u}})^T (\mathbf{Q}_N^i) \mathbf{1} \\ &= \mathbf{1}^T (\mathbf{Q}_N^i) \psi_i(\tilde{\mathbf{u}}) = \mathbf{1}^T \mathbf{B}_N^i \psi_i(\tilde{\mathbf{u}}) \\ &= \mathbf{1}^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) \psi_i(\tilde{\mathbf{u}}_f) \end{aligned}$$

where we have used Corollary 3 in the second to last equality. \square

REMARK. *The only result necessary to prove Theorem 4 is Corollary 3. Thus, when generalizing to curved elements, the proof of entropy stability for the skew-symmetric form will depend only on the curved version of Corollary 3.*

The skew symmetric formulation can also be shown to be locally conservative in the sense of [27], which is sufficient to show the numerical solution convergences to the weak solution under mesh refinement.

THEOREM 5. The formulation (13) is locally conservative such that

$$(16) \quad \mathbf{1}^T \mathbf{W} \frac{\partial (\mathbf{V}_q \mathbf{u})}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{f}_i^* = 0.$$

Proof. To show local conservation, we test (14) with 1. Let \mathbf{e} be the coefficients of 1 under the basis $\phi_j(\mathbf{x})$.

$$(17) \quad \mathbf{1}^T \mathbf{W} \mathbf{V}_q \frac{\partial \mathbf{u}}{\partial t} + \sum_{i=1}^d \mathbf{1}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} + \mathbf{1}^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}) \mathbf{f}_i^* = 0.$$

Because \mathbf{F}_S is symmetric and $(\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T)$ is skew-symmetric, the term

$$\left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right)$$

is a skew-symmetric matrix. Using that $\mathbf{A}, \mathbf{x}^T \mathbf{A} \mathbf{x} = 0$ for any skew symmetric matrix \mathbf{A} , the volume term vanishes

$$\mathbf{1}^T \left((\mathbf{Q}_N^i - (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} = 0. \quad \square$$

Next, we extend this formulation to affinely mapped elements. Derivatives with respect to physical coordinates on D^k are computed in terms of a change of variables formula and derivatives with respect to reference coordinates

$$\frac{\partial u}{\partial x_i} = \sum_{ij} \mathbf{G}_{ij}^k \frac{\partial u}{\partial \hat{x}_j}, \quad \mathbf{G}_{ij}^k = J^k \frac{\partial \hat{x}_j}{\partial x_i},$$

where J^k is the determinant of the Jacobian of the geometric mapping on the element D^k . For affine meshes, the scaled geometric terms \mathbf{G}_{ij}^k are constant over each element. It was shown in [11] that one can define a physical differentiation matrix as a linear combination of reference differentiation matrices. Let $\hat{\mathbf{D}}_N^i$ denote the decoupled SBP operator on the reference element \hat{D} , and let the decoupled SBP operators with respect to the physical coordinates on D^k be defined as

$$(18) \quad \mathbf{D}_N^i = \sum_{j=1}^d \mathbf{G}_{ij} \hat{\mathbf{D}}_N^j.$$

Then, an entropy conservative formulation can be given on D^k as follows:

$$(19) \quad \frac{\partial \mathbf{u}}{\partial t} + \sum_{i=1}^d \left[\begin{array}{cc} \mathbf{P}_q & \mathbf{L}_f \end{array} \right] \left((\mathbf{D}_N^i - \mathbf{W}_N^{-1} (\mathbf{Q}_N^i)^T) \circ \mathbf{F}_S \right) \mathbf{1} + \mathbf{L}_f \text{diag}(\mathbf{n}_i) \mathbf{f}_i^* = 0,$$

$$(\mathbf{F}_S)_{ij} = \mathbf{f}_S(\tilde{\mathbf{u}}_i, \tilde{\mathbf{u}}_j), \quad 1 \leq i, j \leq N_q + N_q^f,$$

$$\mathbf{f}^* = \mathbf{f}_S(\tilde{\mathbf{u}}_f^+, \tilde{\mathbf{u}}_f), \quad \text{on interior interfaces,}$$

where $\tilde{\mathbf{u}}_f^+$ denotes the face value of the entropy-projected conservative variables $\tilde{\mathbf{u}}_f$ on the neighboring element. The formulation can be made entropy stable by adding an appropriate penalization term, such as a Lax-Friedrichs or matrix-based dissipation [28, 9, 11].

5.4. Curvilinear meshes. On affine meshes, it is possible to show entropy stability of the skew-symmetric formulation (13) under a surface quadrature which is only exact for degree N polynomials. However, on curved meshes, a stronger surface quadrature rule is required to demonstrate entropy stability, where the strength of the surface rule depends on the order and approximation of geometric terms.

Let $\mathbf{J}^k \mathbf{G}_{ij}^k$ denote the vector of scaled geometric terms $\mathbf{J}^k \mathbf{G}_{ij}^k$ evaluated at volume and surface quadrature points. Decoupled SBP operators on a curved element D^k can be defined as in [25] by

$$(20) \quad D_N^i = \sum_{j=1}^d \text{diag}(\mathbf{G}_{ij}^k) \hat{D}_N^j + \hat{D}_N^j \text{diag}(\mathbf{G}_{ij}^k).$$

The skew-symmetric formulation (19) can be extended to curvilinear meshes using the definition of D_N^i in (20). However, additional assumptions must be satisfied in order to prove that the resulting formulation is entropy stable.

5.5. Discrete geometric conservation law and surface quadrature accuracy. Add description of how normals factor into this and note that they're computed based on \mathbf{G}_{ij} .

The first assumption which must be satisfied is the discrete geometric conservation law (GCL) [29, 30]. For curved elements, Lemma 2 and Corollary 3 do not necessarily hold at the discrete level. For example, expanding out the condition $\mathbf{Q}_N^i \mathbf{1} = \mathbf{0}$ in terms of (20) yields

$$(21) \quad D_N^i \mathbf{1} = \sum_{j=1}^d \text{diag}(\mathbf{G}_{ij}^k) \hat{D}_N^j \mathbf{1} + \hat{D}_N^j \text{diag}(\mathbf{G}_{ij}^k) \mathbf{1} = \sum_{j=1}^d \hat{D}_N^j (\mathbf{G}_{ij}^k) = 0,$$

where we have used that $\hat{D}_N^j \mathbf{1} = 0$. For degree N isoparametric mappings, the GCL is automatically satisfied in two dimensions due to the fact that the exact geometric terms $\mathbf{G}_{ij}^k \in P^{N-1}$ [30]. However, in three dimensions, the GCL is not automatically preserved due to the fact that $\mathbf{G}_{ij}^k \in P^{2N-2}$, and thus cannot be represented exactly using degree N polynomials. As a consequence, (21) must be enforced through an alternative construction of \mathbf{G}_{ij}^k .

A common approach is to rewrite the geometric terms as the curl of some quantity \mathbf{r}^i , but to interpolate \mathbf{r}^i before applying the curl [31, 30, 32]:

$$(22) \quad \mathbf{r}^i = \frac{\partial \mathbf{x}}{\partial \hat{x}_i} \times \mathbf{x}, \quad \begin{bmatrix} \mathbf{G}_{i1}^k \\ \mathbf{G}_{i2}^k \\ \mathbf{G}_{i3}^k \end{bmatrix} = -\frac{1}{2} \left(\frac{\partial I_R \mathbf{r}^k}{\partial \hat{x}_j} - \frac{\partial I_R \mathbf{r}^j}{\partial \hat{x}_k} \right),$$

where I_R denotes a degree R polynomial interpolation operator with appropriate interpolation nodes.¹ Because the geometric terms are computed by applying the curl, both \mathbf{G}_{ij}^k and $n_i J^f$ are approximated using degree $(R-1)$ polynomials.

Because D_N^i are now defined through (20), Corollary 3 and the proof of entropy stability may not hold for curved elements and must be modified. The introduction of curvilinear meshes will impose slightly different conditions on the accuracy of the surface quadrature, which are summarized in the following curved version of Corollary 3.

¹This interpolation step must be performed using interpolation points with an appropriate number of nodes on each boundary [25]. These include, for example, GLL nodes on tensor product elements, as well as Warp and Blend nodes on non-tensor product elements [33, 34].

LEMMA 6. Assume that the geometric terms $\mathbf{G}_{ij} \in V^N$ satisfy the discrete GCL (21), that $\mathbf{G}_{ij}|_{\partial\hat{D}} \hat{\mathbf{n}}_j \in V^R(\partial\hat{D})$, and that the surface quadrature is exact for polynomials of degree $N + R$. Then,

$$\mathbf{1}^T \mathbf{Q}_N^i \mathbf{u} = \mathbf{1}^T \mathbf{B}_N^i \mathbf{u}, \quad \mathbf{Q}_N^i \mathbf{1} = 0,$$

where

$$\mathbf{B}_N^i = \begin{pmatrix} \mathbf{0} \\ J_f^k \mathbf{W}_f \text{diag}(\mathbf{n}_i) \end{pmatrix}.$$

Proof. The second equality $\mathbf{Q}_N^i \mathbf{1} = 0$ is a consequence of the discrete GCL (21), and the proof is identical to that of [25]. For the first equality, if the surface quadrature is exact for polynomials of degree $2N$ on the trace space, then the stronger matrix SBP property $\mathbf{Q}_N^i = \mathbf{B}_N^i - (\mathbf{Q}_N^i)^T$ holds [25]. Combining this with $\mathbf{Q}_N^i \mathbf{1} = 0$ yields the desired result. We focus on the proof of the variational SBP property for surface quadrature which are exact for polynomials of degree $N + R < 2N$.

Expanding $\mathbf{1}^T \mathbf{Q}_N^i \mathbf{u}$ yields

$$\mathbf{1}^T \mathbf{Q}_N^i \mathbf{u} = \frac{1}{2} \sum_{j=1}^d \mathbf{1}^T \text{diag}(\mathbf{G}_{ij}) \hat{\mathbf{Q}}_N^j \mathbf{u} + \mathbf{1}^T \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}) \mathbf{u}.$$

For the latter term in the sum, Lemma 2 holds and

$$\begin{aligned} \sum_{j=1}^d \mathbf{1}^T \hat{\mathbf{Q}}_N^j \text{diag}(\mathbf{G}_{ij}) \mathbf{u} &= \sum_{j=1}^d \mathbf{1}^T \left(\hat{\mathbf{B}}_N^j - (\hat{\mathbf{Q}}_N^j)^T \right) \text{diag}(\mathbf{G}_{ij}) \mathbf{u} \\ &= \sum_{j=1}^d \mathbf{1}^T \hat{\mathbf{B}}_N^j \text{diag}(\mathbf{G}_{ij}) \mathbf{u}, \end{aligned}$$

where we have used that $\hat{\mathbf{Q}}_N^j \mathbf{1} = \mathbf{0}$. For the former term in the sum, we expand out

$$\begin{aligned} \sum_{j=1}^d \mathbf{1}^T \text{diag}(\mathbf{G}_{ij}) \hat{\mathbf{Q}}_N^j \mathbf{u} &= \sum_{j=1}^d \mathbf{G}_{ij}^T \hat{\mathbf{Q}}_N^j \mathbf{u} \\ &= \sum_{j=1}^d \begin{bmatrix} \mathbf{G}_{ij}^q \\ \mathbf{G}_{ij}^f \end{bmatrix}^T \begin{bmatrix} \hat{\mathbf{Q}}_i - \frac{1}{2} (\mathbf{V}_f \mathbf{P}_q)^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q & \frac{1}{2} (\mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q)^T \\ -\frac{1}{2} \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q & \frac{1}{2} \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \end{bmatrix} \begin{bmatrix} \mathbf{u}_q \\ \mathbf{u}_f \end{bmatrix}. \end{aligned}$$

where $\mathbf{G}_{ij}^q, \mathbf{G}_{ij}^f$ denote the values of \mathbf{G}_{ij} at volume and surface quadrature points.

Because $\mathbf{G}_{ij} \in V^N$, it is equal to its own L^2 projection, and the values at volume and surface quadrature points are related by $\mathbf{G}_{ij}^f = \mathbf{V}_f \mathbf{P}_q \mathbf{G}_{ij}^q$. We can use this to simplify (6)

$$\begin{aligned} \mathbf{1}^T \text{diag}(\mathbf{G}_{ij}) \hat{\mathbf{Q}}_N^j \mathbf{u} &= (\mathbf{G}_{ij}^q)^T \mathbf{Q}_i \mathbf{u}_q - \frac{1}{2} \left(\mathbf{V}_f \mathbf{P}_q \mathbf{G}_{ij}^q + \mathbf{G}_{ij}^f \right)^T \mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{V}_f \mathbf{P}_q \mathbf{u}_q \\ &\quad + \frac{1}{2} \left(\mathbf{V}_f \mathbf{P}_q \mathbf{G}_{ij}^q + \mathbf{G}_{ij}^f \right)^T \frac{1}{2} \mathbf{W}_f \text{diag}(\mathbf{n}_i) \mathbf{u}_f \\ &= (\mathbf{G}_{ij}^q)^T \left(\hat{\mathbf{Q}}_i - (\mathbf{V}_f \mathbf{P}_q \mathbf{G}_{ij}^q)^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q \right) \mathbf{u}_q \\ &\quad + \left(\mathbf{G}_{ij}^f \right)^T \mathbf{W}_f \text{diag}(\hat{\mathbf{n}}_i) \mathbf{u}_f. \end{aligned}$$

371 The first term can be further simplified using Lemma 1

$$372 \quad (\mathbf{G}_{ij}^q)^T \left(\widehat{\mathbf{Q}}_i - (\mathbf{V}_f \mathbf{P}_q \mathbf{G}_{ij}^q)^T \mathbf{W}_f \text{diag}(\widehat{\mathbf{n}}_i) \mathbf{V}_f \mathbf{P}_q \right) \mathbf{u}_q = (\mathbf{G}_{ij}^q)^T \left(\widehat{\mathbf{Q}}_i \right)^T \mathbf{u}_q.$$

373 The latter term can be combined with the surface term of (6) by noting that

$$374 \quad \left(\mathbf{G}_{ij}^f \right)^T \mathbf{W}_f \text{diag}(\widehat{\mathbf{n}}_i) \mathbf{u}_f = \mathbf{1}^T \widehat{\mathbf{B}}_N^j \text{diag}(\mathbf{G}_{ij}) \mathbf{u} = \mathbf{1}^T \text{diag}(\mathbf{G}_{ij}) \widehat{\mathbf{B}}_N^j \mathbf{u},$$

375 where we have used that $\widehat{\mathbf{B}}_N^j$ is diagonal and commutes with diagonal scaling by \mathbf{G}_{ij} .
 376 We can use polynomial exactness and the fact that the geometric terms satisfy the
 377 continuous GCL by construction [25] to show that $\sum_{j=1}^d \widehat{\mathbf{Q}}_j \mathbf{G}_{ij}^q = 0$. Combining this
 378 with (6) and expanding out $\widehat{\mathbf{B}}_N^j$ yields that

$$379 \quad \mathbf{1}^T \mathbf{Q}_N^i \mathbf{u} = \sum_{j=1}^d \left(\frac{1}{2} (\mathbf{G}_{ij}^q)^T \left(\widehat{\mathbf{Q}}_i \right)^T \mathbf{u}_q + \mathbf{1}^T \begin{pmatrix} 0 \\ \mathbf{W}_f \text{diag}(\mathbf{G}_{ij} \circ \widehat{\mathbf{n}}_j) \end{pmatrix} \mathbf{u} \right) \\
 380 \quad = \mathbf{1}^T \begin{pmatrix} 0 \\ \mathbf{W}_f \text{diag} \left(\sum_{j=1}^d \mathbf{G}_{ij} \circ \widehat{\mathbf{n}}_j \right) \end{pmatrix} \mathbf{u} = \mathbf{1}^T \mathbf{B}_N^i \mathbf{u}.$$

382 where we have used in the final step that $\sum_{j=1}^d \mathbf{G}_{ij} \widehat{\mathbf{n}}_j = \mathbf{n}_i \mathbf{J}_f^k$ [35, 25] □

383 Lemma 6 implies that the polynomial degree R of the surface geometric terms
 384 must be compatible with the accuracy of the surface quadrature. This, in turn, will
 385 depend on the way in which geometric terms are approximated. [Finish.](#)

386 **5.6. Simplicial vs tensor product elements.** It was shown in [25] that the
 387 approximation error is $O(h^{R+1})$, and that taking $R = (N + 1)$ approximates the
 388 geometric terms with an error of $O(h^{N+2})$.

389 [Need to note image of curl is in different spaces for simplicial vs tensor product](#)
 390 [elements. Influences proofs that geometric terms satisfy appropriate assumptions in](#)
 391 [those cases.](#)

392 [Kopriva trick to enforcing GCL still results in \$Q^N\$ normals 3D. May not satisfy](#)
 393 [requirement that \$\mathbf{1}^T \mathbf{Q}_N^i \psi_i\(\tilde{\mathbf{u}}\) = \mathbf{1}^T \mathbf{B}_N^i \psi_i\(\tilde{\mathbf{u}}\)\$ because of quadrature inaccuracy. Note](#)
 394 [- this still works for conforming hexes because of the full matrix SBP property.](#)

395 **6. Numerical experiments.** In this section, we present two-dimensional ex-
 396 periments on hybrid meshes consisting of quadrilaterals and triangles, as well as on
 397 non-conforming meshes of quadrilateral elements.

398 **6.1. Reduced surface quadrature.** Affine and curved triangles with GLL sur-
 399 face quadrature (under-integrated).

400 **6.2. Hybrid meshes.**

401 **6.3. Non-conforming meshes.**

402 REFERENCES

- 403 [1] Zhijian J Wang, Krzysztof Fidkowski, Rémi Abgrall, Francesco Bassi, Doru Caraeni, Andrew
 404 Cary, Herman Deconinck, Ralf Hartmann, Koen Hillewaert, Hung T Huynh, et al. High-
 405 order CFD methods: current status and perspective. *International Journal for Numerical*
 406 *Methods in Fluids*, 72(8):811–845, 2013.
 407 [2] Mark Ainsworth. Dispersive and dissipative behaviour of high order discontinuous Galerkin
 408 finite element methods. *Journal of Computational Physics*, 198(1):106–130, 2004.

- [3] Miguel R Visbal and Datta V Gaitonde. High-order-accurate methods for complex unsteady subsonic flows. *AIAA journal*, 37(10):1231–1239, 1999.
- [4] Travis C Fisher and Mark H Carpenter. High-order entropy stable finite difference schemes for nonlinear conservation laws: Finite domains. *Journal of Computational Physics*, 252:518–557, 2013.
- [5] Mark H Carpenter, Travis C Fisher, Eric J Nielsen, and Steven H Frankel. Entropy Stable Spectral Collocation Schemes for the Navier–Stokes Equations: Discontinuous Interfaces. *SIAM Journal on Scientific Computing*, 36(5):B835–B867, 2014.
- [6] Gregor J Gassner, Andrew R Winters, and David A Kopriva. Split form nodal discontinuous Galerkin schemes with summation-by-parts property for the compressible Euler equations. *Journal of Computational Physics*, 327:39–66, 2016.
- [7] Gregor J Gassner, Andrew R Winters, Florian J Hindenlang, and David A Kopriva. The BR1 Scheme is Stable for the Compressible Navier-Stokes Equations. *arXiv preprint arXiv:1704.03646*, 2017.
- [8] Jared Crean, Jason E Hicken, David C Del Rey Fernández, David W Zingg, and Mark H Carpenter. High-Order, Entropy-Stable Discretizations of the Euler Equations for Complex Geometries. In *23rd AIAA Computational Fluid Dynamics Conference*. American Institute of Aeronautics and Astronautics, 2017.
- [9] Tianheng Chen and Chi-Wang Shu. Entropy stable high order discontinuous Galerkin methods with suitable quadrature rules for hyperbolic conservation laws. *Journal of Computational Physics*, 345:427–461, 2017.
- [10] Jared Crean, Jason E Hicken, David C Del Rey Fernández, David W Zingg, and Mark H Carpenter. Entropy-stable summation-by-parts discretization of the Euler equations on general curved elements. *Journal of Computational Physics*, 356:410–438, 2018.
- [11] Jesse Chan. On discretely entropy conservative and entropy stable discontinuous Galerkin methods. *Journal of Computational Physics*, 362:346 – 374, 2018.
- [12] Matteo Parsani, Mark H Carpenter, Travis C Fisher, and Eric J Nielsen. Entropy Stable Staggered Grid Discontinuous Spectral Collocation Methods of any Order for the Compressible Navier–Stokes Equations. *SIAM Journal on Scientific Computing*, 38(5):A3129–A3162, 2016.
- [13] Lucas Friedrich, Andrew R Winters, David C Fernández, Gregor J Gassner, Matteo Parsani, and Mark H Carpenter. An Entropy Stable h/p Non-Conforming Discontinuous Galerkin Method with the Summation-by-Parts Property. *arXiv preprint arXiv:1712.10234*, 2017.
- [14] Matteo Parsani, Mark H Carpenter, and Eric J Nielsen. Entropy stable wall boundary conditions for the three-dimensional compressible navier–stokes equations. *Journal of Computational Physics*, 292:88–113, 2015.
- [15] Magnus Svärd, Mark H Carpenter, and Matteo Parsani. Entropy stability and the no-slip wall boundary condition. *SIAM Journal on Numerical Analysis*, 56(1):256–273, 2018.
- [16] Thomas JR Hughes, LP Franca, and M Mallet. A new finite element formulation for computational fluid dynamics: I. Symmetric forms of the compressible Euler and Navier-Stokes equations and the second law of thermodynamics. *Computer Methods in Applied Mechanics and Engineering*, 54(2):223–234, 1986.
- [17] Michael S Mock. Systems of conservation laws of mixed type. *Journal of Differential equations*, 37(1):70–88, 1980.
- [18] Amiram Harten. On the symmetric form of systems of conservation laws with entropy. *Journal of computational physics*, 49(1):151–164, 1983.
- [19] Constantine M Dafermos. *Hyperbolic conservation laws in continuum physics*. Springer, 2005.
- [20] Jesse Chan, Zheng Wang, Axel Modave, Jean-Francois Remacle, and T Warburton. GPU-accelerated discontinuous Galerkin methods on hybrid meshes. *Journal of Computational Physics*, 318:142–168, 2016.
- [21] Gregor J Gassner. A skew-symmetric discontinuous Galerkin spectral element discretization and its relation to SBP-SAT finite difference methods. *SIAM Journal on Scientific Computing*, 35(3):A1233–A1253, 2013.
- [22] David C Del Rey Fernández, Pieter D Boom, and David W Zingg. A generalized framework for nodal first derivative summation-by-parts operators. *Journal of Computational Physics*, 266:214–239, 2014.
- [23] Hendrik Ranocha. Generalised summation-by-parts operators and variable coefficients. *Journal of Computational Physics*, 362:20 – 48, 2018.
- [24] Jason E Hicken, David C Del Rey Fernández, and David W Zingg. Multidimensional summation-by-parts operators: General theory and application to simplex elements. *SIAM Journal on Scientific Computing*, 38(4):A1935–A1958, 2016.
- [25] Jesse Chan and Lucas Wilcox. On discretely entropy conservative and entropy stable discon-

- tinuous Galerkin methods: curvilinear meshes. In preparation.
- [26] Eitan Tadmor. The numerical viscosity of entropy stable schemes for systems of conservation laws. I. *Mathematics of Computation*, 49(179):91–103, 1987.
 - [27] Cengke Shi and Chi-Wang Shu. On local conservation of numerical methods for conservation laws. *Computers and Fluids*, 2017.
 - [28] Andrew R Winters, Dominik Derigs, Gregor J Gassner, and Stefanie Walch. A uniquely defined entropy stable matrix dissipation operator for high Mach number ideal MHD and compressible Euler simulations. *Journal of Computational Physics*, 332:274–289, 2017.
 - [29] PD Thomas and CK Lombard. Geometric conservation law and its application to flow computations on moving grids. *AIAA journal*, 17(10):1030–1037, 1979.
 - [30] David A Kopriva. Metric identities and the discontinuous spectral element method on curvilinear meshes. *Journal of Scientific Computing*, 26(3):301–327, 2006.
 - [31] Miguel R Visbal and Datta V Gaitonde. On the use of higher-order finite-difference schemes on curvilinear and deforming meshes. *Journal of Computational Physics*, 181(1):155–185, 2002.
 - [32] Florian Hindenlang, Gregor J Gassner, Christoph Altmann, Andrea Beck, Marc Staudenmaier, and Claus-Dieter Munz. Explicit discontinuous Galerkin methods for unsteady problems. *Computers & Fluids*, 61:86–93, 2012.
 - [33] T Warburton. An explicit construction of interpolation nodes on the simplex. *Journal of engineering mathematics*, 56(3):247–262, 2006.
 - [34] Jesse Chan and T Warburton. A comparison of high order interpolation nodes for the pyramid. *SIAM Journal on Scientific Computing*, 37(5):A2151–A2170, 2015.
 - [35] Philippe G Ciarlet. *The finite element method for elliptic problems*. Elsevier, 1978.